

**CONJOINT ADMINISTRATION OF MORPHOGENS AND ACE
INHIBITORS IN TREATMENT OF CHRONIC RENAL FAILURE**

Government Funding

5 A portion of this invention was made with U.S. government support under grant number DK59602, DK09976 from the National Institutes of Health (NIH), and under grant number DK20579 from the NIH supported Diabetes Research and Training Center (DRTC) of Washington University. The government may have certain rights in this invention.

10 **Related Applications**

 This application claims priority to the filing date of U.S. Provisional Application No. 60/406,431, filed August 28, 2002, entitled "Conjoint Administration of Morphogens and ACE Inhibitors In Treatment of Chronic Renal Failure," the entire teachings of which are hereby incorporated by reference.

15 **Background of the Invention**

 The mammalian renal system serves primary roles both in the removal of catabolic waste products from the bloodstream and in the maintenance of fluid and electrolyte balances in the body. Renal failures are, therefore, life-threatening conditions in which the build-up of catabolites and other toxins, and/or the
20 development of significant imbalances in electrolytes or fluids, may lead to the failure of other major organs systems and death. As a general matter, renal failure is classified as "acute" or "chronic." As detailed below, the differences between these two conditions are not merely a matter of severity or rapidity but, rather, reflect differences in etiology, prognosis, and treatment.

25 **Acute renal failure:**

 Acute renal failure is defined as an abrupt cessation or substantial reduction of renal function and, in as many as 90-95% of cases, may be secondary to trauma, surgery or another acute medical condition. Acute renal failure may be due to pre-

renal causes (e.g., decreased cardiac output, hypovolemia, altered vascular resistance) or to post-renal causes (e.g., obstructions or constrictions of the ureters, bladder or urethra) which do not directly involve the kidneys and which, if treated quickly, will not entail significant loss of nephrons or other damage to the kidneys.

5 Alternatively, acute renal failure may be due to intrinsic renal causes which involve a more direct insult or injury to the kidneys, and which may entail permanent damage to the nephrons or other kidney structures. Intrinsic causes of acute renal failure include but are not limited to infectious diseases (e.g., various bacterial, viral or parasitic infections), inflammatory diseases (e.g., glomerulonephritis, systemic
10 lupus erythematosus), ischemia (e.g., renal artery occlusion), toxic syndromes (e.g., heavy metal poisoning, side-effects of antimicrobial treatments or chemotherapy), and direct traumas.

The diagnosis and treatment of acute renal failure is as varied as its causes. In human patients, oliguria (urine output < 400 ml/day) or anuria (urine output < 50
15 ml/day) may be present in 50-70% of cases, BUN levels may climb 10-20 mg/dL/day or faster, plasma creatinine levels may climb 0.5-1.0 mg/dL/day, and metabolic acidosis is almost always present. If not treated, the electrolyte and fluid imbalances (e.g., hyperkalemia, acidosis, edema) associated with acute renal failure may lead to life-threatening arrhythmia, congestive heart failure, or multiple organ
20 system failures. Present therapies are typically directed at the underlying causes of the acute renal failure (e.g., pre-renal, post-renal, or infectious causes) and management of the complications. Due to the severity of acute renal failure, episodes rarely last longer than several weeks without mortality and are treated on an in-patient basis.

25 Chronic renal failure:

Chronic renal failure (CRF) is the progressive loss of kidney function. The kidneys attempt to compensate for renal damage by hyperfiltration (excessive straining of the blood) within the remaining functional nephrons (filtering units that consist of a glomerulus and corresponding tubule). Over time, hyperfiltration causes
30 further loss of function.

Chronic loss of function causes generalized wasting (shrinking in size) and progressive scarring within all parts of the kidneys. In time, overall scarring obscures the site of the initial damage. Yet, it is not until over 70% of the normal combined function of both kidneys is lost that most patients begin to experience
5 symptoms of kidney failure. Thus, chronic renal failure may be defined as a progressive, permanent and significant reduction of the glomerular filtration rate (GFR) due to a significant and continuing loss of nephrons.

Chronic renal failure typically begins from a point at which a chronic renal insufficiency (i.e., a permanent decrease in renal function of at least 50-60%) has
10 resulted from some insult to the renal tissues which has caused a significant loss of nephron units. The initial insult may or may not have been associated with an episode of acute renal failure. Irrespective of the nature of the initial insult, chronic renal failure manifests a "final common path" of signs and symptoms as nephrons are progressively lost and GFR progressively declines. This progressive
15 deterioration in renal function is slow, typically spanning many years or decades in human patients, but seemingly inevitable.

The early stage of chronic renal failure typically begins when GFR has been reduced to approximately one-third of normal (e.g., 30-40 ml/min for an average human adult). As a result of the significant nephron loss, and in an apparent
20 "attempt" to maintain the overall GFR with fewer nephrons, the average single nephron GFR (SNGFR) is increased by adaptations of the remaining nephrons. at both the structural and functional level. One structural manifestation of this adaptation, readily detectable by microscopic examination of biopsy samples, is a "compensatory hypertrophy" of both the glomeruli and the tubules of the kidney, a
25 process which literally increases the volume of filtrate which can be produced by each remaining nephron by literal enlargement of the glomeruli and tubules. Indeed, as a result of the hypertrophy or dilation of the collecting ducts, the urine of subjects with chronic renal failure often contains broad "casts," typically 2-6 times normal diameter, which aid in diagnosis and have also been referred to as "renal failure
30 casts." At the same time, there are functional changes in the remaining nephrons, such as decreased absorption or increased secretion of normally excreted solutes,

which may be responses to hormonal or paracrine changes elsewhere in the body (e.g., increasing levels of parathyroid hormone (PTH) in response to changes in serum levels of calcium and phosphate).

5 These adaptations in early stage chronic renal failure are not successful in completely restoring GFR or other parameters of renal function and, in fact, subject the remaining nephrons to increased risk of loss. For example, the increased SNGFR is associated with mechanical stresses on the glomerulus due to hyper-tension and hyper-perfusion. The loss of integrity of podocyte junctures leads to increased permeability of the glomerulus to macromolecules or "leakiness" of the glomerular capsule. Proliferative effects are also observed in mesangial, epithelial and
10 endothelial cells, as well as increases in the deposition of collagen and other matrix proteins. Sclerosis of both the glomeruli and tubules is another common symptom of the hypertrophied nephrons and the risk of coagulation in the glomerulus is increased. In particular, these adaptations of the remaining nephrons, by pushing the
15 SNGFR well beyond its normal level, actually decrease the capacity of the remaining nephrons to respond to acute changes in water, solute, or acid loads and, therefore, actually increase the probability of additional nephron loss.

 As chronic renal failure progresses, and GFR continues to decline to less than 10% of normal (e.g., 5-10 mL/min), the subject enters end-stage renal disease
20 (ESRD). During this phase, the inability of the remaining nephrons to adequately remove waste products from the blood, while retaining useful products and maintaining fluid and electrolyte balance, leads to a rapid decline in which many organ systems, and particularly the cardiovascular system, may begin to fail. For example, BUN and creatinine levels may be expected to rise and, at BUN levels of
25 60-100 mg/dL and serum creatinine levels of 8-12 mg/dL, a uremic syndrome will typically develop in which the kidneys can no longer remove the end products of nitrogen metabolism. At this point, renal failure will rapidly progress to death unless the subject receives renal replacement therapy (i.e., chronic hemodialysis, continuous peritoneal dialysis, or kidney transplantation).

30 Approximately 600 patients per million receive chronic dialysis each year in the United States, at an average cost approaching \$60,000-\$80,000 per patient per

year. Of the new cases of end-stage renal disease each year, approximately 28-33% are due to diabetic nephropathy (or diabetic glomerulopathy or diabetic renal hypertrophy), 24-29% are due to hypertensive nephrosclerosis (or hypertensive glomerulosclerosis), and 15-22% are due to glomerulonephritis.

- 5 The 5-year survival rate for all chronic dialysis patients is approximately 40%, but for patients over 65, the rate drops to approximately 20%.

Morphogens and Growth Factors:

A great many proteins have now been identified which appear to act as morphogenetic or growth factors, regulating cell proliferation or differentiation.

- 10 Typically these growth factors exert their effects on specific sets or subsets of cells or tissues. Thus, for example, epidermal growth factors, nerve growth factors, fibroblast growth factors, various hormones, and many other proteins inducing or inhibiting cell proliferation or differentiation have been identified and shown to affect some subgroup of cells or tissues.

- 15 One group of morphogenetic proteins, referred to herein as "morphogens," includes members of the family of osteogenic proteins/bone morphogenetic proteins (OP/BMPs) which were initially identified by their ability to induce ectopic, endochondral bone morphogenesis.

- Subsequent characterization of the nucleic acid and amino acid sequences of
20 the BMPs has shown them to be a subgroup of the TGF- β superfamily of growth factors. Members of this morphogen family have now been shown to include the mammalian osteogenic protein-1 (OP-1, also known as BMP-7), osteogenic protein-2 (OP-2), osteogenic protein-3 (OP-3), BMP-2 (also known as BMP-2A or CBMP-2A), BMP-3, BMP-4 (also known as BMP-2B or CBMP-2B), BMP-5, BMP-6, Vgr-
25 1, and GDF- 1, as well as the *Xenopus* homologue Vgl and the *Drosophila* homologues DPP and 60A. Members of this family encode secreted polypeptides, that share common structural features and that are similarly processed from pro-proteins to yield carboxy terminal mature proteins having a conserved pattern of cysteines. The active forms of these proteins are either disulfide-bonded
30 homodimers of a single family member, or heterodimers of two different members

(see, e.g., Massague (1990) *Annu. Rev. Cell Biol.* 6:597; Sampath, et al., (1990) *J. Biol. Chem.* 265:13198).

The members of the morphogen family of proteins are expressed in a variety of tissues during development. BMP-3 for, example, has been shown to be
5 expressed in developing human lung and kidney (Vukicevic et al. (1994) *J. Histochem. Cytochem.* 42:869-875), BMP-4 has been-shown to be expressed in the developing limbs, heart, facial processes and condensed mesenchyme associated with early whisker follicles in embryonic mice (Jones, et al. (1991) *Development* 111: 531-542), and OP-1 (BMP-7) has been shown immunohistochemically to be
10 associated with basement membranes in human embryos, including those of the developing lungs, pancreas, skin, and convoluted tubules of kidneys (Vukicevic, et al. (1994) *Biochem. Biophys. Res. Commun.* 198: 693-700). Some of the morphogens (e.g., OP-2 and BMP-2) were not detected in analyses of adult tissues, suggesting only an early developmental role for these morphogens (Ozkaynak, et al.
15 (1992) *J. Biol. Chem.* 267: 25220-25227). In contrast, high levels of murine OP-1 expression have been observed in adult mouse kidneys (Ozkaynak, et al. (1991) *Biochem. Biophys. Res. Commun.* 179: 116-123). This suggests a possible role for OP-1 synthesized in the kidney as a paracrine regulator of bone growth, and would be consistent with the role of the kidneys in both calcium regulation and bone
20 homeostasis.

A great variety of growth factors have been considered which may participate in the regulation of the growth and repair of renal tissues (reviewed in, e.g., Toback (1992) *Kidney Intl.* 41: 226-246). For example, EGF, TGF- α , TGF- β , IGF-I, IGMI, PDGF, FGF, Renin / Angiotensin II, IL-1 and OP-1 have all been
25 found to be expressed by various adult renal cells or tissues and to have effects on renal cell proliferation or differentiation (see, Toback (1992) *supra*, Ozkaynak, et al. (1991) *supra*). In addition, several of these have been found to be expressed in the developing kidney, including IGF-I, TGF- β and OP-1 (reviewed in, e.g., Bard, et al. (1994) *Mech. Development* 48: 3-11).

30 Interestingly, TGF- β has been shown in a murine metanephric organ culture system to retard overall growth and segmental differentiation of all segments of

developing nephrons except the thick ascending limb-early distal tubules (Avner and Sweeney (1990) *Pediatr. Nephrol.* 4: 372-377). In addition, TGF- β expression has been found to be increased in several models of renal disease, suggesting that TGF- β mediated increases in the synthesis of extracellular matrix components may be
5 involved in the etiology of diabetic nephropathy (or diabetic glomerulopathy or diabetic renal hypertrophy), renal fibrosis, glomerulosclerosis and glomerulonephritis, interstitial fibrosis, and hypertensive nephrosclerosis (Shankland, et al. (1994) *Kidney Intl.* 46: 430-442; Yamamoto, et al. (1994) *Kidney Intl.* 45:916-927; Yamamoto, et al. (1993) *PNAS* 90: 1814 Tamaki, et al. (1994)
10 *Kidney Intl.* 45:525-536; Border, et al. (1990) *Nature* 346: 371-374; Hamaguchi, et al. (1995) *Hypertension* 26: 199-207).

Also of interest is the fact that serum levels of human growth hormone (GH) are elevated in subjects with chronic renal failure (Wright et al. (1968) *Lancet* 2: 798; Samaan and Freeman (1970) *Metabolism* 19: 102). Recombinant GH has been
15 shown to help maintain protein balance in malnourished chronic renal failure patients, and to promote "catch-up" growth in children with chronic renal failure. It has been suggested that these effects are mediated by IGF-I (see, e.g., Kopple (1992) *Miner. Electrolyte Metab.* 18: 269-275). Although some studies have found that the administration of IGF-I increases renal plasma flow and GFR in chronic renal failure
20 patients (e.g., Guler, et al. (1989) *PNAS* 86:2868-2872; Hirschberg, et al. (1993) *Kidney Intl.* 43:387-397), other studies have found that this effect is merely transient (Miller, et al. (1994) *Kidney Intl.* 46: 201-207).

Thus, although some growth factors have been shown to be expressed in both developing and adult renal tissues, and although at least one has been shown to
25 increase renal function in the short term, none has yet been shown to be of therapeutic benefit in preventing, inhibiting, or delaying the progressive loss of renal function that characterizes chronic renal failure. A need remains, therefore, for treatments which will prevent the progressive loss of renal function which causes hundreds of thousand of patients to become dependent upon chronic dialysis, and
30 which results in the premature deaths of tens of thousands each year.

Summary of the Invention

The present invention is directed to methods of treatment, and pharmaceutical preparations for use in the treatment, of vertebrate subjects (preferably mammalian subjects) in, or at risk of, chronic renal failure, or at risk of the need for renal replacement therapy by using a combination of an Angiotensin-Converting Enzyme inhibitor (ACEI) and a morphogen. Suitable subjects include subjects already afflicted with chronic renal failure, or which have already received renal replacement therapy, as well as any subject reasonably expected to suffer a progressive loss of renal function associated with progressive loss of functioning nephron units. Whether a particular subject is at risk is a determination which may routinely be made by one of ordinary skill in the relevant medical or veterinary art. Subjects in, or at risk of, chronic renal failure, or at risk of the need for renal replacement therapy, include but are not limited to the following: subjects which may be regarded as afflicted with chronic renal failure, end-stage renal disease, chronic diabetic nephropathy, hypertensive nephrosclerosis, chronic glomerulonephritis, hereditary nephritis, and/or renal dysplasia; subjects having a biopsy indicating glomerular hypertrophy, tubular hypertrophy, chronic glomerulosclerosis, and/or chronic tubulointerstitial sclerosis; subjects having an ultrasound, MRI, CAT scan, or other non-invasive examination indicating renal fibrosis; subject shaving an unusual number of broad casts present in urinary sediment; subjects having a GFR which is chronically less than about 50%, and more particularly less than about 40%, 30% or 20%, of the expected GFR for the subject; human male subjects weighing at least about 50 kg and having a GFR which is chronically less than about 50 ml/min, and more particularly less than about 40 ml/min, 30 ml/min or 20 ml/min; human female subjects weighing at least about 40 kg and having a GFR which is chronically less than about 40 mL/min, and more particularly less than about 30 mL/min, 20 ml/min or 10 ml/min; subjects possessing a number of functional nephron units which is less than about 50%, and more particularly less than about 40%, 30% or 20%, of the number of functional nephron units possessed by a healthy but otherwise similar subject; subjects which have a single kidney; and subjects which are kidney transplant recipients.

The methods and compositions of this invention capitalize in part upon the discovery that certain morphogens of eukaryotic origin and inhibitors of ACE (Angiotensin-Converting Enzyme) may be used conjointly as therapeutic agents in the treatment of subjects at risk, as defined herein, of chronic renal failure or the
5 need for renal replacement therapy. Generally, these proteins are members of the osteogenic protein/bone morphogenetic protein (OP/BMP) family of proteins or analogs thereof. In a preferred embodiment, the ACE inhibitor is enalapril.

Thus, useful OP/BMP morphogens of the invention include polypeptides, or functional variants of polypeptides, comprising at least the C-terminal six- or seven-
10 cysteine domain of a mammalian protein selected from OP-1, OP-2, OP-3, BMP2, BMP3, BMP4, BMP5, BMP6, BMP9, and proteins which exhibit at least 70% or, more preferably, 75% or 80%, 85%, 90%, 95%, 99% amino acid sequence homology, or at least 50% identity, more preferably 55%, 60%, 65%, 70%, 80%, 90%, 99% or more identity, with the amino acid sequence of the seven-cysteine
15 domain of any of the morphogens described above, such as human OP-1; and which are (a) capable of inducing chondrogenesis in the Reddi-Sampath ectopic bone assay (Sampath and Reddi (1981), Proc. Natl. Acad. Sci. USA 78: 7599-7603) or a substantially equivalent assay, (b) capable of significantly preventing, inhibiting, delaying or alleviating the progressive loss of renal function in a standard animal
20 model of chronic renal failure, or (c) capable of causing a clinically significant improvement in a standard marker of renal function when administered to a mammal in, or at risk of, chronic renal failure. More generally speaking, the invention provides for the use of "morphogens" which are dimeric proteins that induce morphogenesis of one or more eukaryotic (e.g., mammalian) cells, tissues or organs.

25 Of particular interest herein are morphogens that induce morphogenesis at least of mammalian renal tissue, including formation of functional renal epithelium and, in particular, functional glomerular and tubular epithelium. Morphogens comprise a pair of polypeptides that, when folded, adopt a configuration suitable for the resulting dimeric protein to elicit morphogenetic responses in cells and tissues
30 displaying receptors specific for said morphogen. That is, morphogens generally induce all of the following biological functions in a morphogenically permissive

environment: stimulating proliferation of progenitor cells; stimulating the differentiation of progenitor cells; stimulating the proliferation of differentiated cells; and supporting the growth and maintenance of differentiated cells.

"Progenitor" cells are uncommitted cells that are competent to differentiate into one or more specific types of differentiated cells, depending on their genomic repertoire and the tissue specificity of the permissive environment in which morphogenesis is induced. Morphogens further can delay or mitigate the onset of senescence- or quiescence-associated loss of phenotype and/or tissue function. Morphogens still further can stimulate phenotypic expression of differentiated cells, including expression of metabolic and/or functional, e.g., secretory, properties thereof. In addition, morphogens can induce redifferentiation of committed cells under appropriate environmental conditions. As noted above, morphogens that induce proliferation and/or differentiation at least of mammalian renal tissue, and/or support the growth, maintenance and/or functional properties of mammalian nephrons, are of particular interest herein.

In preferred embodiments, the pair of morphogen polypeptides have amino acid sequence search comprising a sequence that shares a defined relationship with an amino acid sequence of a reference morphogen. Herein, preferred morphogen polypeptides share a defined relationship with a sequence present in morphogenically active human OP-1. However, anyone or more of the naturally occurring or biosynthetic sequences disclosed herein similarly could be used as a reference sequence. Preferred morphogen polypeptides share a defined relationship with at least the C-terminal six cysteine domain of human OP-1, residues 43-139 of SEQ ID NO: 1. Preferably, morphogen polypeptides share a defined relationship with at least the C-terminal seven cysteine domain of human OP-1, residues 38-139 of SEQ ID NO: 1. That is, preferred morphogen polypeptides in a dimeric protein with morphogenic activity each comprise a sequence that corresponds to a reference sequence or is functionally equivalent thereto.

Functionally equivalent sequences include functionally equivalent arrangements of cysteine residues disposed within the reference sequence, including amino acid insertions or deletions which alter the linear arrangement of these

cysteines, but do not materially impair their relationship in the folded structure of the dimeric morphogen protein, including their ability to form such intra- or inter-chain disulfide bonds as may be necessary for morphogenic activity.

Functionally equivalent sequences further include those wherein one or more amino acid residues differ from the corresponding residues of a reference morphogen sequence, e.g., the C-terminal seven cysteine domain (or "skeleton") of human OP-1, provided that this difference does not destroy morphogenic activity. Accordingly, conservative substitutions of corresponding amino acids in the reference sequence are preferred. Amino acid residues that are "conservative substitutions" for corresponding residues in a reference sequence are those that are physically or functionally similar to the corresponding reference residues, e.g., that have similar size, shape, electric charge, chemical properties including the ability to form covalent or hydrogen bonds, or the like. Particularly preferred conservative substitutions are those fulfilling the criteria defined for an "accepted point mutation" in Dayhoff et al. (1978), 5 Atlas of Protein Sequence and Structure, Suppl. 3, ch. 22 (pp. 354-352), *Nad. Biomed. Res. Found.*, Washington, D.C. (see below), the teachings of which are incorporated by reference herein.

In certain embodiments, a polypeptide suspected of being functionally equivalent to a reference morphogen polypeptide is aligned therewith using the method of Needleman, et al. (1970), *J. Mol. Biol.* 48: 443-453, implemented conveniently by computer programs such as the Align program or other improved successors / variants (DNASTar, Inc.). For example, the MegAlign program of the Lasergene 5.0 (DNASTar, Inc.) offers several multi-sequence alignment methods (J. Hein method, see Hein, J.J. (1990). "Unified approach to alignment and phylogenies." In *Methods in Enzymology*, Vol. 183: pp. 626-645; Clustal V method, see Higgins, D.G. and P.M. Sharp (1989). "Fast and sensitive multiple sequence alignments on a microcomputer." *CABIOS*, Vol. 5, No. 2: pp. 151-153; Clustal W method, see J.D. Thompson, et al. (1994). *Nucleic Acids Research*, Vol 22, pp. 4673-80), each with different algorithms, and each offers user opportunities to define parameters such as gaps. Specifically, gaps and insertions are arranged to achieve the highest degree of correlation between the amino acids of the two

sequences being compared, with user specified penalties – the so-called Gap Penalty (the amount deducted from the alignment score for each gap in the alignment. Gaps of different sizes carry the same penalty), and the Gap Length Penalty (the value deducted from the alignment score after first multiplying it by the length of gaps.

5 Longer gaps have a greater penalty than shorter gaps). The aligned sequences will then be used to calculate a percent identity (homology) between the candidate and reference sequences. Sequence homology between a section of the aligned sequences can also be generated. In a preferred embodiment, each amino acid of a gap or insertion counts as a mismatch for measuring % identity purpose.

10 Of particular interest herein are morphogens, which, when provided to the kidney of a mammal, induce or maintain the normal state of differentiation and growth of nephron units. Of still more particular interest herein are morphogens which, when administered to a mammal, prevent, inhibit or delay the development of compensatory hypertrophy, including glomerular hypertrophy and/or tubular
15 hypertrophy. Such morphogens can be used to treat a mammal in, or at risk of, chronic renal failure by preventing, inhibiting or delaying the progressive loss of functional nephron units and the consequent progressive loss of renal function.

The present invention alternatively can be practiced with methods and compositions comprising a morphogen stimulating agent or morphogen inducer in
20 lieu of a morphogen. A "morphogen inducer" is a compound that stimulates in vivo production, e.g., expression, of a therapeutically effective concentration of an endogenous morphogen in the body of a mammal sufficient to regenerate or maintain renal tissue and/or to inhibit additional loss thereof. Such compounds are understood to include substances which, when administered to a mammal, act on
25 cells of tissue(s) or organ(s) that normally are competent to produce and/or secrete a morphogen encoded within the genome of the mammal, and which cause the endogenous level of the morphogen in the mammal's body to be altered. Endogenous or administered morphogens can act as endocrine, paracrine or autocrine factors. That is, endogenous morphogens can be synthesized by the cells in
30 which morphogenetic responses are induced, by neighboring cells, or by cells of a distant tissue, in which circumstances the secreted endogenous morphogen is

transported to the site of morphogenesis, e.g., by the individual's bloodstream. In preferred embodiments, the agent stimulates expression and/or secretion of an endogenous morphogen so as to increase amounts thereof in renal tissues.

In still other embodiments, an agent which acts as an agonist of a morphogen
5 receptor may be administered instead of the morphogen itself. An "agonist" of a receptor means a compound which binds to the receptor and for which such binding has a similar functional result as binding of the natural, endogenous ligand of the receptor. That is, the compound must, upon interaction with the receptor, produce the same or substantially similar transmembrane and/or intracellular effects as the
10 endogenous ligand. Thus, an agonist of a morphogen receptor binds to the receptor and such binding has the same or a similar functional result as morphogen binding (e.g., induction of morphogenesis). The activity or potency of an agonist can be less than that of the natural ligand, in which case the agonist is said to be a "partial agonist," or it can be equal to or greater than that of the natural ligand, in which case
15 it is said to be a "full agonist." Thus, for example, a small peptide or other molecule which can mimic the activity of a morphogen in binding to and activating the morphogen's receptor may be employed as an equivalent of the morphogen. Preferably the agonist is a full agonist, but partial morphogen receptor agonists may also be advantageously employed. Methods of identifying such agonists are known
20 in the art and include assays for compounds which induce morphogen-mediated responses (e.g., induction of differentiation of metanephric mesenchyme, induction of endochondral bone formation, and the like). Such an agent may also be referred to as a morphogen "mimic," "mimetic," or "analog."

The OP/BMP morphogens, or the morphogen inducers / agonists of
25 morphogen receptors, or ACEI of the invention, may be administered by any route of administration which is compatible with the selected agent, and may be formulated with any pharmaceutically acceptable carrier appropriate to the route of administration. Preferred routes of administration are parenteral and, in particular, intravenous, intraperitoneal, and renal intracapsular. Treatments are also preferably
30 conducted over an extended period on an out-patient basis. Daily dosages of the morphogens are expected to be in the range of about 0.01-1000 $\mu\text{g/kg}$ body weight,

and more preferably about 10-300 µg/kg body weight, although precise dosages will vary depending upon the particular therapeutic agent employed and the particular subject's medical condition and history.

Finally, in yet further embodiments, renal cells may be implanted into the
5 kidney of a subject in, or at risk of, chronic renal failure, or at risk of needing renal replacement therapy, in order to serve as a source of morphogen and/or to provide a source of additional functional renal tissue. Preferably, the cells are induced to undergo metanephric differentiation by treatment with a morphogen (e.g., OP-1) either before or after implantation.

10 These cells may be renal mesenchymal progenitor cells, or renal mesenchymal progenitor cells which have been induced to undergo metanephric differentiation. The cells may be derived from a donor (e.g., a tissue-type matched donor, sibling, identical twin), or may be derived from a tissue culture (e.g., undifferentiated renal mesenchyme culture, fetal renal tissue culture), or may be
15 explanted from the subject and then be re-implanted after proliferation and/or differentiation.

The methods of the present invention are useful in preventing, inhibiting or delaying the progressive loss of functional nephron units, and the consequent progressive loss of renal function, which typify chronic renal failure. As such they
20 are of great value in preventing or delaying the need for chronic dialysis or renal replacement therapy in subjects with chronic renal insufficiency, or reducing the necessary frequency of chronic renal dialysis in subjects with end-stage renal disease. As such, they are useful in prolonging the lives, and in maintaining the quality of life, of subjects at risk of, or already afflicted with, chronic renal failure.

25 In a related aspect, the invention also contemplates conjoint administration of Angiotensin II Receptor Antagonists / Blockers (AIIRAs) with certain protein-based morphogens to subjects in, or at risk of, chronic renal failure, in order to reduce mortality and/or morbidity rates, and to prevent, inhibit, delay or alleviate the progressive loss of renal function which characterizes chronic renal failure.
30 Alternatively, or in addition, conjoint administration of angiotensin II receptor blockers with the morphogens of the present invention can prevent, inhibit or delay

the progressive loss of functional nephron units and the progressive decline in glomerular filtration rate (GFR) which slowly but inevitably leads to the need for renal replacement therapy (i.e., renal transplant or chronic dialysis) or death. In preferred embodiments, the therapeutic agents of the invention are members of the osteogenic protein / bone morphogenetic protein (OP/BMP) family within the TGF- β superfamily of proteins, and angiotensin II receptor blockers.

Brief Description of the Figures

Figure 1 The long term streptozotocin induced model of diabetic nephropathy. DM was induced at week 0 and the rats maintained as described in methods. (A) At 16 weeks, kidney weights had increased 1.8 fold in DM compared to normal (1.42 ± 0.02 versus 0.81 ± 0.02 g, $p < 0.01$). (B) GFR also increased 3.2 fold compare to normal (1.56 ± 0.27 versus 0.49 ± 0.04 ml/min/100g body wt, $p < 0.01$). After 16 weeks of vehicle treatment, kidney weights had not changed significantly (A), but the GFR was decreased 75% to even lower than normal at 32 weeks (0.34 ± 0.02 versus 0.55 ± 0.02) (B).

Figure 2 Effects of BMP-7 and enalapril treatments on DM induced renal hypertrophy. DM was induced at week 0. Treatment of BMP-7, or enalapril, or vehicle was began at week 16 and finished at week 32. At 16 weeks, kidney weight increased to 1.42 in DM compared to normal. After 16 weeks of vehicle treatment, the kidney weight was still elevated (1.44 ± 0.04 g). The kidney weights of the BMP-7 and enalapril treated groups were significantly decreased to 1.10 ± 0.03 in BMP-7 high dose group and 1.09 ± 0.04 in enalapril treated group, $p < 0.01$ compared to DM vehicle treated.

Figure 3 Effects of BMP-7 and enalapril treatments on GFR. In DM rats the GFR was increased 3.2-fold compare to normal at 16 weeks (1.56 ± 0.27 versus 0.49 ± 0.04 ml/min/100g body wt, $p < 0.01$), but by 32 weeks the GFR was decreased to lower than normal during vehicle treatment (0.34 ± 0.02 versus 0.55 ± 0.02). The BMP-7 and enalapril

treatments restored GFR to normal or slightly above normal. The GFR of the BMP-7 high dose animals were significantly greater than the GFR of DM group (vehicle treated) (0.70 ± 0.08 versus 0.34 ± 0.02 , $p < 0.05$). There was a dose-dependent organization of the GFR in the BMP-7 treated groups (0.59 ± 0.07 , 0.62 ± 0.09 , and 0.70 ± 0.08 , respectively).

Figure 4 Effects of BMP-7 and enalapril treatments on urine protein excretion in DM. DM was induced at week 0. Treatment of BMP-7, or Enalapril, or vehicle was began at week 16 and finished at week 32. Diabetic rats exhibited a pronounced increase in albumin excretion rate compared with nondiabetic rats at both 16 weeks (35.63 ± 13.35 versus 3.76 ± 0.39 mg/day) and 32 weeks (174.4 ± 52.50 versus 8.24 ± 1.28 , $p < 0.01$). This response was markedly reduced by BMP-7 and enalapril treatment ($p < 0.01$, DM versus BMP10; $p < 0.001$, DM versus BMP30, BMP100 and Enalapril). There was a dose-dependent inverse order in the levels of urinary protein in the BMP-7 treated groups from low to high dose (59.46 ± 21.84 , 33.02 ± 9.11 , and 14.27 ± 3.50 , respectively). The Enalapril treatment group had urinary protein excretion levels similar to the intermediate dose BMP-7 group.

Figure 5 Coronal sections of kidneys stained with periodic acid Schiff to highlight basement membranes and mesangial matrix. Panel A is a section of a kidney from a 16 week normal animal. B is a section of a kidney from a 16 week diabetic animal. Glomerular hypertrophy and early increases in mesangial matrix were present. There was evidence of glomerular (arrowhead) and tubular (arrow) basement membrane thickening. C is a section of a kidney from a 32 week diabetic vehicle-treated animal. One glomerulus is sclerotic and both are hypertrophied. There are segments of collapsed glomerular tuft with sparseness of normal cellular elements. D is a section of a kidney from an animal treated with BMP-7 30 μ g/kg body wt IV twice a

week. E is a section of a kidney from an animal treated with BMP-7 100 µg/kg body wt IV twice a week. F is a section of a kidney from an animal treated with Enalapril 100 mg/l in drinking water. All of the three BMP-7 dosages and enalapril treatment decreased glomerular sclerosis, but the enalapril treatment had more mesangial matrix accumulation.

Figure 6 Coronal sections of kidneys stained with Gomori's Trichrome for collagen. With this stain, collagen fibrils stain blue, whereas the cells stain red. A and B are sections from kidneys of two normal control animals maintained in the animal facility with the diabetic animals for 32 weeks. C is a section from a diabetic vehicle-treated animal. Arrow shows early interstitial matrix accumulation. The glomeruli are hypertrophic and one has segmental sclerosis. D, E, and F are sections of kidneys from animals treated with BMP-7 10, 30, and 100 µg/kg body wt. IV twice a week, respectively. Kidneys from the 10 µg/kg body wt. dose animals had greater mesangial matrix accumulation than the two higher doses which were very similar to normal except for residual glomerular hypertrophy.

Figure 7 Effects of BMP-7 and enalapril treatments on glomerular and interstitial areas in DM rats. (A) Glomerular area. Diabetic rats had a significantly larger glomerular area than normal control rats (1.28 ± 0.03 versus $0.90 \pm 0.02 \times 10^4 \mu\text{m}^2$, $p < 0.001$). All the treatment groups partially reversed the glomerular hypertrophy ($p < 0.001$). (B) Interstitial area. The cortical interstitial area was increased from $9.0 \pm 0.6\%$ sections in nondiabetic rat kidneys to $13.1 \pm 0.7\%$ in the DM rats. The BMP-7 high dose and enalapril treatment significantly decreased interstitial area ($10.7 \pm 0.3\%$, $p < 0.05$; $10.3 \pm 0.4\%$, $p < 0.01$, respectively) when compared to vehicle treated DM.

Figure 8 Effects of BMP-7 and enalapril treatments on glomerulosclerosis. Diabetic rats exhibited a significant increase in prevalence of sclerotic glomeruli compared with nondiabetic rats at 32 weeks (10.7

$\pm 4.0\%$ versus $0.7 \pm 0.2\%$, $p < 0.001$). This response was markedly reduced by BMP-7 and enalapril treatment. There was a dose-dependent ordering in the BMP-7 treated groups ($4.1 \pm 1.4\%$, $p < 0.05$; $3.5 \pm 0.8\%$, $p < 0.01$; and $2.1 \pm 0.3\%$, $p < 0.001$, respectively). The effect of high dose BMP-7 was significantly better than that of Enalapril.

Figure 9 Effects of Enalapril and BMP-7 therapies on systolic blood pressure of diabetic rats. Blood pressures were obtained by the artery cuff method. By 16 weeks the diabetic animals were significantly hypertensive. The hypertension was stable in vehicle treated rats until week 28 when systolic blood pressures began to increase again. Over two months Enalapril therapy restored blood pressure to normal. BMP-7 therapy did not affect blood pressure until the last four weeks of therapy.

Figure 10 Loss of BMP-7 renal expression in diabetes and restoration with therapy. (A) The 32-week vehicle treated diabetic kidneys had complete loss of BMP-7 message. Both BMP-7 and Enalapril therapy restored the normal distribution of BMP-7 expression. The levels of BMP-7 message in the BMP-7 and Enalapril treated kidneys appeared to be higher than the normal animals. (B) Glomerular expression of BMP-7 during BMP-7 therapy. Bright and dark field sections demonstrate significant BMP-7 expression in a glomerulus of a BMP-7 treated animal. These results were consistent in three separate experiments with different kidneys.

Figure 11 Induction of Wnt 4 expression by diabetes and effects of Enalapril and BMP-7 therapy. Wnt4 was not significantly expressed in normal kidneys. However, there was generalized renal expression of Wnt4 in vehicle treated diabetic rats. The expression was both in tubular epithelial cells and the glomeruli. BMP-7 therapy and Enalapril therapy had no effect on Wnt4 expression. The results were consistent in three separate experiments involving different kidneys.

- Figure 12 Schematic diagram of the five-sixths nephrectomy (5/6 NPX) Chronic Renal Failure (CRF) injury model.
- Figure 13 OP-1 did not dramatically lower the blood pressure level in the 5/6 nephrectomy model of Chronic Renal Failure.
- 5 Figure 14 OP-1 significantly reduced the proteinuria level in the 5/6 nephrectomy model of Chronic Renal Failure.
- Figure 15 In animals conjointly administered with morphogen (OP-1) and ACE inhibitor (enalapril), there is no additional benefit in reducing the blood pressure of nephrectomized animals to normal level as
10 compared to animals treated by the ACE inhibitor (enalapril) alone.
- Figure 16 Conjoint administration of morphogen (OP-1) and ACEI (enalapril) is more effective in reducing the proteinuria level in nephrectomized animals than ACEI treatment alone.
- Figure 17 Schematic diagram of the Unilateral Ureteral Obstruction (UUO) Renal Fibrosis Model.
15
- Figure 18 Morphogen OP-1 / BMP-7 Inhibits renal fibrosis in the Unilateral Ureteral Obstruction model.
- Figure 19 The mechanism of morphogen (OP-1)- induced renal protection is associated with prevention of tubular atrophy, an effect not shared
20 with ACEI enalapril.
- Figure 20 Both morphogen and ACEI improves renal function as measured by GFR. However, morphogen OP-1 is more efficacious than ACEI enalapril in improving the glomerular filtration rate as evidenced by the inulin clearance rate in the Unilateral Ureteral Obstruction model.
- 25 Figure 21 Morphogen (OP-1), but not ACE inhibitor, significantly reduced the loss of medullary tissue in the kidney in the Unilateral Ureteral Obstruction model.
- Figure 22 Schematic diagram of Streptozotocin-induced rat model of diabetic nephropathy.

- Figure 23 Morphogen (OP-1), but not ACEI enalapril, significantly increased the glomerular filtration rate (GFR) in the Streptozotocin-induced rat model of diabetic nephropathy.
- 5 Figure 24 Administration of morphogen (OP-1) alone, or ACEI enalapril alone, or conjoint administration of morphogen and ACEI significantly reduced the proteinuria level in the Streptozotocin-induced rat model of diabetic nephropathy.
- Figure 25 Schematic diagram of the Alloxan-induced rat model of diabetic nephropathy.
- 10 Figure 26 Morphogen OP-1 dramatically reduces the serum creatinine level, while ACEI elalapril reduces the serum creatinine level at a lesser degree in the Alloxan-induced rat model of diabetic nephropathy.
- Figure 27 Morphogen OP-1 and ACEI enalapril both reduce the preteinuria level. However, conjoint administration of morphogen and ACEI synergistically reduce the preteinuria level in the Alloxan-induced rat model of diabetic nephropathy.
- 15
- Figure 28 Comparison of percent amino acid sequence homology / similarity and percent identity within the C-terminal seven cysteine skeletons of various representative members of the TGF- β superfamily proteins, using OP-1 as the reference sequence. The percent homologies recited in the figure are based on aligning the sequences using the MegaAlign Program (DNASTar, Inc.).
- 20

Detailed Description of the Invention

I. Overview

- 25 The present invention depends, in part, upon the surprising discovery that conjoint administration of ACE inhibitors with certain protein-based morphogens to subjects in, or at risk of, chronic renal failure, can reduce mortality and/or morbidity rates, and prevent, inhibit, delay or alleviate the progressive loss of renal function which characterizes chronic renal failure. Alternatively, or in addition, conjoint

administration of ACE inhibitors with the morphogens of the present invention can prevent, inhibit or delay the progressive loss of functional nephron units and the progressive decline in glomerular filtration rate (GFR) which slowly but inevitably leads to the need for renal replacement therapy (i.e., renal transplant or chronic
5 dialysis) or death. In preferred embodiments, the therapeutic agents of the invention are members of the osteogenic protein / bone morphogenetic protein (OP/BMP) family within the TGF- β superfamily of proteins, and inhibitors of the ACE family of proteins.

The present invention also contemplates conjoint administration of
10 Angiotensin II Receptor Antagonists / Blockers (AIIIRAs) with certain protein-based morphogens to subjects in, or at risk of, chronic renal failure, in order to reduce mortality and/or morbidity rates, and to prevent, inhibit, delay or alleviate the progressive loss of renal function which characterizes chronic renal failure. Alternatively, or in addition, conjoint administration of angiotensin II receptor
15 blockers with the morphogens of the present invention can prevent, inhibit or delay the progressive loss of functional nephron units and the progressive decline in glomerular filtration rate (GFR) which slowly but inevitably leads to the need for renal replacement therapy (i.e., renal transplant or chronic dialysis) or death. In preferred embodiments, the therapeutic agents of the invention are members of the
20 osteogenic protein / bone morphogenetic protein (OP/BMP) family within the TGF- β superfamily of proteins, and angiotensin II receptor blockers.

II. Definitions

In order to more clearly and concisely point out the subject matter of the claimed invention, the following definitions are provided for specific terms used in
25 the following written description and appended claims.

OP/BMP morphogen. As used herein, the term "OP/BMP morphogen" means a polypeptide, or a functional variant of a polypeptide, comprising at least the C-terminal six- or seven-cysteine domain of a mammalian protein selected from the group consisting of OP-1, OP-2, OP-3, BMP2, BMP3, BMP4, BMP5, BMP6,
30 BMP9, and proteins which exhibit at least 65% or, more preferably, 70%, 75% 80%,

85%, 90%, or 95% amino acid sequence homology, or at least 50%, more preferably 55%, 60%, 70%, 80%, 90%, 99% or more identity, with the amino acid sequence of the seven-cysteine domain of any one of the morphogens described above, such as human OP-1 (SEQ ID NO: 1); and which is (a) capable of inducing chondrogenesis
5 in the Reddi-Sampath ectopic bone assay (Sampath and Reddi (1981), Proc. Natl. Acad. Sci. (USA) 78: 7599-7603) or a substantially equivalent assay, (b) capable of significantly preventing, inhibiting, delaying or alleviating the progressive loss of renal function in a standard animal model of chronic renal failure, or (c) capable of causing a clinically significant improvement in a standard marker of renal function
10 when administered to a mammal in, or at risk of, chronic renal failure.

As used herein, "amino acid sequence homology" or a percentage "homology" between two amino acid sequences is understood herein to include both amino acid sequence identity and conserved substitution. Thus, as used herein, a percentage "homology" between two amino acid sequences indicates the percentage
15 of amino acid residues which are identical or are conserved substitution between the sequences. "Conservative substitutions" of amino acids fulfill the criteria defined for an "accepted point mutation" in Dayhoff et al. (1978), Atlas of Protein Sequence and Structure Vol. 5 (Suppl. 3), pp. 354-352, Natl. Biomed. Res. Found., Washington, D.C. Thus, "conservative substitutions" are residues that are physically or
20 functionally similar to the corresponding reference residues, having similar size, shape, electric charge, and/or chemical properties such as the ability to form covalent or hydrogen bonds, or the like. Examples of preferred conservative substitutions include the substitution of one amino acid for another with similar characteristics, e.g., substitutions within the following groups: (a) Ser, Thr, Pro, Ala,
25 Gly; (b) Asn, Asp, Glu, Gln; (c) His, Arg, Lys; (d) Met, Ile, Leu, Val; (e) Phe, Tyr, Trp. In a most preferred embodiment, conservative substitutions include the substitution of one amino acid for another within the following groups: (a) glycine, alanine; (b) valine, isoleucine, leucine; (c) aspartic acid, glutamic acid; (d) asparagine, glutamine; (e) serine, threonine; (f) lysine, arginine, histidine; and (g)
30 phenylalanine, tyrosine. See Figure 84 of Dayhoff et al. (1978), Atlas of Protein Sequence and Structure Vol. 5 (Suppl. 3), pp. 354-352, Natl. Biomed. Res. Found., Washington, D.C. The term "conservative substitution" or "conservative variation"

also includes the use of a substituted amino acid in place of an unsubstituted parent amino acid in a given polypeptide chain, provided that the resulting substituted polypeptide chain also has therapeutic efficacy in the present invention.

As used herein, a therapeutic agent (morphogen and/or ACEI) of the
5 invention is said to have "therapeutic efficacy," and an amount of the agent is said to be "therapeutically effective," if administration of that amount of the agent is sufficient to cause a clinically significant improvement in a standard marker of renal function when administered to a mammalian subject (e.g., a human patient) in, or at risk of, chronic renal failure. Such markers of renal function are well known in the
10 medical literature and include, without being limited to, rates of increase in BUN levels, rates of increase in serum creatinine, static measurements of BUN, static measurements of serum creatinine, glomerular filtration rates (GFR), ratios of BUN/creatinine, serum concentrations of sodium (Na^+), urine/plasma ratios for creatinine, urine/plasma ratios for urea, urine osmolarity, daily urine output, and the
15 like (see, for example, Brenner and Lazarus(1994), in Harrison's Principles of internal Medicine, 13th edition, Isselbacher et al., eds., McGraw Hill Text, New York; Luke and Strom (1994), in Internal Medicine, 4th Edition, J.H. Stein, ed., Mosby-Year Book, Inc. St. Louis).

As used herein, "conjoint administration" means administration of two or
20 more agents to a subject of interest as part of a single therapeutic regimen. The administration(s) can be either simultaneous or sequential, i.e., administering one agent followed by administering of a second (and/or a third one, etc.) at a later time, as long as the agents administered co-exist in the subject being treated, or at least one agent will have the opportunity to act upon the same target tissues of other
25 agents while said target tissues are still under the influence of said other agents. In a preferred embodiment, agents to be administered can be included in a single pharmaceutical composition and administered together. In another preferred embodiment, the agents are administered simultaneously, including through separate routes. In yet another preferred embodiment, one or more agents are administered
30 continuously, while other agents are administered only at predetermined intervals (such as a single large dosage, or twice a week at smaller dosages, etc.). For

example, the morphogens can be administered three times a week through direct injection, while the ACE inhibitors can be continuously released by an implant.

The route of administration can be the same or different, depending on needs or suitable methods of administration for each agent. Any suitable route of administration may be employed for providing the patient with effective dosages of an ACEI and a morphogen. For example, oral, rectal, parenteral, transdermal, subcutaneous, intramuscular, inhalation and the like may be employed. Dosage forms include tablets, troches, dispersions, suspensions, solutions, capsules, patches and the like. In certain embodiments, the morphogen can be administered via direct injection, while the ACE inhibitors can be administered through drinking water.

As used herein, “prevent” or “prevention” means reducing the probability / risk of developing a condition in a subject (cell, tissue, organ, or organism, etc.), or delaying the onset of a condition in the subject, or to lessening the severity of one or more symptoms of a condition that may develop in the subject, or any combination thereof.

Filtration. The kidneys’ main function is to remove toxins (uremic wastes) that accumulate in the blood as a result of the body’s metabolism. The body continuously uses digested elements from foods and stored nutrients to perform normal bodily functions. The by-products of nutrient metabolism and cell function are filtered from the blood by the kidneys, which excrete (discharge) wastes as urine. Every day, approximately 200 liters of blood flow to the kidneys where 2 liters of waste are filtered out.

BUN and Creatinine. The concentration in the blood (blood level) of blood urea nitrogen (BUN), known as urea, and creatinine (Cr) can be measured by routine laboratory tests. BUN and creatinine levels indicate the general function of the kidneys. BUN is a metabolic by-product of protein-rich food such as meat, poultry, and certain vegetables. BUN is filtered out of the blood by the kidneys and excreted in the urine. Creatinine is continuously generated by normal cell metabolism within the muscles. Creatinine is also filtered out of the blood by the kidneys and excreted in the urine.

The amounts of BUN and creatinine in the blood are equal to the amount excreted by the kidneys. The blood levels of BUN and Cr remain unchanged unless there is sudden deterioration of renal (i.e., kidney) function. If the kidneys are *suddenly* unable to function, BUN and Cr increase daily. This condition is known as
5 *acute renal failure*. *Chronic renal failure* is a condition distinguished by a *gradual* increase in BUN and Cr over a long period of time.

Measurement of Kidney Function. When renal function decreases, blood levels of Cr and BUN increase because the kidneys are unable to clean the blood effectively. Factors not related to the kidneys also impact BUN and Cr levels.
10 Creatinine, in particular, is affected by age, sex, weight, and muscle mass.

Renal function is measured to evaluate the rate at which both kidneys are able to clean the blood. To measure renal function, a 24-hour urine sample must be collected. It is of importance that the 24-hour sample is complete (i.e., no urine is missing), so that true renal function will not be underestimated.

15 The amount of Cr in the urine sample is compared to the blood level of Cr. This figure is known as creatinine clearance (CrCl), the rate at which both kidneys clean the blood. The normal CrCl is about 90 to 130 milliliters per minute (mL/min). Many people gradually lose renal function as they age. Alternative renal function measurements rely on tables or formulas that take into consideration age, body
20 weight, sex, and blood creatinine.

Some health care facilities in the United States offer the Glofil-125 assay to evaluate renal function. Sodium iothalamate I-125 (a radiopharmaceutical) is injected into the skin, and blood and urine samples are obtained to determine renal function. The test is easy to perform, is more sensitive than blood creatinine
25 measurements, and provides results within 2 to 3 hours. Measurements of renal function determine the severity of kidney impairment. It is important to monitor renal function over time to document the rate of deterioration or improvement with treatment.

Glomerular Filtration Rate (GFR). The "glomerular filtration rate" or "GFR"
30 is proportional to the rate of clearance into urine of a plasma-borne substance which is not bound by serum proteins, is freely filtered across glomeruli, and is neither

secreted nor reabsorbed by the renal tubules. Thus, as used herein, GFR preferably is defined by the following equation.

$$\text{GFR} = U_{\text{conc}} \times V / P_{\text{conc}}$$

where U_{conc} is the urine concentration of the marker, P_{conc} is the plasma

5 concentration of the marker, and V is the urine flow rate in ml/min. Optionally, GFR is corrected for body surface area. Thus, the GFR values used herein may be regarded as being in units of ml/min/1.73m².

The preferred measure of GFR is the clearance of insulin but, because of the difficulty of measuring the concentrations of this substance, the clearance of
10 creatinine is typically used in clinical settings. For example, for an average size, healthy human male (70 kg, 20-40 yrs), a typical GFR measured by creatinine clearance is expected to be approximately 125 ml/min with plasma concentrations of creatinine of 0.7-1.5 mg/dL. For a comparable, average size woman, a typical GFR measured by creatinine clearance is expected to be approximately 115 ml/min with
15 creatinine levels of 0.5-1.3 mg/dL. During times of good health, human GFR values are relatively stable until about age 40, when GFR typically begins to decrease with age. For subjects surviving to age 85-90, GFR may be reduced to 50% of the comparable values at 40.

Expected Glomerular Filtration Rate (GFR_{exp}). An estimate of the "expected
20 GFR" or " GFR_{exp} " may be provided based upon considerations of a subject's age, weight, sex, body surface area, and degree of musculature, and the plasma concentration of some marker compound (e.g., creatinine) as determined by a blood test. Thus, as an example, an expected GFR or GFR_{exp} maybe estimated as:

$$\text{GFR}_{\text{exp}} = (140 - \text{age}) \times \text{weight (kg)} / [72 \times P_{\text{conc}} (\text{mg/dl})]$$

25 This estimate does not take into consideration such factors as surface area, degree of musculature, or percentage body fat. Nonetheless, using plasma creatinine levels as the marker, this formula has been employed for human males as an inexpensive means of estimating GFR. Because creatinine is produced by striated muscle, the expected GFR or GFR_{exp} of human female subjects is estimated by the

same equation multiplied by 0.85 to account for expected differences in muscle mass. (See Lemann, et al. (1990) Am. J. Kidney Dis. 16(3): 236).

Broad Cast. Microscopic examination of urinary sediment for the presence of formed elements is a standard procedure in urinalysis. Amongst the formed elements
5 which may be present in urine are cylindrical masses of agglutinated materials that typically represent a mold or "cast" of the lumen of a distal convoluted tubule or collecting tubule. In healthy human subjects, such casts typically have a diameter of 15-25 μm . In subjects with chronic renal failure, however, hypertrophy of the tubules may result in the presence of "broad casts" or "renal failure casts" which are
10 2-6 times the diameter of normal casts and often have a homogeneous waxy appearance. Thus, as used herein, a "broad cast" means a urinary sediment cast having a diameter of 2-6 times normal, or about 30-150 μm for human casts.

Chronic. As used herein with respect to clinical indications such as urinary casts, measured GFR, or other markers of renal function, "chronic" means persisting
15 for a period of at least three, and more preferably, at least six months. Thus, for example, a subject with a measured GFR chronically below 50% of GFR_{exp} is a subject in which the GFR has been measured and found to be below 50% of GFR_{exp} in at least two measurements separated by at least three, and more preferably, by at least six months, and for which there is no medically sound reason to believe that
20 GFR was substantially (e.g., 10%) higher during the intervening period.

Subjects in, or at risk of, chronic renal failure. As used herein, a subject is said to be in, or at risk of chronic renal failure, or at risk of the need for renal replacement therapy, if the subject is reasonably expected to suffer a progressive loss of renal function associated with progressive loss of functioning nephron units.
25 Whether a particular subject is in, or at risk of, chronic renal failure is a determination which may routinely be made by one of ordinary skill in the relevant medical or veterinary art. Subjects in, or at risk of, chronic renal failure, or at risk of the need for renal replacement therapy, include but are not limited to the following: subjects which may be regarded as afflicted with chronic renal failure, end-stage
30 renal disease, chronic diabetic nephropathy, hypertensive nephrosclerosis, chronic glomerulonephritis, hereditary nephritis, and/or renal dysplasia; subjects having a

biopsy indicating glomerular hypertrophy, tubular hypertrophy, chronic glomerulosclerosis, and/or chronic tubulointerstitial sclerosis; subjects having an ultrasound, NMR, CAT scan, or other non-invasive examination indicating renal fibrosis; subjects having an unusual number of broad casts present in urinary sediment; subjects having a GFR which is chronically less than about 50%, and more particularly less than about 40%, 30% or 20%, of the expected GFR for the subject; human male subjects weighing at least about 50 kg and having a GFR which is chronically less than about 50 ml/min, and more particularly less than about 40 ml/min, 30 ml/min or 20 ml/min; human female subjects weighing at least about 40 kg and having a GFR which is chronically less than about 40 ml/min, and more particularly less than about 30 ml/min, 20 ml/min or 10 ml/min; subjects possessing a number of functional nephron units which is less than about 50%, and more particularly less than about 40%, 30% or 20%, of the number of functional nephron units possessed by a healthy but otherwise similar subject; subjects which have a single kidney; and subjects which are kidney transplant recipients.

III. Description of the Preferred Embodiments

A. Therapeutic Agents

The morphogens of the present invention are naturally occurring proteins, or functional variants of naturally occurring proteins, in the osteogenic protein / bone morphogenetic protein (OP/BMP) family within the TGF- β superfamily of proteins.

The ACE inhibitors of the invention are generally small organic molecules. "Small molecule" as used herein, is meant to refer to a composition, which has a molecular weight of less than about 5 kD and most preferably less than about 2.5 kD. Small molecules can be nucleic acids, peptides, polypeptides, peptidomimetics, carbohydrates, lipids or other organic (carbon containing) or inorganic molecules.

(i) OP/BMP family of morphogens

The "OP/BMP family" of proteins forms a distinct subgroup, within the loose evolutionary grouping of sequence-related proteins known as the TGF- β superfamily. Members of this protein family comprise secreted polypeptides that share common structural features, and that are similarly processed from a pro-

protein to yield a carboxy-terminal mature protein. This family of proteins is also referred to as morphogens. As noted above, a protein is morphogenic as defined herein if it induces the developmental cascade of cellular and molecular events that culminate in the formation of new, organ-specific tissue.

5 It has been discovered that morphogens enhance survival of neurons and maintain neural pathways. As described herein, morphogens are capable of enhancing survival of neurons, stimulating neuronal CAM expression, maintaining the phenotypic expression of differentiated neurons, inducing the redifferentiation of transformed cells of neural origin, and stimulating axonal growth over breaks in
10 neural processes, particularly large gaps in axons. Morphogens also protect against tissue destruction associated with immunologically related nerve tissue damage. In addition, morphogens may be used as part of a method for monitoring the viability of nerve tissue in a mammal.

 In a preferred embodiment, a morphogen is a dimeric protein, each
15 polypeptide component of which has a sequence that corresponds to, or is functionally equivalent to, at least the conserved C-terminal six or seven cysteine skeleton of human OP-1, included in SEQ ID NO: 2, and/or which shares 70% amino acid sequence homology or 50% identity with OP-1 in this region. The morphogens are generally competent to induce a cascade of events including the
20 following, in a morphogenically permissive environment: stimulating proliferation of progenitor cells; stimulating the differentiation of progenitor cells; stimulating the proliferation of differentiated cells; and supporting the growth and maintenance of differentiated cells. Under appropriate conditions, morphogens are also competent to induce redifferentiation of cells that have undergone abnormal differentiation.
25 Details of how the morphogens useful in this invention were identified, as well as a description on how to make, use and test them for morphogenic activity are disclosed in numerous publications, including U.S. Patent Nos. 5,011,691 and 5,266,683, and the international patent application publications WO 92/15323; WO 93/04692; and WO 94/03200, each of which are incorporated by reference herein.
30 As disclosed therein, the morphogens can be purified from naturally sourced material or recombinantly produced from prokaryotic or eukaryotic host cells, using

the genetic sequences disclosed therein. Alternatively, novel morphogenic sequences can be identified following the procedures disclosed therein.

The naturally occurring morphogens share substantial amino acid sequence homology in their C-terminal sequences (sharing, *e.g.*, a six or seven cysteine skeleton sequence). Typically, a naturally occurring morphogen is translated as a precursor, having an N-terminal signal peptide sequence, typically less than about 35 residues in length, followed by a "pro" domain that is cleaved to yield the mature polypeptide, which includes the biologically active C-terminal skeleton sequence. The signal peptide is cleaved rapidly upon translation, at a cleavage site that can be predicted in a given sequence using the method of Von Heijne, *Nucleic Acids Research* 14: 4683-4691 (1986). The "pro" domain is variable both in sequence and in length, ranging from approximately 200 to over 400 residues. The pro domain is cleaved to yield the "mature" C-terminal domain of approximately 115-180 residues, which includes the conserved six- or seven-cysteine C-terminal domain of 97-106 residues. The pro polypeptide typically is about three times larger than the fully processed, mature C-terminal polypeptide. Under native conditions, the protein is secreted as a mature dimer and the cleaved pro polypeptide is thought to remain associated therewith to form a protein complex, presumably to improve the solubility of the mature dimeric protein. The complexed form of a morphogen is generally observed to be more soluble than the mature form under physiological conditions.

As used herein, the "pro form" of an OP/BMP family member refers to a protein comprising a folded pair of polypeptides, each comprising a pro domain in either covalent or noncovalent association with the mature domains of the OP/BMP polypeptide. The pro form appears to be the primary form secreted from cultured mammalian cells. The "mature form" of the protein refers to mature C-terminal domain which is not associated, either covalently or noncovalently, with the pro domain. Any preparation of OP-1 is considered to contain mature form when the amount of pro domain in the preparation is no more than 5% of the amount of "mature" C-terminal domain.

Natural-sourced morphogenic protein in its mature, native form, is typically a glycosylated dimer, having an apparent molecular weight of about 30-36 kDa as determined by SDS-PAGE. When reduced, the 30 kDa protein gives rise to two glycosylated polypeptide subunits having apparent molecular weights in the range of about 16 kDa and about 18 kDa. The unglycosylated dimeric protein, which also has morphogenic activity, typically has an apparent molecular weight in the range of about 27 kDa. When reduced, the 27 kDa protein gives rise to two unglycosylated polypeptides having molecular weights typically in the range of about 14 kDa to about 16 kDa.

OP/BMP family members useful herein include any of the known naturally occurring native proteins including allelic, phylogenetic counterpart and other variants thereof, whether naturally sourced or biosynthetically produced (e.g., including "muteins" or "mutant proteins"), as well as new, active members of the OP/BMP family of proteins. Particularly useful sequences include those comprising the C-terminal seven cysteine domains of mammalian, preferably human, OP-1, OP-2, OP-3, BMP2, BMP3, BMP4, BMP5, BMP6, BMP8 and BMP9. Other proteins useful in the practice of the invention include active forms of DPP, Vg1, Vgr-1, 60A, GDF-1, GDF-3, GDF-5, GDF-6, GDF-7, BMP10, BMP11; BMP13, BMP15, UNIVIN, NODAL, SCREW, ADMP or NURAL and amino acid sequence variants thereof. In one preferred embodiment, the morphogens of the invention are selected from any one of OP-1, OP-2, OP-3, BMP2, BMP3, BMP4, BMP5, BMP6, and BMP9.

In preferred embodiments, each of the polypeptide subunits of a dimeric morphogenic protein as defined herein comprises an amino acid sequence sharing a defined relationship with an amino acid sequence of a reference morphogen. In one embodiment, preferred morphogenic polypeptide chains share a defined relationship with a sequence present in morphogenically active full-length human OP-1, SEQ ID NO: 3. However, any one or more of the naturally occurring or biosynthetic morphogenic proteins disclosed herein similarly could be used as a reference sequence. Preferred morphogenic polypeptide chains share a defined relationship with at least the C-terminal six cysteine skeleton of human OP-1, residues 335-431

of SEQ ID NO: 3 (or residues 38-139 of SEQ ID NO: 1). Preferably, morphogenic proteins share a defined relationship with at least the C-terminal seven cysteine skeleton of human OP-1, residues 330-431 of SEQ ID NO: 3 (or residues 38-139 of SEQ ID NO: 1).

5 Functionally equivalent sequences include functionally equivalent arrangements of cysteine residues disposed within the reference sequence, including amino acid insertions or deletions which alter the linear arrangement of these cysteines, but do not materially impair their relationship in the folded structure of the dimeric morphogen protein, including their ability to form such intra- or inter-
10 chain disulfide bonds as may be necessary for morphogenic activity. For example naturally occurring morphogens have been described in which at least one internal deletion (of one residue; BMP2) or insertion (of four residues; GDF-1) is present but does not abrogate biological activity. Functionally equivalent sequences further include those wherein one or more amino acid residues differ from the
15 corresponding residue of a reference sequence, *e.g.*, the C-terminal seven cysteine skeleton of human OP-1, provided that this difference does not destroy tissue morphogenic activity. Accordingly, conservative substitutions of corresponding amino acids in the reference sequence are preferred. Amino acid residues that are "conservative substitutions" for corresponding residues in a reference sequence are
20 those that are physically or functionally similar to the corresponding reference residues, *e.g.*, that have similar size, shape, electric charge, chemical properties including the ability to form covalent or hydrogen bonds, or the like. Particularly preferred conservative substitutions are those fulfilling the criteria defined for an accepted point mutation in Dayhoff, *et al.*, 5 ATLAS OF PROTEIN SEQUENCE AND
25 STRUCTURE, Suppl. 3, ch. 22 pp. 354-352 (1978), Natl. Biomed. Res. Found., Washington, D.C. (*supra*), the teachings of which are incorporated by reference herein. The term "conservative substitution" also includes the use of a synthetic or derivatized amino acid in place of the corresponding natural parent amino acid, provided that antibodies raised to the resulting variant polypeptide also immunoreact
30 with the corresponding naturally sourced morphogen polypeptide.

Among the morphogens useful in this invention are proteins originally identified as osteogenic proteins, such as the OP-1, OP-2 and CBMP2 proteins, as well as amino acid sequence-related proteins such as DPP (from *Drosophila*), Vgl (from *Xenopus*), Vgr-1 (from mouse, see U.S. 5,011,691 to Oppermann et al.), GDF-1 (from mouse, see Lee (1991) *PNAS* 88: 4250-4254), all of which are presented in Table II), and the recently identified 60A protein (from *Drosophila*, see Wharton et al. (1991) *PNAS* 88: 9214-9218). As mentioned before, the members of this family, which include members of the TGF- β super-family of proteins, share substantial amino acid sequence homology in their C-terminal regions. The proteins are translated as a precursor, having an N-terminal signal peptide sequence, typically less than about 30 residues, followed by a "pro" domain that is cleaved to yield the mature sequence. The signal peptide is cleaved rapidly upon translation, at a cleavage site that can be predicted in a given sequence using the method of Von Heijne ((1986) *Nucleic Acids Research* 14:4683-4691.)

Table I below summarizes various naturally occurring members of the OP/BMP family identified to date, including their nomenclature as used herein, their Sequence listing references, and publication sources for the amino acid sequences for the full length proteins not included in the Sequence listing. Each of the generic terms set forth in Table I is intended and should be understood to embrace the therapeutic effective proteins expressed from nucleic acids encoding the identified sequence mentioned below and set forth in the Sequence listing, or an active fragment or precursor thereof, or a functional equivalent thereof such as a naturally occurring or biosynthetic variant. Naturally occurring variants include allelic variant forms isolated from other individuals of a single biological species, as well as species variants (homologous) isolated from phylogenetically distinct biological species.

Table I Exemplary Morphogens

30	"OP-1"	Refers generically to the group of morphogenically active proteins expressed from part or all of a DNA sequence encoding OP-1 protein, including allelic and species variants thereof, e.g., human OP-1 ("hOP-1", SEQ ID NOs: 1, mature protein amino acid sequence), or mouse OP-1 ("mOP-1", SEQ ID NO: 4, mature protein amino acid sequence.) The conserved seven cysteine skeleton is defined by
----	--------	--

- 5 residues 38 to 139 of SEQ ID NOs: 1 and 4. The cDNA sequences and the amino acids encoding the full length proteins are provided in SEQ ID NOs: 17 and 3 (hOP-1) and SEQ ID NOs: 18 and 19 (mOP-1). The mature proteins are defined by residues 293-431 (hOP-1) and 292-430 (mOP-1). The "pro" regions of the proteins, cleaved to yield the mature, morphogenically active proteins are defined essentially by residues 30-292 (hOP-1) and residues 30-291 (mOP-1).
- 10 "OP-2" refers generically to the group of active proteins expressed from part or all of a DNA sequence encoding OP-2 protein, including allelic and species variants thereof, e.g., human OP-2 ("hOP-2", SEQ ID No: 5, mature protein amino acid sequence) or mouse OP-2 ("mOP-2", SEQ ID No: 6, mature protein amino acid sequence). The conserved seven cysteine skeleton is defined by residues 38 to 139 of SEQ ID NOs: 5 and 6. The cDNA sequences and the amino acids encoding the full length proteins are provided in SEQ ID NOs: 20 and 21 (hOP-2) and SEQ ID NOs: 22 and 23 (mOP-2). The mature proteins are defined essentially by residues 264-402 (hOP-2) and 261-399 (mOP-2). The "pro" regions of the proteins, cleaved to yield the mature, morphogenically active proteins likely are defined essentially by residues 18-263 (hOP-2) and residues 18-260 (mOP-2). Another cleavage site also occurs 21 residues upstream for both OP-2 proteins.
- 15 "CBMP2" refers generically to the morphogenically active proteins expressed from a DNA sequence encoding the CBMP2 proteins, including allelic and species variants thereof, e.g., human CBMP2A ("CBMP2A(fx)"), SEQ ID NO: 10) or human CBMP2B DNA ("CBMP2B(fx)"), SEQ ID NO: 11). The amino acid sequence for the full length proteins, referred to in the literature as BMP2A and BMP2B, or BMP2 and BMP4, appear in Wozney, et al. (1988) Science 242:1528-1534, the content of which is incorporated by reference herein. The pro-domain for BMP2 (BMP2A) likely includes residues 25-248 or 25-282; the mature protein, residues 249-396 or 283-396. The pro-domain for BMP4 (BMP2B) likely includes residues 25-256 or 25-292; the mature protein, residues 257-408 or 293-408.
- 20 "DPP(fx)" refers to protein sequences encoded by the *Drosophila* DPP gene (DPP protein, see SEQ ID NO: 7) and defining the conserved seven cysteine skeleton. The amino acid sequence for the full length protein appears in Padgett, et al (1987) Nature 325: 81-84, the content of which is incorporated by reference herein. The pro-domain likely extends from the signal peptide cleavage site to residue 456; mature protein likely is defined by residues 457-588. The sequence of DPP(fx) is shown in Table II.
- 25 "Vgl(fx) " refers to protein sequences encoded by the *Xenopus* Vgl gene (Vgl protein, see SEQ ID NO: 8) and defining the conserved seven cysteine

- 5 skeleton. The amino acid sequence for the full length protein appears in Weeks (1987) Cell 51: 861-867, the content of which is incorporated by reference herein. The pro-domain likely extends from the signal peptide cleavage site to residue 246; the mature protein likely is defined by residues 247-360. The sequence of Vgl1(fx) is shown in Table II.
- 10 ""Vgr-1(fx)" refers to protein sequences encoded by the murine vgr-1 gene (Vgr-1 protein, see SEQ ID NO: 9) and defining the conserved seven cysteine skeleton. The amino acid sequence for the full length protein appears in Lyons, et al., (1989) PNAS 86: 4554-4558, the content of which is incorporated by reference herein. The pro-domain likely extends from the signal peptide cleavage site to residue 299; the mature protein likely is defined by residues 300-438. The sequence of Vgr-1(fx) is shown in Table II.
- 15 "GDF-1(fx)" refers to protein sequences encoded by the human GDF-1 gene (GDF-1 protein, see SEQ ID NO: 13) and defining the conserved seven cysteine skeleton. The amino acid sequence for the full length protein is provided in SEQ ID NO: 13. The pro-domain likely extends from the signal peptide cleavage site to residue 214; the mature protein likely is defined by residues 215-372. The sequence of GDF-1(fx) is shown in Table II.
- 20 "60A" refers generically to the morphogenically active proteins expressed from part or all of a DNA sequence (from the *Drosophila* 60A gene) encoding the 60A proteins (see SEQ ID NO: 14). "60A(fx)" refers to the protein sequences defining the conserved seven cysteine skeleton (residues 354 to 455 of SEQ ID NO: 14). The pro-domain likely extends from the signal peptide cleavage site to residue 324; the mature protein likely is defined by residues 325-455. The sequence of 60A(fx) is shown in Table II.
- 25 30 "BMP3(fx)" refers to protein sequences encoded by the human BMP3 gene (BMP3 protein, see SEQ ID NO: 12) and defining the conserved seven cysteine skeleton. The amino acid sequence for the full length protein appears in Wozney et al. (1988) Science 242: 1528-1534, the content of which is incorporated by reference herein. The pro-domain likely extends from the signal peptide cleavage site to residue 290; the mature protein likely is defined by residues 291-472. The sequence of BMP3(fx) is shown in Table II.
- 35 40 "BMP5(fx)" refers to protein sequences encoded by the human BMP5 gene (BMP5 protein, see SEQ ID NO: 15) and defining the conserved seven cysteine skeleton. The amino acid sequence for the full length protein appears in Celeste, et al. (1991) PNAS 87: 9843-9847, the content of which is incorporated by reference herein. The pro-domain likely extends from the signal peptide cleavage site to residue 316; the
- 45

mature protein likely is defined by residues 317-454. The sequence of BMP5(fx) is shown in Table II.

5 "BMP6(fx)" refers to protein sequences encoded by the human BMP6 gene (BMP6
protein, see SEQ ID NO: 16) and defining the conserved seven
cysteine skeleton. The amino acid sequence for the full length protein
appears in Celeste, et al. (1990) PNAS 87: 9843-9847, the content of
10 which is incorporated by reference herein. The pro-domain likely
includes extends from the signal peptide cleavage site to residue 374;
the mature sequence likely includes residues 375-513. The sequence
of BMP6(fx) is shown in Table II.

The OP-2 proteins have an "additional" cysteine residue in this region (e.g.,
15 see residue 41 of SEQ ID NOs: 21 and 23), in addition to the conserved cysteine
skeleton in common with the other proteins in this family. The GDF-1 protein has a
four amino acid insert within the conserved skeleton (compare SEQ ID NO: 19 with
SEQ ID NO: 13) but this insert likely does not interfere with the relationship of the
cysteines in the folded structure. In addition, the CBMP2 proteins are missing one
20 amino acid residue within the cysteine skeleton.

The morphogens are inactive when reduced, but are active as oxidized
homodimers and when oxidized in combination with other morphogens of this
invention (e.g., as heterodimers). Thus, as defined herein, a morphogen is a dimeric
protein comprising a pair of polypeptide chains, wherein each polypeptide chain
25 comprises at least the C-terminal six cysteine skeleton defined by residues 43-139 of
SEQ ID NO: 1, including functionally equivalent arrangements of these cysteines
(e.g., amino acid insertions or deletions which alter the linear arrangement of the
cysteines in the sequence but not their relationship in the folded structure), such that,
when the polypeptide chains are folded, the dimeric protein species comprising the
30 pair of polypeptide chains has the appropriate three-dimensional structure, including
the appropriate intra-or inter-chain disulfide bonds such that the protein is capable of
acting as a morphogen as defined herein. Specifically, the morphogens generally are
capable of all of the following biological functions in a morphogenically permissive
environment: stimulating proliferation of progenitor cells; stimulating the
35 differentiation of progenitor cells; stimulating the proliferation of differentiated
cells; and supporting the growth and maintenance of differentiated cells, including

the "redifferentiation" of transformed cells. In addition, it is also anticipated that these morphogens are capable of inducing redifferentiation of committed cells under appropriate environmental conditions.

The following publications disclose published morphogen polypeptide sequences, as well as relevant chemical and physical properties, of naturally occurring and/or synthetic morphogens: OP-1 and OP-2: U.S. 5,011,691, U.S. 5,266,683, Ozkaynak, *et al.*, *EMBO J.* 9: 2085-2093 (1990); OP-3: WO 94/10203 (PCT US93/10520); BMP-2, BMP-3, and BMP-4: WO 88/00205, Wozney, *et al.*, *Science* 242: 1528-1534 (1988); BMP-5 and BMP-6: Celeste, *et al.*, *PNAS* 87: 9843-9847 (1991); Vgr1: Lyons, *et al.*, *PNAS* 86: 4554-4558 (1989); DPP: Padgett, *et al.*, *Nature* 325: 81-84 (1987); Vg-1: Weeks *Cell* 51: 861-867 (1987); BMP-9: WO 95/33830 (PCT/US95/07084); BMP-10: WO 94/26893 (PCT/US94/05290); BMP-11: WO 94/26892 (PCT/US94/05288); BMP-12: WO 95/16035 (PCT/US94/14030); BMP-13: WO 95/16035 (PCT/US94/14030); GDF-1: WO 92/00382 (PCT/US91/04096) and Lee, *et al.*, *PNAS* 88:4250-4254 (1991); GDF-8: WO 94/21681 (PCT/US94/03019); GDF-9: WO 94/15966 (PCT/US94/00685); GDF-10: WO 95/10539 (PCT/US94/11440); GDF-11: WO 96/01845 (PCT/US95/08543); BMP-15: WO 96/36710 (PCT/US96/06540); *MP121*: WO 96/01316 (PCT/EP95/02552); GDF-5 (CDMP-1, MP52): WO 94/15949 (PCT/US94/00657) and WO 96/14335 (PCT/US94/12814) and WO 93/16099 (PCT/EP93/00350); GDF-6 (CDMP-2, BMP-13): WO 95/01801 (PCT/US94/07762) and WO 96/14335 and WO 95/10635 (PCT/US94/14030); GDF-7 (CDMP-3, BMP-12): WO 95/10802 (PCT/US94/07799) and WO 95/10635 (PCT/US94/14030). In another embodiment, useful proteins include biologically active biosynthetic constructs, including novel biosynthetic morphogenic proteins and chimeric proteins designed using sequences from two or more known morphogens. See also the biosynthetic constructs disclosed in U.S. Pat. 5,011,691 (*e.g.*, COP-1, COP-3, COP-4, COP-5, COP-7, and COP-16). The disclosure of all cited references describing morphogens and other related proteins are incorporated herein by reference.

In certain preferred embodiments, useful morphogenic proteins include those in which the amino acid sequences comprise a sequence sharing at least 70% amino

acid sequence homology (identity or conserved substitution), and preferably 80%, 85%, 90%, 95% or 99% homology, with a reference morphogenic protein selected from the exemplary naturally occurring morphogenic proteins listed herein. Preferably, the reference protein is human OP-1, and the reference sequence thereof

5 is the C-terminal seven cysteine skeleton present in osteogenically active forms of human OP-1, residues 330-431 of SEQ ID NO: 3 (or residues 38-139 of SEQ ID NO: 1). Useful morphogenic proteins accordingly include allelic, phylogenetic counterpart and other variants of the preferred reference sequence, whether naturally occurring or biosynthetically produced (*e.g.*, including "muteins" or "mutant

10 proteins"), as well as novel members of the general morphogenic family of proteins including those set forth and identified above. Certain particularly preferred morphogenic polypeptides share at least 50% amino acid identity with the preferred reference sequence of human OP-1, or any of the other morphogens described above, still more preferably at least 55%, 60%, 65%, 70%, 80%, 85%, 90%, 95%,

15 99% or more amino acid identity therewith.

FIG. 28 recites the percent amino acid sequence homology and percent identity within the C-terminal seven cysteine skeletons of various representative members of the TGF- β superfamily, using OP-1 as the reference sequence. The percent homologies recited in the figure are determined by aligning the sequences

20 using the MegaAlign Program (DNASTar, Inc.). Insertions and deletions from the reference morphogen sequence (the C-terminal, biologically active seven-cysteine skeleton of hOP-1) are ignored for purposes of calculation (details see below).

As is apparent to one of ordinary skill in the art reviewing the sequences for the proteins listed in FIG. 28, significant amino acid changes can be made from the

25 reference sequence while retaining substantial morphogenic activity. Moreover, GDF-1 contains a four amino acid insert (Gly-Gly-Pro-Pro, SEQ ID NO: 31) between the two residues corresponding to residue 372 and 373 of OP-1 (SEQ ID NO: 3). Similarly, BMP3 has a "extra" residue, a valine, inserted between the two residues corresponding to residues 385 and 386 of hOP-1 (SEQ ID NO: 3). Also,

30 BMP2 and BMP4 are both "missing" the amino acid residue corresponding to

residue 389 of OP-1 (SEQ ID NO: 3). None of these "deviations" from the reference sequence appear to interfere substantially with biological activity.

In other preferred embodiments, the family of morphogenic polypeptides useful in the present invention, and members thereof, are defined by a generic amino acid sequence. For example, Generic Sequence 1 (SEQ ID NO: 24) and Generic Sequence 2 (SEQ ID NO: 25) disclosed below, encompass the observed variations between preferred protein family members identified to date, including at least OP-1, OP-2, OP-3, CBMP2A, CBMP2B, BMP3, 60A, DPP, Vgl, BMP5, BMP6; Vgr-1, and GDF-1. The amino acid sequences for these proteins are described herein and/or in the art, as summarized above. The generic sequences include both the amino acid identity shared by these sequences in the C-terminal skeleton, defined by the six and seven cysteine skeletons (Generic Sequences 1 and 2, respectively), as well as alternative residues for the variable positions within the sequence. The generic sequences provide an appropriate cysteine skeleton where inter- or intra-molecular disulfide bonds can form, and contain certain critical amino acids likely to influence the tertiary structure of the folded proteins. In addition, the generic sequences allow for an additional cysteine at position 36 (Generic Sequence 1) or position 41 (Generic Sequence 2), thereby encompassing the morphogenically active sequences of OP-2 and OP-3.

Generic Sequence 1 (SEQ ID NO: 24)

			Leu	Xaa	Xaa	Xaa	Phe	Xaa	Xaa	
			1				5			
	Xaa	Gly	Trp	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Pro
25			10				15			
	Xaa	Xaa	Xaa	Xaa	Ala	Xaa	Tyr	Cys	Xaa	Gly
			20				25			
	Xaa	Cys	Xaa	Xaa	Pro	Xaa	Xaa	Xaa	Xaa	Xaa
			30				35			

	Xaa	Xaa	Xaa	Asn	His	Ala	Xaa	Xaa	Xaa	Xaa
			40					45		
	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
			50					55		
5	Xaa	Xaa	Xaa	Cys	Cys	Xaa	Pro	Xaa	Xaa	Xaa
			60					65		
	Xaa	Xaa	Xaa	Xaa	Xaa	Leu	Xaa	Xaa	Xaa	Xaa
			70					75		
	Xaa	Xaa	Xaa	Val	Xaa	Leu	Xaa	Xaa	Xaa	Xaa
10			80					85		
	Xaa	Met	Xaa	Val	Xaa	Xaa	Cys	Xaa	Cys	Xaa
			90					95		

wherein each Xaa independently is selected from a group of one or more specified amino acids defined as follows: "Res." means "residue" and Xaa at res. 2 = (Tyr or
15 Lys); Xaa at res. 3 = Val or Ile); Xaa at res. 4 = (Ser, Asp or Glu); Xaa at res. 6 = (Arg, Gln, Ser, Lys or Ala); Xaa at res. 7 = (Asp or Glu); Xaa at res. 8 = (Leu, Val or Ile); Xaa at res. 11 = (Gln, Leu, Asp, His, Asn or Ser); Xaa at res. 12 = (Asp, Arg, Asn or Glu); Xaa at res. 13 = (Trp or Ser); Xaa at res. 14 = (Ile or Val); Xaa at res. 15 = (Ile or Val); Xaa at res. 16 (Ala or Ser); Xaa at res. 18 = (Glu, Gln,
20 Leu, Lys, Pro or Arg); Xaa at res. 19 = (Gly or Ser); Xaa at res. 20 = (Tyr or Phe); Xaa at res. 21 = (Ala, Ser, Asp, Met, His, Gln, Leu or Gly); Xaa at res. 23 = (Tyr, Asn or Phe); Xaa at res. 26 = (Glu, His, Tyr, Asp, Gln, Ala or Ser); Xaa at res. 28 = (Glu, Lys, Asp, Gln or Ala); Xaa at res. 30 = (Ala, Ser, Pro, Gln, Ile or Asn); Xaa at res. 31 = (Phe, Leu or Tyr); Xaa at res. 33 = (Leu, Val or Met); Xaa at res.

34 = (Asn, Asp, Ala, Thr or Pro); Xaa at res. 35 = (Ser, Asp, Glu, Leu, Ala or
Lys); Xaa at res. 36 = (Tyr, Cys, His, Ser or Ile); Xaa at res. 37 = (Met, Phe, Gly
or Leu); Xaa at res. 38 = (Asn, Ser or Lys); Xaa at res. 39 = (Ala, Ser, Gly or Pro);
Xaa at res. 40 = (Thr, Leu or Ser); Xaa at res. 44 = (Ile, Val or Thr); Xaa at res. 45
5 = (Val, Leu, Met or Ile); Xaa at res. 46 = (Gln or Arg); Xaa at res. 47 = (Thr, Ala
or Ser); Xaa at res. 48 = (Leu or Ile); Xaa at res. 49 = (Val or Met); Xaa at res. 50
= (His, Asn or Arg); Xaa at res. 51 = (Phe, Leu, Asn, Ser, Ala or Val); Xaa at res.
52 = (Ile, Met, Asn, Ala, Val, Gly or Leu); Xaa at res. 53 = (Asn, Lys, Ala, Glu,
Gly or Phe); Xaa at res. 54 = (Pro, Ser or Val); Xaa at res. 55 = (Glu, Asp, Asn,
10 Gly, Val, Pro or Lys); Xaa at res. 56 = (Thr, Ala, Val, Lys, Asp, Tyr, Ser, Gly, Ile
or His); Xaa at res. 57 = (Val, Ala or Ile); Xaa at res. 58 = (Pro or Asp); Xaa at res.
59 = (Lys, Leu or Glu); Xaa at res. 60 = (Pro, Val or Ala); Xaa at res. 63 = (Ala or
Val); Xaa at res. 65 = (Thr, Ala or Glu); Xaa at res. 66 = (Gln, Lys, Arg or Glu);
Xaa at res. 67 = (Leu, Met or Val); Xaa at res. 68 = (Asn, Ser, Asp or Gly); Xaa at
15 res. 69 = (Ala, Pro or Ser); Xaa at res. 70 = (Ile, Thr, Val or Leu); Xaa at res. 71 =
(Ser, Ala or Pro); Xaa at res. 72 = (Val, Leu, Met or Ile); Xaa at res. 74 = (Tyr or
Phe); Xaa at res. 75 = (Phe, Tyr, Leu or His); Xaa at res. 76 = (Asp, Asn or Leu);
Xaa at res. 77 = (Asp, Glu, Asn, Arg or Ser); Xaa at res. 78 = (Ser, Gln, Asn, Tyr
or Asp); Xaa at res. 79 = (Ser, Asn, Asp, Glu or Lys); Xaa at res. 80 = (Asn, Thr or
20 Lys); Xaa at res. 82 = (Ile, Val or Asn); Xaa at res. 84 = (Lys or Arg); Xaa at res.
85 = (Lys, Asn, Gln, His, Arg or Val); Xaa at res. 86 = (Tyr, Glu or His); Xaa at
res. 87 = (Arg, Gln, Glu or Pro); Xaa at res. 88 = (Asn, Glu, Trp or Asp); Xaa at
res. 90 = (Val, Thr, Ala or Ile); Xaa at res. 92 = (Arg, Lys, Val, Asp, Gln or Glu);

Xaa at res. 93 = (Ala, Gly, Glu or Ser); Xaa at res. 95 = (Gly or Ala) and Xaa at res. 97 = (His or Arg).

Generic Sequence 2 (SEQ ID NO: 25) includes all of Generic Sequence 1 (SEQ ID NO: 24) and in addition includes the following sequence (SEQ ID NO: 26) at its N-terminus:

SEQ ID NO: 26

Cys	Xaa	Xaa	Xaa	Xaa
1				5

Accordingly, beginning with residue 7, each "Xaa" in Generic Sequence 2 is a specified amino acid defined as for Generic Sequence 1, with the distinction that each residue number described for Generic Sequence 1 is shifted by five in Generic Sequence 2. Thus, "Xaa. at res. 2 = (Tyr or Lys)" in Generic Sequence 1 refers to Xaa at res. 7 in Generic Sequence 2. In Generic Sequence 2, Xaa at res. 2 = (Lys, Arg, Ala or Gln); Xaa at res. 3 = (Lys, Arg or Met); Xaa at res. 4 = (His, Arg or Gln); and Xaa at res. 5 = (Glu, Ser, His, Gly, Arg, Pro, Thr, or Tyr).

In another embodiment, useful osteogenic proteins include those defined by Generic Sequences 3 and 4 (SEQ ID NOs: 27 and 28, respectively), described herein above. Specifically, Generic Sequences 3 and 4 are composite amino acid sequences of the following proteins: human OP-1, human OP-2, human OP-3, human BMP2, human BMP3, human BMP4, human BMP5, human BMP6, human BMP8, human BMP9, human BMP10, human BMP11, Drosophila 60A, Xenopus Vgl, sea urchin UNIVIN, human CBMP1 (mouse GDF-5), human CBMP2 (mouse GDF-6, human BMP13), human CBMP3 (mouse GDF-7, human BMP12), mouse GDF3, human GDF-1, mouse GDF-1, chicken DORSALIN, Drosophila DPP, Drosophila SCREW, mouse NODAL, mouse GDF-8, human GDF-8, mouse GDF-9, mouse GDF-10, human GDF-11, mouse GDF-11, human BMP15, and rat BMP3b. Like Generic Sequence 1, Generic Sequence 3 accommodates the C-terminal six cysteine skeleton and, like Generic Sequence 2, Generic Sequence 4 accommodates the seven cysteine skeleton.

Generic Sequence 3 (SEQ ID NO: 27)

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
1				5					10
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Pro	Xaa	Xaa	Xaa
				15					20
Xaa	Xaa	Xaa	Xaa	Cys	Xaa	Gly	Xaa	Cys	Xaa
				25					30
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				35					40
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				45					50
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				55					60
Xaa	Cys	Xaa	Pro	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				65					70
Xaa	Xaa	Leu	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				75					80
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				85					90
Xaa	Xaa	Xaa	Cys	Xaa	Cys	Xaa			
				95					

wherein each Xaa is independently selected from a group of one or more specified amino acids defined as follows: "Res." means "residue" and Xaa at res. 1 = (Phe,

5 Leu or Glu); Xaa at res. 2 = (Tyr, Phe, His, Arg, Thr, Lys, Gln, Val or Glu); Xaa at res. 3 = (Val, Ile, Leu or Asp); Xaa at res. 4 = (Ser, Asp, Glu, Asn or Phe); Xaa at res. 5 = (Phe or Glu); Xaa at res. 6 = (Arg, Gln, Lys, Ser, Glu, Ala or Asn); Xaa at res. 7 = (Asp, Glu, Leu, Ala or Gln); Xaa at res. 8 = (Leu, Val, Met, Ile or Phe); Xaa at res. 9 = (Gly, His or Lys); Xaa at res. 10 = (Trp or Met); Xaa at res. 11 =

10 (Gln, Leu, His, Glu, Asn, Asp, Ser or Gly); Xaa at res. 12 = (Asp, Asn, Ser, Lys, Arg, Glu or His); Xaa at res. 13 = (Trp or Ser); Xaa at res. 14 = (Ile or Val); Xaa at res. 15 = (Ile or Val); Xaa at res. 16 = (Ala, Ser, Tyr or Trp); Xaa at res. 18 = (Glu, Lys, Gln, Met, Pro, Leu, Arg, His or Lys); Xaa at res. 19 = (Gly, Glu, Asp, Lys, Ser, Gln, Arg or Phe); Xaa at res. 20 = (Tyr or Phe); Xaa at res. 21 = (Ala, Ser,

Gly, Met, Gln, His, Glu, Asp, Leu, Asn, Lys or Thr); Xaa at res. 22 = (Ala or Pro);
Xaa at res. 23 = (Tyr, Phe, Asn, Ala or Arg); Xaa at res. 24 = (Tyr, His, Glu, Phe
or Arg); Xaa at res. 26 = (Glu, Asp, Ala, Ser, Tyr, His, Lys, Arg, Gln or Gly); Xaa
at res. 28 = (Glu, Asp, Leu, Val, Lys, Gly, Thr, Ala or Gln); Xaa at res. 30 = (Ala,
5 Ser, Ile, Asn, Pro, Glu, Asp, Phe, Gln or Leu); Xaa at res. 31 = (Phe, Tyr, Leu,
Asn, Gly or Arg); Xaa at res. 32 = (Pro, Ser, Ala or Val); Xaa at res. 33 = (Leu,
Met, Glu, Phe or Val); Xaa at res. 34 = (Asn, Asp, Thr, Gly, Ala, Arg, Leu or Pro);
Xaa at res. 35 = (Ser, Ala, Glu, Asp, Thr, Leu, Lys, Gln or His); Xaa at res. 36 =
(Tyr, His, Cys, Ile, Arg, Asp, Asn, Lys, Ser, Glu or Gly); Xaa at res. 37 = (Met,
10 Leu, Phe, Val, Gly or Tyr); Xaa at res. 38 = (Asn, Glu, Thr, Pro, Lys, His, Gly,
Met, Val or Arg); Xaa at res. 39 = (Ala, Ser, Gly, Pro or Phe); Xaa at res. 40 =
(Thr, Ser, Leu, Pro, His or Met); Xaa at res. 41 = (Asn, Lys, Val, Thr or Gln); Xaa
at res. 42 = (His, Tyr or Lys); Xaa at res. 43 = (Ala, Thr, Leu or Tyr); Xaa at res.
44 = (Ile, Thr, Val, Phe, Tyr, Met or Pro); Xaa at res. 45 = (Val, Leu, Met, Ile or
15 His); Xaa at res. 46 = (Gln, Arg or Thr); Xaa at res. 47 = (Thr, Ser, Ala, Asn or
His); Xaa at res. 48 = (Leu, Asn or Ile); Xaa at res. 49 = (Val, Met, Leu, Pro or
Ile); Xaa at res. 50 = (His, Asn, Arg, Lys, Tyr or Gln); Xaa at res. 51 = (Phe, Leu,
Ser, Asn, Met, Ala, Arg, Glu, Gly or Gln); Xaa at res. 52 = (Ile, Met, Leu, Val,
Lys, Gln, Ala or Tyr); Xaa at res. 53 = (Asn, Phe, Lys, Glu, Asp, Ala, Gln, Gly,
20 Leu or Val); Xaa at res. 54 = (Pro, Asn, Ser, Val or Asp); Xaa at res. 55 = (Glu,
Asp, Asn, Lys, Arg, Ser, Gly, Thr, Gln, Pro or His); Xaa at res. 56 = (Thr, His,
Tyr, Ala, Ile, Lys, Asp, Ser, Gly or Arg); Xaa at res. 57 = (Val, Ile, Thr, Ala, Leu
or Ser); Xaa at res. 58 = (Pro, Gly, Ser, Asp or Ala); Xaa at res. 59 = (Lys, Leu,
Pro, Ala, Ser, Glu, Arg or Gly); Xaa at res.: 60 = (Pro, Ala, Val, Thr or Ser); Xaa

at res. 61 = (Cys, Val or Ser); Xaa at res. 63 = (Ala, Val or Thr); Xaa at res. 65 =
 (Thr, Ala, Glu, Val, Gly, Asp or Tyr); Xaa at res. 66 = (Gln, Lys, Glu, Arg or Val);
 Xaa at res. 67 = (Leu, Met, Thr or Tyr); Xaa at res. 68 = (Asn, Ser, Gly, Thr, Asp,
 Glu, Lys or Val); Xaa at res. 69 = (Ala, Pro, Gly or Ser); Xaa at res. 70 = (Ile, Thr,
 5 Leu or Val); Xaa at res. 71 = (Ser, Pro, Ala, Thr, Asn or Gly); Xaa at res. 2 = (Val,
 Ile, Leu or Met); Xaa at res. 74 = (Tyr, Phe, Arg, Thr, Tyr or Met); Xaa at res. 75 =
 (Phe, Tyr, His, Leu, Ile, Lys, Gln or Val); Xaa at res. 76 = (Asp, Leu, Asn or Glu);
 Xaa at res. 77 = (Asp, Ser, Arg, Asn, Glu, Ala, Lys, Gly or Pro); Xaa at res. 78 =
 (Ser, Asn, Asp, Tyr, Ala, Gly, Gln, Met, Glu, Asn or Lys); Xaa at res. 79 = (Ser,
 10 Asn, Glu, Asp, Val, Lys, Gly, Gln or Arg); Xaa at res. 80 = (Asn, Lys, Thr, Pro,
 Val, Ile, Arg; Ser or Gln); Xaa at res. 81 = (Val, Ile, Thr or Ala); Xaa at res. 82 =
 (Ile, Asn, Val, Leu, Tyr, Asp or Ala); Xaa at res. 83 = (Leu, Tyr, Lys or Ile); Xaa
 at res. 84 = (Lys, Arg, Asn, Tyr, Phe, Thr, Glu or Gly); Xaa at res. 85 = (Lys, Arg,
 His, Gln, Asn, Glu or Val); Xaa at res. 86 = (Tyr, His, Glu or Ile); Xaa at res. 87 =
 15 (Arg, Glu, Gln, Pro or Lys); Xaa at res. 88 = (Asn, Asp, Ala, Glu, Gly or Lys);
 Xaa at res. 89 = (Met or Ala); Xaa at res. 90 = (Val, Ile, Ala, Thr, Ser or Lys); Xaa
 at res. 91 = (Val or Ala); Xaa at res. 92 = (Arg, Lys, Gln, Asp, Glu, Val, Ala, Ser
 or Thr); Xaa at res. 93 = (Ala, Ser, Glu, Gly, Arg or Thr); Xaa at res. 95 = (Gly,
 Ala or Thr); Xaa at res. 97 = (His, Arg, Gly, Leu or Ser). Further, after res. 53 in
 20 rBMP-3b and mGDF-10 there is an Ile; after res. 54 in GDF-1 there is a T; after
 res. 54 in BMP3 there is a V; after res. 78 in BMP8 and Dorsalin there is a G; after
 res. 37 in hGDF-1 there is Pro, Gly, Gly, Pro.

Generic Sequence 4 (SEQ ID NO: 28) includes all of Generic Sequence 3
 (SEQ ID NO: 27) and in addition includes the following sequence (SEQ ID NO: 26)
 25 at its N-terminus:

SEQ ID NO: 26

Cys	Xaa	Xaa	Xaa	Xaa
1				5

Accordingly, beginning with residue 6, each "Xaa" in Generic Sequence 4 is a specified amino acid defined as for Generic Sequence 3, with the distinction that each residue number described for Generic Sequence 3 is shifted by five in Generic Sequence 4. Thus, "Xaa at res. 1 = (Tyr, Phe, His, Arg, Thr, Lys, Gln, Val or Glu)" in Generic Sequence 3 refers to Xaa at res. 6 in Generic Sequence 4. In Generic Sequence 4, Xaa at res. 2 = (Lys, Arg, Gln, Ser, His, Glu, Ala, or Cys); Xaa at res. 3 = (Lys, Arg, Met, Lys, Thr, Leu, Tyr, or Ala); Xaa at res. 4 = (His, Gln, Arg, Lys, Thr, Leu, Val, Pro, or Tyr); and Xaa at res. 5 = (Gln, Thr, His, Arg, Pro, Ser, Ala, Gln, Asn, Tyr, Lys, Asp, or Leu).

Based upon alignment of the naturally occurring morphogens within the definition of Generic Sequence 4, it should be clear that gaps and/or insertions of one or more amino acid residues can be tolerated (without abrogating or substantially impairing biological activity) at least between or involving residues 11-12, 42-43, 59-60, 68-69 and 83-84.

Particularly useful sequences for use as morphogens in this invention include the C-terminal domains, e.g., the C-terminal 96-102 amino acid residues of Vgl, Vgr-1, DPP, OP-1, OP-2, CBMP2A, CBMP2B, GDF-1 (see Table II, below, and SEQ ID NOs: 1-13), as well as proteins comprising the C-terminal domains of 60A, BMP3, BMP5 and BMP6 (see SEQ ID NOs: 12, 14-16), all of which include at least the conserved six or seven cysteine skeleton. In addition, biosynthetic constructs designed from the generic sequences, such as COP-1, 3-5, 7, 16, disclosed in U.S. Pat. No. 5,011,691, also are useful. Other sequences include the inhibins/activin proteins (see, for example, U.S. Pat. Nos. 4,968,590 and 5,011,691). Accordingly, other useful sequences are those sharing at least 70% amino acid sequence homology or 50% identity, and preferably 80% homology or 70% identity with any of the sequences above. These are anticipated to include allelic and species variants and mutants, and biosynthetic muteins, as well as novel members of this

morphogenic family of proteins. Particularly envisioned in the family of related proteins are those proteins exhibiting morphogenic activity and wherein the amino acid changes from the preferred sequences include conservative changes.

Information regarding conserved amino acid changes are well-known in the art. For example, Dayhoff et al. described in *Atlas of Protein Sequence and Structure*; vol. 5, Suppl. 3, pp. 345-362, (M.O. Dayhoff, ed., Nat'l BioMed. Research Fdn., Washington, D.C. 1978) that certain amino acids substitutions among evolutionary conserved proteins occur at higher than expected frequency than random chance would allow. Thus, conserved amino acid substitutions can be determined according to Figure 84 (*supra*). As used herein, potentially useful sequences are aligned with a known morphogen sequence using the method of Needleman et al. ((1970) *J. Mol. Biol.* 48:443-453) and identities calculated by the MegaAlign program (DNASTar, Inc.).

Table II, set forth below, compares the amino acid sequences of the active regions of native proteins that have been identified as morphogens, including human OP-1 (hOP-1, SEQ ID NOs: 1-3), mouse OP-1 (mOP-1, SEQ ID NOs: 4 and 19), human and mouse OP-2 (SEQ ID NOs: 5, 6, 21, and 23), CBMP2A (SEQ ID NO: 10), CBMP2B (SEQ ID NO: 11), BMP3 (SEQ ID NO: 12), DPP (from *Drosophila*, SEQ ID NO: 7), Vgl (from *Xenopus*, SEQ ID NO: 8), Vgr-1 (from mouse, SEQ ID NO: 9), GDF-1 (from mouse, SEQ ID NOs: 13), 60A protein (from *Drosophila*, SEQ ID NOs: 14), BMP5 (SEQ ID NO: 15) and BMP6 (SEQ ID NO: 16). The sequences are aligned essentially following the method of Needleman et al. (1970) *J. Mol. Biol.*, 48: 443-453, calculated using the Align Program (DNASTar, Inc.) In the table, three dots indicates that the amino acid in that position is the same as the amino acid in hOP-1. Three dashes indicates that no amino acid is present in that position, and are included for purposes of illustrating homologies. For example, amino acid residue 60 of CBMP-2A and CBMP-2B is "missing". Of course, both these amino acid sequences in this region comprise Asn-Ser (residues 58, 59), with CBMP-2A then comprising Lys and Ile, whereas CBMP-2B comprises Ser and Ile.

TABLE II

hOP-1	Cys	Lys	Lys	His	Glu	Leu	Tyr	Val
mOP-1
hOP-2	...	Arg	Arg
mOP-2	...	Arg	Arg
DPP	...	Arg	Arg	...	Ser
Vgl	Lys	Arg	His
Vgr-1	Gly
CBMP-2A	Arg	...	Pro
CBMP-2B	...	Arg	Arg	...	Ser
BMP3	...	Ala	Arg	Arg	Tyr	...	Lys	...
GDF-1	...	Arg	Ala	Arg	Arg
60A	...	Gln	Met	Glu	Thr
BMP5
BMP6	...	Arg
	1				5			

hOP-1	Ser	Phe	Arg	Asp	Leu	Gly	Trp	Gln	Asp
mOP-1
hOP-2	Gln	Leu	...
mOP-2	Ser	Leu	...
DPP	Asp	...	Ser	...	Val	Asp	...
Vgl	Glu	...	Lys	...	Val	Asn
Vgr-1	Gln	...	Val
CBMP-2A	Asp	...	Ser	...	Val	Asn	...
CBMP-2B	Asp	...	Ser	...	Val	Asn	...
BMP3	Asp	...	Ala	...	Ile	Ser	Glu
GDF-1	Glu	Val	His	Arg
60A	Asp	...	Lys	His	...
BMP5
BMP6	Gln
		10					15		

hOP-1	Trp	Ile	Ile	Ala	Pro	Glu	Gly	Tyr	Ala
mOP-1
hOP-2	...	Val	Gln	Ser
mOP-2	...	Val	Gln	Ser
DPP	Val	Leu	Asp
Vgl	...	Val	Gln	Met
Vgr-1	Lys
CBMP-2A	Val	Pro	His
CBMP-2B	Val	Pro	Gln
BMP3	Ser	...	Lys	Ser	Phe	Asp
GDF-1	...	Val	Arg	...	Phe	Leu
60A	Gly
BMP5

BMP6	Lys
			20				25		
hOP-1	Ala	Tyr	Tyr	Cys	Glu	Gly	Glu	Cys	Ala
mOP-1
hOP-2	Ser
mOP-2
DPP	His	...	Lys	...	Pro
Vgl	...	Asn	Tyr	Pro
Vgr-1	...	Asn	Asp	Ser
CBMP-2A	...	Phe	His	...	Glu	...	Pro
CBMP-2B	...	Phe	His	...	Asp	...	Pro
BMP3	Ser	...	Ala	...	Gln
GDF-1	...	Asn	Gln	...	Gln
60A	...	Phe	Ser	Asn
BMP5	...	Phe	Asp	Ser
BMP6	...	Asn	Asp	Ser
				30					35
hOP-1	Phe	Pro	Leu	Asn	Ser	Tyr	Met	Asn	Ala
mOP-1
hOP-2	Asp	...	Cys
mOP-2	Asp	...	Cys
DPP	Ala	Asp	His	Phe	...	Ser
Vgl	Tyr	Thr	Glu	Ile	Leu	...	Gly
Vgr-1	Ala	His
CBMP-2A	Ala	Asp	His	Leu	...	Ser
CBMP-2B	Ala	Asp	His	Leu	...	Ser
BMP3	Leu	...	Val	Ala	Leu	Ser	Gly	Ser†	...
GDF-1	Met	Pro	Lys	Ser	Leu	Lys	Pro
60A	Ala	His
BMP5	Ala	His	Met
BMP6	Ala	His	Met
					40				
hOP-1	Thr	Asn	His	Ala	Ile	Val	Gln	Thr	Leu
mOP-1
hOP-2	Leu	...	Ser	...
mOP-2	Leu	...	Ser	...
DPP	Val
Vgl	Ser	Leu
Vgr-1
CBMP-2A
CBMP-2B
BMP3	Ser	Thr	Ile	...	Ser	Ile
GDF-1	Leu	Val	Leu	Arg	Ala	...

60A
BMP5
BMP6
	45					50			
hOP-1	Val	His	Phe	Ile	Asn	Pro	Glu	Thr	Val
mOP-1	Asp
hOP-2	...	His	Leu	Met	Lys	...	Asn	Ala	...
mOP-2	...	His	Leu	Met	Lys	...	Asp	Val	...
DPP	...	Asn	Asn	Asn	Gly	Lys	...
Vgl	Ser	...	Glu	Asp	Ile
Vgr-1	Val	Met	Tyr	...
CBMP-2A	...	Asn	Ser	Val	...	Ser	---	Lys	Ile
CBMP-2B	...	Asn	Ser	Val	...	Ser	---	Ser	Ile
BMP3	...	Arg	Ala†	Gly	Val	Val	Pro	Gly	Ile
GDF-1	Met	...	Ala	Ala	Ala	...	Gly	Ala	Ala
60A	Leu	Leu	Glu	...	Lys	Lys	...
BMP5	Leu	Met	Phe	...	Asp	His	...
BMP6	Leu	Met	Tyr	...
		55					60		
hOP-1	Pro	Lys	Pro	Cys	Cys	Ala	Pro	Thr	Gln
mOP-1
hOP-2	Ala	Lys
mOP-2	Ala	Lys
DPP	Ala	Val
Vgl	...	Leu	Val	Lys
Vgr-1	Lys
CBMP-2A	Ala	Val	Glu
CBMP-2B	Ala	Val	Glu
BMP3	...	Glu	Val	...	Glu	Lys
GDF-1	Asp	Leu	Val	...	Ala	Arg
60A	Arg
BMP5	Lys
BMP6	Lys
			65					70	
hOP-1	Leu	Asn	Ala	Ile	Ser	Val	Leu	Tyr	Phe
mOP-1
hOP-2	...	Ser	...	Thr	Tyr
mOP-2	...	Ser	...	Thr	Tyr
Vgl	Met	Ser	Pro	Met	...	Phe	Tyr
Vgr-1	Val
DPP	...	Asn	Ser	Val	Ala	Met	Leu
CBMP-2A	...	Ser	Met	Leu
CBMP-2B	...	Ser	Met	Leu

BMP3	Met	Ser	Ser	Leu	...	Ile	...	Phe	Tyr
GDF-1	...	Ser	Pro	Phe	...
60A	...	Gly	...	Leu	Pro	His
BMP5
BMP6
				75					80
hOP-1	Asp	Asp	Ser	Ser	Asn	Val	Ile	Leu	Lys
mOP-1
hOP-2	...	Ser	...	Asn	Arg
mOP-2	...	Ser	...	Asn	Arg
DPP	Asn	...	Gln	...	Thr	...	Val
Vgl	...	Asn	Asn	Asp	Val	...	Arg
Vgr-1	Asn
CBMP-2A	...	Glu	Asn	Glu	Lys	...	Val
CBMP-2B	...	Glu	Tyr	Asp	Lys	...	Val
BMP3	...	Glu	Asn	Lys	Val
GDF-1	...	Asn	...	Asp	Val	...	Arg
60A	Leu	Asn	Asp	Glu	Asn
BMP5
BMP6	Asn
					85				
hOP-1	Lys	Tyr	Arg	Asn	Met	Val	Val	Arg	
mOP-1	
hOP-2	...	His	Lys	
mOP-2	...	His	Lys	
DPP	Asn	...	Gln	Glu	...	Thr	...	Val	
Vgl	His	...	Glu	Ala	...	Asp	
Vgr-1	
CBMP-2A	Asn	...	Gln	Asp	Glu	
CBMP-2B	Asn	...	Gln	Glu	Glu	
BMP3	Val	...	Pro	Thr	...	Glu	
GDF-1	Gln	...	Glu	Asp	Asp	
60A	Ile	...	Lys	
BMP5	
BMP6	Trp	
	90						95		
hOP-1	Ala	Cys	Gly	Cys	His				
mOP-1				
hOP-2				
mOP-2				
DPP	Gly	Arg				
Vgl	Glu	Arg				
Vgr-1				

CBMP-2A	Gly	Arg
CBMP-2B	Gly	Arg
BMP3	Ser	...	Ala	...	Arg
GDF-1	Glu	Arg
60A	Ser
BMP5	Ser
BMP6
100					

† Between residues 56 and 57 of BMP3 is a Val residue; between residues 43 and 44 of GDF-1 lies the amino acid sequence Gly-Gly-Pro-Pro.

As is apparent from the foregoing amino acid sequence comparisons,
5 significant amino acid changes can be made within the generic sequences while retaining the morphogenic activity.

As noted above, certain preferred morphogenic polypeptide sequences useful in this invention have greater than 50% identity, preferably greater than 55%, 60%, 65%, 70%, 80%, 85%, 90%, 95% or even 99% identity, with the amino acid
10 sequence defining the preferred reference sequence of hOP-1 (especially the conserved six-seven cysteine skeleton of hOP-1, e.g., residues 39-139 of SEQ ID No: 1), or equivalent regions from other morphogens described in the application. These particularly preferred sequences include allelic and phylogenetic counterpart variants of the OP-1 and OP-2 proteins, including the *Drosophila* 60A protein, as
15 well as the closely related proteins BMP5, BMP6 and Vgr-1. Accordingly, in certain particularly preferred embodiments, useful morphogenic proteins include active proteins comprising pairs of polypeptide chains within the generic amino acid sequence herein referred to as "OPX" (SEQ ID NO: 29), which defines the seven cysteine skeleton and accommodates the homologies between several identified
20 variants of OP-1 and OP-2. Accordingly, each "Xaa" at a given position in OPX independently is selected from the residues occurring at the corresponding position in the C-terminal sequence of mouse or human OP-1 or-OP-2. Specifically, each "Xaa" is independently selected from a group of one or more specified amino acids as defined below:

25

Cys 1	Xaa	Xaa	His	Glu 5	Leu	Tyr	Val	Ser	Phe 10	Xaa	Asp	Leu	Gly	Trp 15
Xaa	Asp	Trp	Xaa	Ile 20	Ala	Pro	Xaa	Gly	Tyr 25	Xaa	Ala	Tyr	Tyr	Cys 30
Glu	Gly	Glu	Cys	Xaa 35	Phe	Pro	Leu	Xaa	Ser 40	Xaa	Met	Asn	Ala	Thr 45
Asn	His	Ala	Ile	Xaa 50	Gln	Xaa	Leu	Val	His 55	Xaa	Xaa	Xaa	Pro	Xaa 60
Xaa	Val	Pro	Lys	Xaa 65	Cys	Cys	Ala	Pro	Thr 70	Xaa	Leu	Xaa	Ala	Xaa 75
Ser	Val	Leu	Tyr	Xaa 80	Asp	Xaa	Ser	Xaa	Asn 85	Val	Ile	Leu	Xaa	Lys 90
Xaa	Arg	Asn	Met	Val 95	Xaa	Ala	Cys	Gly	Cys 100	His				

wherein Xaa at res. 2 = (Lys or Arg); Xaa at res. 3 = (Lys or Arg); Xaa at res. 11 = (Arg or Gln); Xaa at res. 16 = (Gln or Leu); Xaa. at res. 19 = (Ile or Val); Xaa at res. 23 = (Glu or Gln); Xaa at res. 26 = (Ala or Ser); Xaa at res. 35 = (Ala or Ser); Xaa at res. 39 = (Asn or Asp); Xaa at res. 41 = (Tyr or Cys); Xaa at res. 50 = (Val or Leu); Xaa at res. 52 = (Ser or Thr); Xaa at res. 56 = (Phe or Leu); Xaa at res. 57 = (Ile or Met); Xaa at res. 58 = (Asn or Lys); Xaa at res. 60 = (Glu, Asp or Asn); Xaa at res. 61 = (Thr, Ala or Val); Xaa at res. 65 = (Pro or Ala); Xaa at res. 71 = (Gln or Lys); Xaa at res. 73 = (Asn or Ser); Xaa at res. 75 = (Ile or Thr); Xaa at res. 80 = (Phe or Tyr); Xaa at res. 82 = (Asp or Ser); Xaa at res. 84 = (Ser or Asn); Xaa at res. 89 = (Lys or Arg); Xaa at res. 91 = (Tyr or His); and Xaa at res. 97 = (Arg or Lys).

The following patents or publications or patent applications disclose morphogens or formula of useful / active morphogens, the entire contents of which are hereby incorporated by reference herein: EP 601106, and USSN 08/937,755 (filed on September 25, 1997).

The morphogens useful in the methods, composition and devices of this invention include proteins comprising any of the polypeptide chains described above, whether isolated from naturally occurring sources, or produced by recombinant DNA or other synthetic techniques, and includes allelic and species

variants of these proteins, naturally occurring or biosynthetic mutants thereof, as well as various truncated and fusion constructs. Deletion or addition mutants also are envisioned to be active, including those which may alter the conserved C-terminal cysteine skeleton, provided that the alteration does not functionally disrupt the relationship of these cysteines in the folded structure. Accordingly, such active forms are considered the equivalent of the specifically described constructs disclosed herein.

The proteins may include forms having varying glycosylation patterns, varying N-termini, a family of related proteins having regions of amino acid sequence homology, and active truncated or mutated forms of native or biosynthetic proteins, produced by expression of recombinant DNA in host cells.

The morphogenic proteins can be expressed from intact or truncated cDNA or from synthetic DNAs in prokaryotic or eukaryotic host cells, and purified, cleaved, refolded, and dimerized to form morphogenically active compositions. Currently preferred host cells include *E. coli* or any suitable mammalian host cells, such as CHO, COS or BSC cells. A detailed description of the morphogens useful in the methods, compositions and devices of this invention is disclosed in copending US patent application Serial Nos. 08/937755, filed September 25, 1997, and issued European Patent EP 601106, the contents of which are all incorporated by reference herein. Thus, in view of this disclosure, skilled genetic engineers can isolate genes from cDNA or genomic libraries of various different species which encode appropriate amino acid sequences, or construct DNAs from oligonucleotides, and then can express them in various types of host cells, including both prokaryotes and eukaryotes, to produce large quantities of active proteins capable of protecting tissues and organs from immune cell-mediated tissue destruction, including substantially inhibiting such damage and/or regenerating the damaged tissue in a variety of mammals, including humans.

In still another preferred embodiment, useful morphogenically active proteins have polypeptide chains with amino acid sequences comprising a sequence encoded by a nucleic acid that hybridizes with DNA or RNA encoding reference morphogen sequences, e.g., C-terminal sequences defining the conserved seven

cysteine skeletons of OP-1, OP-2, BMP2, BMP4, BMP5, BMP6, 60A, GDF-3, GDF-5, GDF-6, GDF-7 and the like. As used herein, high stringency hybridization conditions are defined as hybridization according to known techniques in 40% formamide, 5 X SSPE, 5 X Denhardt's Solution, and 0.1% SDS at 37 °C overnight, and washing in 0.1 X SSPE, 0.1% SDS at 50 °C. Standard stringency conditions are well characterized in standard molecular biology cloning texts. See, for example, MOLECULAR CLONING-A LABORATORY MANUAL, 2nd Ed., ed. by Sambrook, Fritsch and Maniatis (Cold Spring Harbor Laboratory Press: 1989); DNA CLONING, Volumes I and II (D.N. Glover ed., 1985); OLIGONUCLEOTIDE SYNTHESIS (M.J. Gait ed., 1984); NUCLEIC ACID HYBRIDIZATION (B. D. Hames & S.J. Higgins eds. 1984); and B. Perbal, A PRACTICAL GUIDE TO MOLECULAR CLONING (1984).

In other embodiments, as an alternative to the administration of a morphogenic protein, an effective amount of an agent competent to stimulate or induce increased endogenous morphogen expression in a mammal may be administered by any of the routes described herein. Such a morphogen inducer may be provided to a mammal, *e.g.*, by systemic administration to the mammal or by direct administration to the neural tissue. A method for identifying and testing inducers (stimulating agents) competent to modulate the levels of endogenous morphogens in a given tissue is described in published applications WO 93/05172 and WO 93/05751, each of which is incorporated by reference herein. Briefly, candidate compounds are identified and tested by incubation in vitro with test tissue or cells, or a cultured cell line derived therefrom, for a time sufficient to allow the compound to affect the production, *i.e.*, cause the expression and/or secretion, of a morphogen produced by the cells of that tissue. Suitable tissues, or cultured cells of a suitable tissue, are preferably selected from renal epithelium, ovarian tissue, fibroblasts, and osteoblasts.

In yet other embodiments, an agent which acts as an agonist of a morphogen receptor may be administered instead of the morphogen itself. Such an agent may also be referred to as a morphogen "mimic," "mimetic," or "analog." Thus, for example, a small peptide or other molecule which can mimic the activity of a morphogen in binding to and activating the morphogen's receptor- may be employed

as an equivalent of the morphogen. Preferably the agonist is a full agonist, but partial morphogen receptor agonists may also be advantageously employed. Methods of identifying such agonists are known in the art and include assays for compounds which induce morphogen-mediated responses (e.g., induction of
5 differentiation of metanephric mesenchyme, induction of endochondral bone formation). For example, methods of identifying morphogen inducers or agonists of morphogen receptors may be found in U.S.S.N. 08/478,097 filed June 7, 1995; U.S.S.N. 09/791946, filed Feb. 22, 2001; U.S. Pat. No. 5,834,188; U.S. Pat. No. 6,273,598; WO 97/26277; EP 0876401; U.S. Provisional Application No. 60/080032, filed on March 30, 1998; U.S. Provisional Application No. 60/296291,
10 filed on Jun. 10, 2001; U.S. Provisional Application No. 60/354820, filed on Feb. 5, 2002; and U.S. Provisional Application filed on April 10, 2002 (first named inventor Peter Keck, title: "MORPHOGEN ANALOGS AND METHODS FOR PRODUCING THEM"), the disclosures of which are incorporated herein by
15 reference.

The OP/BMP family of morphogens of the invention are also characterized by biological activities which may be readily ascertained by those of ordinary skill in the art. Specifically, a morphogen of the present invention is (a) capable of inducing chondrogenesis in the Reddi-Sampath ectopic bone assay (Sampath and
20 Reddi (1981), Proc. Natl. Acad. Sci. USA 78:7599-7603) or a substantially equivalent assay, (b) capable of significantly preventing, inhibiting, delaying or alleviating the progressive loss of renal function in a standard animal model of chronic renal failure, or (c) capable of causing a clinically significant improvement in a standard marker of renal function when administered to a mammal in, or at risk
25 of, chronic renal failure.

The Reddi-Sampath ectopic bone assay is well known in the art as an assay of chondrogenic activity. The assay, which can be easily performed, is described and discussed in, for example, Sampath and Reddi (1981), Proc. Natl. Acad. Sci. USA 78: 7599-7603; and function which characterizes chronic renal failure.

30 Finally, the morphogens of the present invention may be evaluated for their therapeutic efficacy in causing a clinically significant improvement in a standard

marker of renal function when administered to a mammalian subject (e.g., a human patient) in, or at risk of, chronic renal failure. Such markers of renal function are well known in the medical literature and include, without being limited to, rates of increase in BUN levels, rates of increase in serum creatinine, static measurements of
5 BUN, static measurements of serum creatinine, glomerular filtration rates (GFR), ratios of BUN/creatinine, serum concentrations of sodium (Na^+), urine/plasma ratios for creatinine, urine/plasma ratios for urea, urine osmolality, daily urine output, and the like (see, for example, Brenner and Lazarus (1994), in Harrison's Principles of Internal Medicine, 13th edition, Isselbacher et al., eds., McGraw Hill Text, New
10 York; Luke and Strom (1994), in Internal Medicine, 4th Edition, J.H. Stein, ed., Mosby-Year Book, Inc. St. Louis).

The morphogens contemplated herein can be expressed from intact or truncated genomic or cDNA or from synthetic DNAs in prokaryotic or eukaryotic host cells. The dimeric proteins can be isolated from the culture media and/or
15 refolded and dimerized in vitro to form biologically active compositions, Heterodimers can be formed in vitro by combining separate, distinct polypeptide chains. Alternatively, heterodimers can be formed in a single cell by co-expressing nucleic acids encoding separate, distinct polypeptide chains. See, for example, WO93/09229, or U.S. Pat. No. 5,411,941, for several exemplary recombinant
20 heterodimer protein production protocols. Currently preferred host cells include, without limitation, prokaryotes including *E. coli*, or eukaryotes including yeast (such as *S. cerevisiae*), insect cells, or any suitable mammalian host cells, such as CHO, COS or BSC cells. One of ordinary skill will appreciate that other host cells can be used to advantage. A detailed description of the morphogens useful in the methods,
25 compositions and devices of this invention, including how to make, use and test them for chondrogenic activity, are disclosed in numerous publications, including U.S. Pat. Nos. 5,266,683 and 5,011,691, the specifications of which are incorporated herein by reference.


As a general matter, methods of the present invention may be applied to the
30 treatment of any mammalian subject at risk of or afflicted with a neural tissue insult or neuropathy. The invention is suitable for the treatment of any primate, preferably

a higher primate such as a human. In addition, however, the invention may be employed in the treatment of domesticated mammals which are maintained as human companions (*e.g.*, dogs, cats, horses), which have significant commercial value (*e.g.*, goats, pigs, sheep, cattle, sporting or draft animals), which have
 5 significant scientific value (*e.g.*, captive or free specimens of endangered species, or inbred or engineered animal strains), or which otherwise have value.


(ii) Inhibitors of ACE (Angiotensin-Converting Enzyme)

Angiotensin I-converting enzyme (EC 3.4.15.1), or kininase II, is a dipeptidyl carboxypeptidase that plays an important role in blood pressure regulation
 10 and electrolyte balance by hydrolyzing angiotensin I into angiotensin II, a potent vasopressor, and aldosterone-stimulating peptide. The enzyme is also able to inactivate bradykinin, a potent vasodilator. The ACE gene encodes 2 isozymes. The somatic ACE isozyme is expressed in many tissues, including vascular endothelial cells, renal epithelial cells, and testicular Leydig cells, whereas the testicular or
 15 germinal ACE isozyme is expressed only in sperm (Ramaraj et al., *J. Clin. Invest.* 102: 371-378, 1998).

The angiotensin converting enzyme (ACE) inhibitors of the present invention may include 3-amino-[1]benzazepin-2-one-1-alkanoic acids and derivatives, as disclosed in U.S. Patent Nos. 4,473,575 and 4,410,520 (the entire contents of which
 20 are incorporated herein by reference), and characterized by formula (I) shown below:



wherein R_A and R_B are radicals of the formula:



respectively, in which:

R₀ is carboxy or a functionally modified carboxy;

5 R₁ is hydrogen, lower alkyl, amino(lower)alkyl, aryl, aryl(lower)alkyl, cycloalkyl, cycloalkyl(lower)alkyl, acylamino(lower) alkyl, mono- or di-(lower)alkylamino(lower)alkyl, lower alkylthio(lower)alkyl, carboxy(lower)alkyl, esterified carboxy(lower)alkyl, carbamoyl(lower)alkyl, etherified or acylated hydroxyl(lower)alkyl, aryloxy(-lower)alkyl, aryl-(thio-, sulfinyl-, or sulfonyl-)lower
10 alkyl, aryl-N-(lower)alkylamino(lower)alkyl, or arylamino(lower)alkyl;

R₂ is hydrogen or lower alkyl;

R₃ and R₄, each independently, represent hydrogen, lower alkyl, lower alkoxy, lower alkanoyloxy, hydroxyl, halogen, trifluoromethyl, or


R₃ and R₄ taken together represent lower alkylenedioxy;

15 R₅ is hydrogen or lower alkyl, and

X represents oxo, two hydrogens, or one hydroxyl or acylated hydroxy together with one hydrogen; and wherein the carbocyclic ring may also be hexahydro or 6,7,8,9-tetrahydro; and salts and complexes thereof.

The functionally modified carboxyl group in the meaning of the symbol R₀,
20 may be, for example, an esterified carboxyl group or a carbamoyl group optionally substituted on the nitrogen atom. More specifically one or both of R₀ represented by COR₆ in radical R_A and represented by COR₇ in radical R_B may independently represent carboxy, esterified carboxy, carbamoyl or substituted carbamoyl.

ACE inhibitors of the present invention may also include bicyclic
25 compounds and their derivatives disclosed in U.S. Patent No. 4,385,051 (the entire contents of which are incorporated herein by reference), and represented by formula (II) shown below:



wherein R_1 and R_2 are the same or different and each represents hydrogen, hydroxyl or lower alkoxy;

5 R_3 is hydrogen or lower alkyl;

R_4 is hydrogen, lower alkyl, amino-lower-alkyl or acylamino-lower-alkyl;


R_5 is hydrogen, lower alkyl or aralkyl which may be substituted;

R_6 is hydroxyl, lower alkoxy, amino or lower alkylamino;

and m and n each means 1 or 2, and salts thereof.

10

ACE inhibitors of the present invention may also include phosphinylalkanoyl substituted proline compounds disclosed in U.S. Patent No. 4,337,201 (the entire contents of which are incorporated herein by reference), and represented by formula (III) shown below:




15


and salts thereof, wherein R_1 is alkyl, aryl, arylalkyl, cycloalkyl, or cycloalkyl(alkyl);

R_2 and R_4 each is independently hydrogen, alkyl, arylalkyl or

20







wherein X is hydrogen, alkyl, or phenyl and Y is hydrogen, alkyl, phenyl or alkoxy, or together X and Y are $-(CH_2)_2-$, $-(CH_2)_3-$, $-CH=CH-$, or




R_3 is hydrogen or alkyl;

5 $-R_5-COOR_4$ is



R_6 is hydrogen, hydroxyl, alkyl, halogen, azido, amino, cycloalkyl, aryl, arylalkyl, carbamoyloxy;

10



N,N-dialkylcarbamoyloxy, or $-Z-R_9$;

15 R_7 and R_7 are the same and each is halogen or $-Z-R_{10}$, or R_7 and R_7 together are $=O$, $-O-(CH_2)_m-O-$ or $-S-(CH_2)_m-S-$;

R_8 is hydrogen and R_8 is phenyl, 2 hydroxyphenyl or 4-hydroxyphenyl or R_8 and R_8 together are $=O$;


R_9 is alkyl, aryl, arylalkyl, 1- or 2-naphthyl, or biphenyl;

R_{10} is alkyl, aryl or arylalkyl;

Z is oxygen or sulfur;


n is 0 or 1; and

5 m is 1 or 2; with the proviso that if $-R_5-COOR_4$ is




at least one of R_2 and R_4 is

10



ACE inhibitors of the present invention may also include azetidine-2-
 carboxylic acid derivative compounds disclosed in U.S. Patent No. 4,046,889 (the
 15 entire contents of which are incorporated herein by reference), and represented by
 formula (IV) shown below:



wherein R is hydroxy, NH_2 or lower alkoxy;

R_1 and R_4 each is hydrogen, lower alkyl or phenyl-lower alkyl;

20 R_2 is hydrogen or R_5-CO ;

R_3 is hydrogen, hydroxy or lower alkyl;


R_5 is lower alkyl, phenyl or phenyl-lower alkyl;

m is 1 to 3;

n is 0 to 2, and the asterisks indicate asymmetric carbon atoms. The carbon in the acyclic side chain is asymmetric when R_1 is other than hydrogen.

5

ACE inhibitors of the present invention may also include carboxyalkyl dipeptide compounds and derivatives thereof, disclosed in U.S. Patent No. 4,374,829 (the entire contents of which are incorporated herein by reference), and represented by formula (V) shown below:



15 wherein R and R_6 are the same or different and are hydroxy, lower alkoxy, lower alkenoxy, dilower alkylamino lower alkoxy (dimethylaminoethoxy), acylamino lower alkoxy (acetylaminomethoxy), acyloxy lower alkoxy (pivaloyloxymethoxy), aryloxy such as phenoxy, aryl(lower)alkoxy such as benzyloxy; substituted aryloxy or substituted aryl(lower)alkoxy wherein the substituent is methyl, halo or methoxy, amino, lower alkylamino, di(lower)alkylamino, hydroxyamino, or aryl(lower)alkylamino such as benzylamino;

20 R_1 is hydrogen, alkyl of from 1 to 20 carbon atoms which include branched and cyclic and unsaturated (such as allyl) alkyl groups, substituted lower alkyl wherein the substituent can be halo, hydroxy, lower alkoxy, aryloxy such as phenoxy, amino, dilower alkylamino, acylamino, such as acetamido and benzamidoaryl amino, guanidino, imidazolyl, indolyl, mercapto, lower alkylthio, arylthio such as phenylthio, carboxy or carboxyamido, carbolower alkoxy, aryl such as phenyl or naphthyl, substituted aryl such as phenyl wherein the substituent is
 25 lower alkyl, lower alkoxy or halo, aryl lower alkyl, aryl lower alkenyl, heteroaryl lower alkyl or heteroaryl lower alkenyl such as benzyl, styryl or indolyl ethyl, substituted aryl lower alkyl, substituted aryl lower alkenyl, substituted heteroaryl lower alkyl, or substituted heteroaryl lower alkenyl, wherein the substituent(s) is

halo, dihalo, lower alkyl, hydroxy, lower alkoxy, amino, aminomethyl, acylamino
(acetylamino or benzoylamino) dilower alkylamino, lower alkylamino, carboxyl,
halolower alkyl, cyano or sulfonamide, aryl lower alkyl or heteroaryl lower alkyl
substituted on the alkyl portion by amino or acylamino (acetylamino or
5 benzoylamino);

R_2 and R_7 are the same or different and are hydrogen or lower alkyl;


R_3 is hydrogen, lower alkyl, phenyl lower alkyl, aminomethyl phenyl lower
alkyl, hydroxy phenyl lower alkyl, hydroxy lower alkyl, acylamino lower alkyl
(such as benzoylamino lower alkyl, acetylamino lower alkyl) amino lower alkyl,
10 dimethylamino lower alkyl, halo lower alkyl, guanidino lower alkyl, imidazolyl
lower alkyl, indolyl lower alkyl, mercapto lower alkyl, lower alkyl thio lower alkyl;

R_4 is hydrogen or lower alkyl;

R_5 is hydrogen, lower alkyl, phenyl, phenyl lower alkyl, hydroxyl phenyl
lower alkyl, hydroxyl lower alkyl, amino lower alkyl, guanidino lower alkyl,
15 imidazolyl lower alkyl, indolyl lower alkyl, imidazolyl lower alkyl, indolyl lower
alkyl, mercapto lower alkyl or lower alkyl thio lower alkyl;

R_4 and R_5 may be connected together to form an alkylene bridge of from 2 to
4 carbon atoms, an alkylene bridge of from 2 to 3 carbon atoms and one sulfur atom,
an alkylene bridge of from 3 to 4 carbon atoms containing a double bond or an
20 alkylene bridge as above substituted with hydroxyl, lower alkoxy, lower alkyl or
dilower alkyl.

ACE inhibitors of the present invention may also include substituted
iminodiacid compounds as disclosed in U.S. Patent No. 4,508,729 (the entire
25 contents of which are incorporated herein by reference), and represented by formula
(VI) shown below:




wherein the ring A is saturated and n=0 or 1, or the ring A is a benzene ring and n=1,

R₁ represents a lower alkyl group having from 1 to 4 carbon atoms which can carry an amino group,

R₂ represents a hydrogen atom or an alkyl group having from 1 to 4 carbon atoms,

R₃ represents a straight or branched alkyl group, a mono- or dicycloalkylalkyl or phenylalkyl group having no more than a total of 9 carbon atoms, or a substituted alkyl group of the formula:




with R₄ = H, a lower alkyl (C₁ to C₄) or a cycloalkyl (C₃ to C₆) group,

R₅=H, a lower alkyl (C₁ to C₄), a cycloalkyl (C₃ to C₆) or an alkoxy carbonyl group,

Y=S or >N-Q where Q=H, or an acetyl or benzyloxycarbonyl group, and p=1 or 2, and q=0 or 1.

The ACE inhibitor compounds of the present invention may also include substituted acyl derivatives of 1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid compounds disclosed in U.S. Patent No. 4,344,949 (the entire contents of which are incorporated herein by reference), and having formula (VII) as shown below:



where R is hydrogen, lower alkyl or aralkyl;

R₁ is hydrogen, lower alkyl, or benzyl;


R₂ is hydrogen or lower alkyl, and Ar is phenyl or phenyl substituted with 1
5 or 2 substituents selected from the group consisting of fluorine, chlorine, bromine,
lower alkyl, lower alkoxy, hydroxy or amino;

X and Y are independently hydrogen, lower alkyl, lower alkoxy, lower
alkylthio, lower alkylsulfinyl, lower alkylsulfonyl, hydroxy, or X and Y together are
methylenedioxy;

10 m is 0 to 3;

and pharmaceutically acceptable salts thereof.

The ACE inhibitors may also include phosphinylalkanoyl proline compounds
disclosed in U.S. 4,168,267 (the entire content of which is incorporated herein by
15 reference), which have formula (VIII) as shown below:



wherein R₁ is lower alkyl, phenyl or phenyl-lower alkyl;


R₂ is hydrogen, phenyl-lower alkyl or a metal ion;

R_3 is hydrogen or lower alkyl;

R_4 is hydrogen, lower alkyl, phenyl-lower alkyl or a metal ion; and

n is 0 or 1.

5 The ACE inhibitor compounds of the present invention may also include ether and thioether mercaptoacyl proline compounds disclosed in U.S. Patent No. 4,316,906 (the entire content of which is incorporated herein by reference), and having formula (IX) as shown below:



10 wherein the group $X-R_1$ is located at the 3- or 4-position in the ring;


X is oxygen or sulfur;

R is hydrogen or lower alkyl;

R_1 is lower alkyl, lower alkenyl, lower alkynyl, cycloalkyl, 1- or 2-adamantyl, aryl, substituted aryl, phenyl-lower alkylene or substituted phenyl-lower alkylene.

R_2 and R_3 are independently selected from hydrogen, lower alkyl, and trifluoromethyl;

R_4 is hydrogen, $R_5 - CO -$, or




20

R_5 is lower alkyl, phenyl, phenyl-lower alkylene; substituted phenyl, or substituted phenyl-lower alkylene;

n is 0, 1 or 2; and salts thereof.


The ACE inhibitor compounds of the present invention also may include proline derivatives and related compounds disclosed in U.S. Patent No. 4,105,776 (the entire contents of which are incorporated herein by reference), which have the general formula (X) as shown below:



wherein R is hydroxy, NH₂ or lower alkoxy;

R₁ and R₄ each is hydrogen, lower alkyl, phenyl or phenyl-lower alkyl;


R₂ is hydrogen, lower alkyl, phenyl, substituted phenyl wherein the phenyl substituent is halo, lower alkyl or lower alkoxy, phenyl-lower alkyl, diphenyl-lower alkyl, triphenyl-lower alkyl, lower alkylthiomethyl, phenyl-lower alkythiomethyl, lower alkanoyl-amidomethyl,



R₃ is hydrogen, hydroxyl or lower alkyl;

R₅ is lower alkyl, phenyl or phenyl-lower alkyl;

R₆ is lower alkyl, phenyl, substituted phenyl, (wherein the phenyl substituent is halo, lower alkyl or lower alkoxy), hydroxyl-lower alkyl or amino(carboxy)lower alkyl;




M is O or S;

m is 1 to 3; and

n and *p* each is 0 to 2.

- 5 The asterisks indicate asymmetric carbon atoms. Each of the carbons bearing a substituent R₁, R₃ and R₄ is asymmetric when that substituent is other than hydrogen.

- 10 The ACE inhibitors of the present invention may also include bicyclic pyridazo [1,2-A][1,2] diazepine compounds disclosed in U.S. Patent No. 4,512,924 (the entire contents of which are incorporated herein by reference), having the general formula (XI) as shown below:




- 15 wherein B represent a methylene (—CH₂—) , ethylene (—CH₂—CH₂—) or vinylene (—CH=CH—) group;

- 20 R₁ represents a hydrogen atom or an alkyl, aralkyl, amino-alkyl, monoalkylamino-alkyl, dialkylaminoalkyl, acylamino-alkyl, phthalimido-alkyl, alkoxycarbonylamino-alkyl, aryloxy carbonylamino-alkyl, aralkoxycarbonylamino-alkyl, alkylaminocarbonylaminoalkyl, arylaminocarbonylamino-alkyl,

aralkylaminocarbonylamino-alkyl, alkylsulphonylamino-alkyl or arylsulphonylamino-alkyl group;

R₂ represents a carboxyl, alkoxycarbonyl or aralkoxycarbonyl group or a group of the formula:

5




R₃ represents a carboxyl, alkoxycarbonyl or aralkoxycarbonyl group;

R₄ and R₅ each represent a hydrogen atom or R₄ and R₅ together represent an oxo group;

10 R₆ and R₇ each represent a hydrogen atom or an alkyl or aralkyl group or R₆ and R₇ together with the nitrogen atom to which they are attached represent a saturated 5 membered or 6-membered heteromonocyclic ring which may contain a further nitrogen atom or an oxygen or sulphur atom, and n stands for zero, 1 or 2, and pharmaceutically acceptable salts thereof.

15

The ACE inhibitors may also include pyroglutamic acid derivatives as disclosed in U.S. Patent No. 4,234,489 (the entire contents of which are incorporated herein by reference), having the formula (XII) as shown below:




20 and salts thereof, wherein

R is hydrogen, alkyl or diphenylmethyl;

R₁ is hydrogen, alkyl or trifluoromethyl;

R₂ is hydrogen,



5


R₃ is hydrogen, alkyl, phenyl, or phenylalkyl;

X is oxygen or sulfur; and

n is 0 or 1.







10

The ACE inhibitor compounds may also include the phosphoramidate substituted amino or imino acids and salts thereof of U.S. Patent No. 4,432,971 (the entire contents of which are incorporated herein by reference), and of the formula (XIII) as shown below:







15



wherein X is an imino or amino acid of the formula:




5




10



R₇ is hydrogen, lower alkyl, halogen, keto, hydroxyl,




5




10

a 1- or 2-naphthyloxy of the formula:




15



or a 1- or 2-naphthylthio of the formula:



5 ---O--- lower alkyl, a 1- or 2-naphthyloxy of the formula:




10 or a 1- or 2-naphthylthio of the formula:



15

R_{10} is halogen or ---Y---R_{16} ,

R_{11} , R_{11} , R_{12} and R_{12} are independently selected from hydrogen and lower alkyl or R_{11} , R_{12} and R_{12} are hydrogen and R_{11} is



5 R_{13} is hydrogen, lower alkyl of 1 to 4 carbons, lower alkythio of 1 to 4 carbons, chloro, bromo, fluoro, trifluoromethyl, hydroxyl, phenyl, phenoxy, phenylthio, or phenylmethyl.

R_{14} is hydrogen, lower alkyl of 1 to 4 carbons, lower alkoxy of 1 to 4 carbons, lower alkylthio of 1 to 4 carbons, chloro, bromo, fluoro, trifluoromethyl, or hydroxy.


10 m is zero, one two or three;

p is one, two or three provided that p is more than one only if R_{13} or R_{14} is hydrogen, methyl, methoxy, chloro, or fluoro;

R_{15} is hydrogen or lower alkyl of 1 to 4 carbons;

Y is oxygen or sulfur;


15 R_{16} is lower alkyl of 1 to 4 carbons;




or the R_{16} groups join to complete an unsubstituted 5- or 6-membered ring or said ring in which one or more of the carbons has a lower alkyl of 1 to 4 carbons or a di(lower alkyl of 1 to 4 carbons) substituent.

20 R_4 is hydrogen, lower alkyl, $-(CH_2)_m-$ cycloalkyl,

or



R_5 is hydrogen, lower alkyl,




10





r is an integer from 1 to 4.

R_1 is hydrogen, lower alkyl, or cycloalkyl.

R_2 is hydrogen, lower alkyl, halo substituted lower alkyl,


15






or R_1 and R_2 taken together are $\text{---}(\text{CH}_2)_n\text{---}$ wherein n is an integer from 2 to 4.

10 R_3 and R_6 are independently selected from hydrogen, lower alkyl, benzyl, benzhydryl, or



15 wherein R_{17} is hydrogen, lower alkyl, cycloalkyl, or phenyl, and R_{18} is hydrogen, lower alkyl, lower alkoxy, phenyl, or R_{17} and R_{18} taken together are










R_{19} is lower alkyl, benzyl, or phenethyl;

R_{20} is hydrogen, lower alkyl, benzyl or phenethyl;

R_{21} is alkyl of 1 to 10 carbons;

5




10

wherein q is zero or an integer from 1 to 7, s is an integer from 1 to 8, and R_{13} , and p is as defined above.







15

The ACE inhibitor compounds of the present invention may also include the phosphonate substituted amino or imino acids and salts thereof, as disclosed in U.S. Patent No. 4,452,790 (the entire contents of which are incorporated herein by reference), of the general formula (XIV) as shown below:







20



X is an imino or amino acid of the formula:





5




10




R₄ is hydrogen, lower alkyl, halogen, keto, hydroxy,






5


a 1- or 2-naphthyl of the formula:



10




a 1- or 2-naphthyloxy of the formula:




15


or a 1- or 2-naphthylthio of the formula:



R₅ is keto, halogen,




5 ---O--- lower alkyl, a 1- or 2-naphthyloxy of the formula:




---S--- lower alkyl,

10




or a 1- or 2-naphthylthio of the formula:



15


R₇ is keto or



20

each R₈ is independently halogen or ---Y---R_{14} ;

R₉, R₉, R₁₀ are independently selected from hydrogen and lower alkyl or R₉,
R₁₀ and R₁₀ are hydrogen and R₉ is



5 R_{11} is hydrogen, lower alkyl of 1 to 4 carbons, lower alkoxy of 1 to 4 carbons, lower alkythio of 1 to 4 carbons, chloro, bromo, fluoro, trifluormethyl, hydroxyl, phenyl, phenoxy, phenylthio, or phenylmethyl;

R_{12} is hydrogen, lower alkyl of 1 to 4 carbons, lower alkoxy of 1 to 4 carbons, lower alkythio of 1 to 4 carbons, chloro, bromo, fluoro, trifluoromethyl, or hydroxy;


10 m is zero, one two or three;

p is one, two or three provided that p is more than one only if R_{11} or R_{12} is hydrogen, methyl, methoxy, chloro, fluoro;

R_{13} is hydrogen or lower alkyl of 1 to 4 carbons;


Y is oxygen or sulfur;

15 R_{14} is lower alkyl of 1 to 4 carbons;













20 or the R_{14} groups join to complete an unsubstituted 5- or 6-membered ring or said ring in which one or more of the carbons has a lower alkyl of 1 to 4 carbons or a di(lower alkyl of 1 to 4 carbons) substituent;

R_{21} is hydrogen, lower alkyl, cycloalkyl, phenyl, or






R_{22} is hydrogen, lower alkyl,





5

r is an integer from 1 to 4;



R₁ is alkyl of 1 to 10 carbons, aminoalkyl, haloalkyl;





wherein q is zero or an integer from 1 to 7 and R₁₂ and p are defined as above;


- 5 R₁₉ and R₂₀ are independently selected from hydrogen, lower alkyl, halo substituted lower alkyl;



10


wherein m, R₁₁, and p are as defined above;

R₂ is hydrogen, lower alkyl, halo substituted lower alkyl;




5 wherein r is defined above.

R_3 and R_6 are independently selected from hydrogen, lower alkyl, benzyl, alkali metal such as Li, Na or K, benzhydryl, or




10 wherein R_{15} is hydrogen, lower alkyl, cycloalkyl, or phenyl, and R_{16} is hydrogen, lower alkyl, lower alkoxy, phenyl, or R_{15} and R_{16} taken together are $\text{---}(\text{CH}_2)_2\text{---}$, $\text{---}(\text{CH}_2)_3\text{---}$, ---CH=CH--- , or



R_{17} is lower alkyl, benzyl, or phenethyl;

R_{18} is hydrogen, lower alkyl, benzyl or phenethyl.

The ACE inhibitor compounds may also include the compounds disclosed in EP Patent No. 0060668 (the entire contents of which are incorporated herein by reference), having formula (XV) as shown below:



5 or a pharmaceutically acceptable salt thereof, wherein

m is 0 to 3;

n is 1 to 5;

R_1 is hydrogen or C_{1-6} alkyl;

R_2 is hydrogen, C_{1-4} alkyl, $-(CH_2)_p-NH_2$

10 wherein p is 1 to 4, or $-NHCOR_5$ wherein R_5 is C_{1-4} alkyl;

R_3 is hydrogen or C_{1-6} alkyl;


R_4 is C_{1-4} alkyl, C_{1-4} alkoxy, halogen or CF_3 ; and

X is CH_2 or S ;

The dihydrobenzofuranyl moiety may be bonded to the rest of the structure
15 at the 2- or 3-position, preferably, the 2-position;

Preferably X is CH_2 .

The ACE inhibitor compounds may also be compounds as disclosed in EP
Patent No. 0080822 (the entire contents of which are incorporated herein by
20 reference), having formula (XVI) as shown below:



or a pharmaceutically acceptable salt thereof,


wherein R₁ is C₁₋₅ alkyl optionally substituted by NHR₆, (wherein R₆ is hydrogen or C₁₋₅) alkylcarbonyl) or by phenyl or naphthyl optionally substituted by halogen, C₁₋₅ alkyl or C₁₋₅ alkoxy or by dihydrobenzofuran-2-yl, optionally substituted in the benzo moiety by C₁₋₅ alkyl, C₁₋₅ alkoxy, halogen or trifluoromethyl;

R₂ and R₅ are the same or different and each is hydroxyl, C₁₋₅ alkoxy, C₂₋₆ alkylcarbonyl or amino optionally substituted by C₁₋₅ alkyl;

R₃ is C₁₋₅ alkyl optionally substituted by the group -NHR₇, wherein R₇ is hydrogen, C₁₋₅ alkyl or C₂₋₆ alkylcarbonyl; and

R₄ is phenyl optionally substituted by halogen, C₁₋₅ alkoxy, trifluoromethyl or C₁₋₅ alkyl.

The ACE inhibitor compounds of the present invention may also include phosphory aminoacid derivators as disclosed in EP Patent No. 0009183 (the entire contents of which are incorporated herein by reference), and having the general formula (XVII) as shown below:



wherein n is 0 or 1;

R is hydrogen, lower alkyl, phenyl lower alkyl, hydroxy phenyl lower alkyl, hydroxy lower alkyl, amino lower alkyl, guanidino lower alkyl, imidazol lower alkyl, indolyl lower alkyl, mercapto lower alkyl, lower alkyl mercapto lower alkyl;

R₃ is hydrogen;

5 R₄ is hydrogen, lower alkyl, phenyl lower alkyl, hydroxyl phenyl lower alkyl, hydroxyl lower alkyl, aminolower alkyl, guanidino lower alkyl, guanidino lower alkyl, imidazol lower alkyl, indolyl lower alkyl, mercapto lower alkyl, lower alkyl mercapto lower alkyl;

10 R₃ and R₄ may be connected together to form an alkylene bridge of from 2 to 4 carbon atoms or an alkylene bridge of from 2 to 3 carbon atoms one sulfur atom;


X is 0, NR₅, S where R₅ = H or lower alkyl;

R₁ is hydrogen, lower alkyl, aralkyl or aryl;

R₂ is hydrogen, lower alkyl, aralkyl or aryl and pharmaceutically acceptable salts thereof.

15


The ACE inhibitors may also include the carbamate derivatives of mercaptoacyl hydroxyl prolines as disclosed in U.S. Patent No. 4,217,359 (the entire contents of which are incorporated herein by reference), and which have the formula (XVIII) as shown below:



wherein R, R₂ and R₃ each is hydrogen or lower alkyl;

R₀ and R₁ each is hydrogen, lower alkyl, cyclo-lower alkyl, allyl, propargyl, phenyl or substituted phenyl; or R₀ and R₁ can join with the nitrogen to form a 5- or 6-membered heterocyclic;

25 R₄ is hydrogen or hydrolysable organic protecting group of the formula R₅-CO- or




R₅ is lower alkyl, phenyl, substituted phenyl, phenyl-lower alkyl, substituted
5 phenyl-lower alkyl, cycloalkyl, thienyl, or furyl;

n is 0, 1 or 2; and salts thereof.

The asterisks indicate centers of asymmetry. The carbon in the acyclic side
chain is asymmetric when R₂ and/or R₃ is other than hydrogen. Each of the centers
of asymmetry provide D and L forms which can be separated by conventional
10 methods as described below.

The ACE inhibitors of the present invention also may include substituted
acyl derivatives of amino acids disclosed in U.S. Patent No. 4,129,571 (the entire
contents of which are incorporated herein by reference), and which have the general
15 formula (XIX) as shown below:




and salts thereof, wherein:

R is hydroxyl or lower alkoxy;

R₁ is hydrogen, lower alkanoyl or amino(imino)-methyl;

20 R₂ is hydrogen, lower alkyl or phenyl-lower alkylene;

R₃ is hydrogen, lower alkanoyl, benzoyl or



A is hydrogen, lower alkyl or hydroxyl-lower alkylene;

B is hydrogen, lower alkyl, phenyl, phenyl-lower alkylene, hydroxyl-lower
 5 alkylene, hydroxyphenyl-lower alkylene, amino-lower alkylene, guanidino-lower
 alkylene, mercapto-lower alkylene, lower alkyl-thio-lower alkylene, imidazolyl-
 lower alkylene, indolyl-lower alkylene, carbamoyl-lower alkylene or carboxy-lower
 alkylene;

or A and B together form a $(\text{CH}_2)_p$ bridge which completes a ring of 5 or 6
 10 atoms with the nitrogen and carbon to which they are joined, one carbon optionally
 bearing a hydroxy group;


n is 0 or 1;

m 0, 1, 2, 3 or 4; at least one of m and n is other than 0; and

p is 3 or 4.


15 The asterisks denote centers of asymmetry.

ACE inhibitors may also include halogenated compounds as disclosed in
 U.S. Patent No. 4,154,935 (the entire contents of which are incorporated herein by
 reference), which have the general formula (XX) as shown below:



20

wherein R is hydrogen, lower alkanoyl or



R₁ is hydrogen or lower alkyl;

R₂ and R₂' each independently represent hydrogen or halogen;

R₃ and R₄ each independently represent hydrogen, lower alkyl or


- 5 trifluoromethyl, not more than one being trifluoromethyl, and at least one of R₂, R₂', R₃ or R₄ is a halogen or trifluoromethyl substituent represented by the named symbol as defined above;

m is 2; and

n is 0 or 1.

- 10 The asterisks indicate asymmetric carbon atoms.

- ACE inhibitor compounds may also include carboxyalkylacylamino acids and related compounds disclosed in U.S. independently represent 4,052,511 (the entire contents of which are incorporated herein by reference), which are derivatives of proline, pipecolic acid, azetidine-2-carboxylic acid and which have the general formula (XXI) as shown below:
- 15



wherein

R is hydroxy, amino or lower alkoxy;

- 20 R₁ and R₄ each is hydrogen, lower alkyl or phenyl-lower alkyl;

R₂ is hydroxy, amino, hydroxyamino or lower alkoxy;


R₃ is hydrogen, hydroxy or lower alkyl;

m is 1 to 3;

n is 0 to 2.

The asterisks indicate asymmetric carbon atoms. The carbons in the acyclic side chain are asymmetric when R₁ or R₄ are other than hydrogen.


The ACE inhibitor compounds may also include the dehydrocyclicimino acid compounds disclosed in U.S. independently represent 4,129,566 (the entire contents of which are incorporated herein by reference), which have the general formula (XXII) as shown below:



wherein:

R and R₂ each is hydrogen or lower alkyl;


R₁ is hydrogen, lower alkanoyl or



m and n each is 0 or 1.

The asterisks indicate asymmetric carbon atoms. The carbon in the acyclic side chain is asymmetric when R₁ is other than hydrogen.

The ACE inhibitor compounds may also include compounds disclosed in U.S. independently represent 4,053,651 (the entire contents of which are incorporated herein by reference), having the formula (XXIII) as shown below:




5 or a salt thereof, wherein

R₁ is hydrogen, lower alkyl, phenyl-lower alkylene, hydroxyl-lower alkylene, amino-lower alkylene, guanidino-lower alkylene, imidazolyl-lower alkylene, indolyl-lower alkylene, mercapto-lower alkylene, lower alkylmercapto-lower alkylene, carbamoyl-lower alkylene or carboxy-lower alkylene;

10 R₂, R₃ and R₄ each is hydrogen, lower alkyl or phenyl-lower alkylene;


R₅ is hydrogen, lower alkanoyl, benzoyl or



15 The asterisks denote centers of asymmetry.

ACE inhibitors may also include derivatives of mercaptoacyl prolines and pipercolic acids as disclosed in U.S. independently represent 4,311,697 (the entire contents of which are incorporated herein by reference), of the formula (XXIV) as shown below:


20



R and R₆ are independently selected from hydrogen and lower alkyl provided that R₆ is lower alkyl only if R₃ is also lower alkyl;


R₃ and R₄ are independently selected from hydrogen, lower alkyl, lower alkylthio, $-(CH_2)_n-SH$, and halo substituted lower alkyl;

X₁, X₂ and X₃ are independently selected from lower alkyl, lower alkenyl, lower alkynyl, cycloalkyl, halo substituted lower alkyl, hydroxyl substituted lower alkyl;




or R₁ and R₂ join in a polymethylene chain to complete an unsubstituted or substituted 5- or 6-membered ring.

When R₁ and R₂ are joined together in a polymethylene chain of 2 or 3 carbons, these cyclic ketal and thioketals can be represents as follows:




wherein t is 2 or 3 and R₉ and R₁₀ are both hydrogen, both lower alkyl, or one is hydrogen and the other is lower alkyl, halo substituted lower alkyl, hydroxyl substituted lower alkyl,



preferably, only one carbon of the polymethylene chain will be substituted.

R_7 is hydrogen, lower alkyl of 1 to 4 carbons, especially methyl, lower alkoxy of 1 to 4 carbons, especially methyl, lower alkoxy of 1 to 4 carbons, especially methoxy, lower alkythio 1 to 4 carbons, especially methylthio, chloro, bromo, fluoro, trifluoromethyl, or hydroxy;

R_5 is hydrogen, a hydrolyzably removable protecting group, a chemically removable protecting group, or when R_3 and R_4 are other than $-(CH_2)_m-SH$ a sulfide of the formula



m is zero, one or two;


n is one, two or three;

p and q are each one or two provided that both are not two.

The asterisk in the above formula indicates a center of asymmetry in the ring. In the case of praline, i.e., p and q are both one, this center is in the L-configuration. In the case of pipecolic acid, i.e., one of p and q is two, this center is in the D, L or L-configuration.

Asymmetric centers can also be present in the mercaptoacyl sidechain depending upon the definition of R_3 , R_4 and R_6 . Another asymmetric center may also be present in the ring when X_1-R_1 and X_2-R_2 are different. The products can accordingly exist in stereoisomeric forms or as racemic mixtures thereof.

The ACE inhibitors may also include the imido, amido and amino derivative compounds of mercaptoacyl prolines and pipecolic acids disclosed in U.S. Patent No. 4,310,461 (the entire contents of which are incorporated herein by reference),
 5 and having the formula (XXV) as shown below:



and salts thereof, and the symmetrical dimer thereof, wherein

R_1 and R_2 are the same or different and are hydrogen, alkyl, cycloalkyl, 1-adamantyl, aryl, arylalkyl, alkylcarbonyl, arylcarbonyl, arylalkylcarbonyl,
 10 alkysulfonyl, arylsulfonyl, or arylvinylcarbonyl ($\text{aryl}-\text{CH}=\text{CH}-\text{CO}-$), or together with the nitrogen atom to which they are attached R_1 and R_2 are 1-pyrrolidinyl, 1-piperidinyl, 1-homopiperidinyl, 4-morpholinyl, 4-alkyl-1-piperazinyl, 4-aryl-1-piperazinyl, 1-imidazolyl, 1-pyrrolidinyl-2,5-dione(succinimido), 3-alkyl-1-pyrrolidinyl-2,5-dione, 3-aryl-1-pyrrolidinyl-2,5-dione, 1-piperidinyl-2,6-dione, 2H-isoindol-2-yl-1,3-dione(phthalimido), hexahydro-2H-isoindol-2-yl-1,3-dione(hexahydrophthalimido), 2,5-dihydro-2,5-dioxo-1H-pyrrol-1-yl(maleimido),
 15 1,1,3-trioxo-1,2-benziso-thiazol-2(3H)-yl(2-saccharinyl), or 1,3-dihydro-1,3-dioxo-2H-benz[de]isoquinolin-2-yl(1,8-naphthalenedicarboximido);


R_3 is hydrogen, alkyl, aryl, arylalkyl, or a hydrolyzable acyl protecting group
 20 such as alkanoyl or arylcarbonyl;

R_4 is hydrogen, alkyl, alkythio or trifluoromethyl;

R_5 is hydrogen, alkyl, or arylalkyl;

n is 0, 1 or 2; and p is 1 or 2.

The ACE inhibitors may also include phosphonoacyl prolines and related compounds as disclosed in U.S. Patent No. 4,151,172 (the entire contents of which are incorporated herein by reference), which have the formula (XXVI) as shown below:




R₁ and R₂ each is hydrogen, lower alkyl, lower alkenyl, unsubstituted or substituted phenyl-lower alkyl or a metal ion;

R₃ is hydrogen or lower alkyl;

R₄ is hydrogen, lower alkyl, phenyl-lower alkyl or a metal ion; and



n is 0 or 1.


The ACE inhibitor compounds may also include mercaptoacyl derivatives of various 4-cis substituted prolines and salts thereof as disclosed in U.S. Patent No. 4,316,905 (the entire contents of which are incorporated herein by reference), and which have the formula (XXVII) as shown below:



R represents hydrogen or lower alkyl;

R₁ represents -(CH₂)_m-cycloalkyl, 1-cyclohexenyl, 1,4-cyclohexadienyl;






R_2 and R_3 are independently selected from hydrogen, lower alkyl, lower alkylthio and halo substituted lower alkyl;

5 n is zero, one or two;

R_4 is hydrogen, a hydrolyzably removable protecting group, a chemically removable protecting group, or



m is zero, one, two or three;

10 R_5 is hydrogen, lower alkyl of 1 to 4 carbons, especially methyl, lower alkoxy of 1 to 4 carbons, especially methoxy, lower alkylthio of 1 to 4 carbons, especially methylthio, chloro, bromo, fluoro, trifluoromethyl, hydroxy, phenyl, phenoxy, phenylthio, or phenylmethyl. The hydroxy substituted compounds are obtained by heating the corresponding methoxy substituted compound with pyridine
15 HCl;

q is one, two or three provided that q is more than one only if R_5 is hydrogen, methyl, methoxy, chloro or fluoro;

X is oxygen or sulfur.

20 The ACE inhibitors also preferably include: alylmercapto and mercaptoalkanoyl prolines (U.S. Patent No. 4,046,889) such as captopril (1-[(2S)-3-mercapto-2-methylpropionyl]-L-proline) (U.S. Patent No. 4,105,776) and ether or thioether mercaptoacyl prolines such as zofenopril (U.S. Patent No. 4,316,906);

carboxyalkyl dipeptides such as enalapril (N-(1-ethoxycarbonyl-3-phenylpropyl)-L-ananyl-L-proline) (U.S. Patent No. 4,374,829), lisinopril (U.S. Patent No. 4,374,829), quinapril (U.S. Patent No. 4,344,949), and ramipril (U.S. Patent No. 4,508,729); carboxyalkyl dipeptide mimics such as cilazapril (U.S. Patent No. 4,512,924) and benazapril (U.S. Patent No. 4,410,520); phosphinylalkanoyl prolines (U.S. Patent No. 4,168,267) such as fosinopril (U.S. Patent No. 4,337,201) and trandolopril; phosphonamidate substituted amino or imino acids (U.S. Patent No. 4,432,971); phosphonate substituted amino or imino acids and salts thereof, including ceronapril ((S)-1-[6-amino-2-[[hydroxyl(4-phenylbutyl)phosphinyl]oxy]-1-oxohexyl]-L-proline) (U.S. Patent No. 4,452,790); Beecham's BRL 36,378 (EP Nos. 80822 and 60668); Chugai's MC-838 (disclosed in CA 102:72588v and Jap.J.Pharmacol. 40:373 (1986)); Ciba-Geigy's CGS 14824 (3-([1-ethoxycarbonyl-3-phenyl-(1S)-propyl]-amino)-2,3,4,5-tetrahydro-2-oxo-1-(3S)-benzazepine-1 acetic acid HCL) (U.K. Patent No. 2103614) and CGS 16,617 (3(S)-[[(1S)-5-amino-1-carboxypentyl]amino]2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-ethanoic acid) (U.S. Patent No. 4,473,575); Cetapril (alacepril, Dainippon) (Eur.Therap.Res. 39:671 (1986); 40:543 (1986)); Ru 44570 (Hoechst) (Arzneimittelforschung 35:1254 (1985)); Cilazapril (Hoffman-LaRoche) (J.Cardiovasc.Pharmacol. 9:39 (1987); Ro 31-2201 (Hoffman-LaRoche) (FEBS Lett. 165:201 (1984); Lisinopril (Merck) (Curr.Therap.Res. 37:342 (1985) and Eur. Patent App. No. 12-401); Indalapril (delapril) (U.S. Patent No. 4,385,051); Rentiapril (fentiapril, Santen) (Clin.Exp. Pharmacol. Physiol. 10:131 (1983); Indolapril (Schering) (J.Cardiovasc.Pharmacol. 5:643, 655 (1983)); Spirapril (Schering) (Acta.Pharmacol.Toxicol. 59 (Supp. 5):173 (1986); Perindopril (Servier) (Eur.J.Clin.Pharmacol. 31:519 (1987); Quinapril (Warner-Lambert) (U.S. Patent No. 4,344,949); CI 925 (Warner-Lambert) ([3S-[2[R(*)R(*)]3R(*)]-2-[2-[1-(ethoxycarbonyl)-3-phenylpropyl]amino]-1-oxopropyl]-1,2,3,4-tetrahydro-6,7-dimethoxy-3-isoquinolinecarboxylic acid HCL) (Pharmacologist 26:243, 266 (1984)); WY-44221 (Wyeth) (J.Med.Chem. 26:394 (1983)); Mercapto containing compounds such as pivopril and YS980 (U.S. Patent No. 6,127,370); Omapatrilat (Drugs R.D. 1999 1(4):350-1) (U.S. Patent No. 6,300,503); Alacepril (Jap. Patent App. No. 78/82809); moveltopril; quinaprilat; moexipril; perinodpril (S-9490) (Annual Drug Data Report

7, 99 (1985)); pentopril; ancovenin (Annual Drug Data Report 6, 20 (1984)); phenacein (Annual Drug Data Report 7, 20 (1985)); and nicotianamin (East German Patent DD 226 880). The contents of all cited patents, applications, publications are hereby incorporated herein by reference.

5 The most preferred ACEI is Enalapril, i.e. (N-(1-ethoxycarbonyl-3-phenylpropyl)-L-ananyl-L-proline), (U.S. Patent No. 4,374,829, entire contents incorporated herein by reference) and other similar or derivative carboxyalkyl dipeptides. Other preferred ACE inhibitors include lisinopril or captopril.

(iii) Angiotensin II Receptor Blockers / Antagonists

10 Angiotensin is formed from a precursor, angiotensinogen, which is produced by the liver and found in the alpha-globulin fraction of plasma. The lowering of blood pressure is a stimulus to secretion of renin by the kidney into the blood. Renin cleaves from angiotensinogen a terminal decapeptide, angiotensin I. This is further altered by the enzymatic removal of a dipeptide, by Angiotensin Convertin Enzyme
15 (ACE), to form angiotensin II. Angiotensin II is a potent regulator of blood pressure and of water and electrolyte balance. Angiotensin II interacts with two pharmacologically distinct subtypes of cell surface receptors, types 1 and 2. Whereas AGTR1, the type-1 receptor for Angiotensin II, mediates the vasopressive and aldosterone-secreting effects of angiotensin II, the function of the type-2
20 Angiotensin receptor (AGTR2) was relatively unclear, although it is expressed in both adult and embryonic life. Recent evidence indicates that the type-2 Angiotension receptor is not required for embryonic development, but plays a role in the central nervous system and cardiovascular functions that are mediated by the renin-angiotensin system (Hein et al., *Nature* 377: 744-748, 1995).

25 Ichiki et al. (*Nature* 377: 748-750, 1995) reported the unexpected finding that the targeted disruption of the mouse AGTR2 gene resulted in a significant increase in blood pressure and increased sensitivity to the pressor action of angiotensin II. The authors concluded that the type 2 receptor mediates a depressor effect and antagonizes the AGTR1-mediated pressor action of angiotensin II. In
30 addition, disruption of the AGTR2 gene attenuated exploratory behavior and lowered blood pressure. Their results indicated that angiotensin II activates AGTR1

and AGTR2, which have mutually counteracting hemodynamic effects, and that AGTR2 regulates central nervous system functions, including behavior. Ichiki et al. (1995) commented on the fact that Hein et al. (1995) did not find an increase in basal blood pressure and they suggested that this could be due to differences in genetic background of the mice studied.

Angiotensin-II receptor antagonists act by binding to specific membrane-bound receptors that displace Angiotensin II from its type 1-receptor subtype (AGTR1). These drugs therefore function as selective blockers. AT-II pressor effects are mediated by AGTR1. Unlike angiotensin-converting enzyme inhibitors, they do not inhibit bradykinin metabolism or enhance prostaglandin synthesis. Angiotensin-II receptor antagonists are well tolerated. Cough occurs much less often with these agents than with angiotensin-converting enzyme inhibitors, and they do not adversely affect lipid profiles or cause rebound hypertension after discontinuation. There has been a rapid growth in members of this new class of drugs. The angiotensin-II receptor antagonists that have been labeled for use in hypertension by the U.S. Food and Drug Administration (FDA) are Losartan (Cozaar[®]), Valsartan (Diovan[®]), Irbesartan (Avapro[®]), Candesartan (Atacand[®]) and Telmisartan (Micardis[®]). Other angiotensin-II receptor antagonists currently under investigation include tasosartan, zolarsartan, Teveten (eprosartan mesylate). At the present time four are being actively marketed in Canada: Losartan (Cozaar[®]), Valsartan (Diovan[®]), Irbesartan (Avapro[®]), Candesartan (Atacand[®]).

Losartan. Losartan (U.S. Pat. No. 5,153,197) was the first angiotensin-II receptor antagonist to be introduced (1995). Compared with the parent drug, the active metabolite (EXP3174) has a longer half-life and antihypertensive effects that correlate more with plasma concentration. Double-blind studies have shown that losartan is well tolerated and as efficacious as enalapril and nifedipine for lowering blood pressure. The mean blood pressure reduction achieved with losartan in a dosage of 50 to 150 mg once daily is 5.5 to 10.5 mm Hg for systolic pressure and 3.5 to 7.5 mm Hg for diastolic pressure. One review of the efficacy and safety of losartan in the treatment of essential hypertension indicated a slowly developing

response, with blood pressure becoming lower over several weeks of continued treatment.

5 The starting dosage of losartan is 50 mg once daily. The duration of activity for a dose is 24 hours. Twice-daily dosing can be used if the antihypertensive effect measured at a trough is inadequate. However, a comparison of losartan in dosages of 100 mg once daily and 50 mg twice daily showed no significant difference in antihypertensive efficacy.

10 A hydrochlorothiazide-losartan combination (Hyzaar) is also available. This combination drug contains 12.5 mg of hydrochlorothiazide and 50 mg of losartan. Some investigators advocate the use of this combination instead of escalation of a single drug, because dose-dependent adverse effects are less likely to occur. Dosing is once or twice daily.

15 **Valsartan.** (U.S. Pat. Nos. 5,399,578; 6,294,197) Placebo-controlled trials have found valsartan to be both safe and effective for the treatment of hypertension. With valsartan taken in a dosage of 80 to 320 mg once daily, the mean reduction in diastolic blood pressure is 6 to 9 mm Hg, and the mean reduction in systolic pressure is 3 to 6 mm Hg. Studies have shown that valsartan is as effective as ACE inhibitors enalapril, lisinopril and amlodipine in the treatment of mild to moderate hypertension.

20 The affinity of valsartan for the AT₁ receptor is about 20,000 times greater than its affinity for the AT₂ receptor. In comparison, the affinity of losartan for the AT₁ receptor is about 1,000 times greater than its affinity for AT₂ receptors. The clinical implication of receptor affinity is not yet clear.

25 Valsartan is also available as a combination product with hydrochlorothiazide (Diovan HCT). This combination drug contains 80 or 160 mg of valsartan and 12.5 mg of hydrochlorothiazide. With the addition of hydrochlorothiazide, blood pressure decreases even more (i.e., by 6 mm Hg systolic and 3 mm Hg diastolic). Dosing is once daily.

30 **Irbesartan.** Irbesartan (U.S. Pat. Nos. 5,270,317; 5,994,348; 6,342,247) is a safe and effective angiotensin-II receptor antagonist with an affinity for the AT₁

receptor that is more than 8,500 times greater than its affinity for the AT₂ receptor. This agent has a higher bioavailability (60 to 80 percent) than other drugs in its class.

5 In one study, 530 patients with mild to moderate hypertension were given placebo, losartan in a dosage of 100 mg per day or irbesartan in a dosage of 150 or 300 mg per day. After only one week of therapy, blood pressure trough reduction was significantly greater with irbesartan in a dosage of 300 mg per day than with losartan in a dosage of 100 mg per day.

10 Placebo-controlled trials have shown that irbesartan in a dosage of 150 to 300 mg per day lowers mean systolic blood pressure by 8 to 12 mm Hg and mean diastolic pressure by 5 to 8 mm Hg. Irbesartan has also been found to be as effective as enalapril and atenolol in reducing blood pressure.

A combination product that contains both irbesartan and hydrochlorothiazide is being developed.

15 **Candesartan.** Candesartan cilexetil (U.S. Pat. No. 5,196,444) has been shown to be effective for the treatment of hypertension. Candesartan itself is poorly absorbed after oral administration; the ester prodrug, candesartan cilexetil, improves bioavailability. With oral administration of candesartan cilexetil, conversion to the active compound occurs rapidly and completely during gastrointestinal absorption.
20 The affinity of candesartan for the AT₁ receptor is more than 10,000 times greater than its affinity for the AT₂ receptor.

Candesartan is both safe and well tolerated in dosages of 8 to 32 mg per day. With these dosages, systolic blood pressure is reduced by 8 to 12 mm Hg and diastolic pressure is reduced by 4 to 8 mm Hg.

25 Comparable reductions of diastolic blood pressure have been achieved with candesartan in a dosage of 8 mg per day and enalapril in a dosage of 10 mg per day. In one trial, significant reductions in mean sitting diastolic pressures occurred after 12 weeks of treatment with candesartan in a dosage of 8 or 12 mg per day and enalapril in a dosage of 10 mg per day ($P < 0.01$), but not with candesartan in a
30 dosage of 4 mg per day ($P = 0.074$). The same study compared losartan in a dosage

of 50 mg per day with candesartan in dosages of 8 and 16 mg per day. The 16-mg dosage of candesartan reduced diastolic blood pressure by an adjusted mean of 3.7 mm Hg more than the 50-mg losartan dosage.

5 **Telmisartan.** Telmisartan (U.S. Pat. No. 5,591,762) is the most recently labeled angiotensin-II receptor antagonist. Its affinity for the AT₁ receptor is more than 3,000 times greater than its affinity for the AT₂ receptor. Nonlinear pharmacokinetics yield a greater than proportional increase in plasma telmisartan concentrations with increasing dosages.

10 The efficacy of telmisartan in the treatment of hypertension has been demonstrated in placebo-controlled trials. A three-month study of 440 patients showed that telmisartan in a dosage of 40, 80, 120 or 160 mg per day produced a slightly greater antihypertensive effect than enalapril in a dosage of 20 mg per day. In this study, diastolic blood pressure reductions with telmisartan ranged from 8.6 to 9.3 mm Hg, and systolic blood pressure reductions ranged from 10 to 11.9 mm Hg.
15 The decreases in diastolic and systolic blood pressures for enalapril were 7.2 and 8.2 mm Hg, respectively.

 Like the other angiotensin-II receptor antagonists, telmisartan has been shown to have a side effect profile similar to that of placebo. Clinical trials have demonstrated no rebound hypertension or first-dose orthostatic effect.

20 Most recently, the FDA has approved a new angiotensin II receptor blocker called olmesartan medoxomil (Benicar), for the treatment of hypertension. A 20 mg-starting dose of olmesartan medoxomil has been shown to reduce systolic pressure by an average of 15 mm Hg and diastolic pressure by an average of 12 mm Hg. The manufacturers Sankyo Pharma Inc. stated that studies have shown their drug to be
25 superior to losartan, and the launch of Benicar is expected within the first half of 2002.

 In addition, Iyer et al. (*Proc. Nat. Acad. Sci.* 93: 9960-9965, 1996) explored the possibility of using gene therapy to inhibit AGTR1. They demonstrated that the delivery of angiotensin type 1 receptor antisense by a retrovirally-mediated delivery
30 system resulted in a selective attenuation of the cellular actions of angiotensin II in the neurons of the spontaneously hypertensive (SH) rat model. A single injection

normalized blood pressure in the SH rat on a long-term basis. The use of this approach in patients was proposed. Thus, this type of AGTR1 antagonist is also within the scope of the AIIRA of the instant invention.

Although both ACE inhibitors and AIIRAs prevent the activation of AGTR1, there is a difference between the effects of these two classes of compounds. This is because non-ACE pathways can also produce some angiotensin II. ACE inhibitors also decrease bradykinin breakdown and this action could be involved in some of the beneficial and adverse effects of that class of drugs. Therefore, a potential for differential clinical effects exists for these two classes of drugs. For example, angiotensin receptor blockers are indicated in patients who require an ACE inhibitor but who cannot tolerate it due to drug-induced dry cough. Similar consideration can also be helpful in conjoint administration of morphogen with either ACE inhibitor or AIIRA.

B. Formulations and Methods of Treatment

In one embodiment, the invention comprises a pharmaceutical composition comprising a therapeutically effective amount an ACE inhibitor and an OP/BMP morphogen formulated with pharmaceutically acceptable salt, carrier, excipient or diluent. In one embodiment, the ACE inhibitor is Enalapril. In another embodiment, the ACE ACEI is: any one compound of the formulas I-XXVIII or their salts thereof; acylmercapto and mercaptoalkanoyl prolines; captopril (1-[(2S)-3-mercapto-2-methylpropionyl]-L-proline); ether or thioether mercaptoacyl prolines; zofenopril; carboxyalkyl dipeptides; enalapril (N-(1-ethoxycarbonyl-3-phenylpropyl)-L-ananyl-L-proline); lisinopril; quinapril; ramipril; carboxyalkyl dipeptide mimics; cilazapril; benazapril; phosphinylalkanoyl prolines; fosinopril;trandolopril; phosphonamidate substituted amino or imino acids; phosphonate substituted amino or imino acids and salts thereof; ceronapril ((S)-1-[6-amino-2-[[hydroxyl(4-phenylbutyl)phosphinyl]oxy]-1-oxohexyl]-L-proline); BRL 36,378; MC-838; CGS 14824 (3-([1-ethoxycarbonyl-3-phenyl-(1S)-propyl]-amino)-2,3,4,5-tetrahydro-2-oxo-1-(3S)-benzazepine-1 acetic acid HCL); CGS 16,617 (3(S)-[[[(1S)-5-amino-1-carboxypentyl]amino]2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-ethanoic acid); Cetapril (alacepril, Dainippon); Ru 44570; Cilazapril; Ro 31-2201;

Lisinopril; Indalapril (delapril); Rentiapril (fentiapril, Santen); Indolapril; Spirapril; Perindopril; Quinapril; CI 925 ([3S-[2[R(*)R(*)]]3R(*)]-2-[2-[[1-(ethoxy-carbonyl)-3-phenylpropyl]amino[-1-oxopropyl]]-1,2,3,4-tetrahydro-6,7-dimethoxy-3-isoquinolinecarboxylic acid HCL); WY-44221; mercapto-containing compounds; 5 pivopril; YS980; Omapatrilat; Alacepril; moveltopril; quinaprilat; moexipril; perinodpril (S-9490); pentopril; ancovenin; phenacein; or nicotianamin.

In another embodiment, the invention comprises a pharmaceutical composition comprising a therapeutically effective amount an AIIRA and an OP/BMP morphogen formulated with pharmaceutically acceptable salt, carrier, 10 excipient or diluent. In one embodiment AIIRA is: Losartan (Cozaar[®]), Valsartan (Diovan[®]), Irbesartan (Avapro[®]), Candesartan (Atacand[®]), Telmisartan (Micardis[®]), tasosartan, zolarsartan, Teveten (eprosartan mesylate) or olmesartan medoxomil (Benicar).

In one embodiment, the morphogen in any of the above pharmaceutical 15 composition embodiments is the polypeptide of SEQ ID NO: 3.

In another embodiment, the morphogen in any of the above pharmaceutical composition embodiments is a first polypeptide including at least a C-terminal cysteine domain of a protein selected from: a pro form, a mature form, or a soluble form of a second polypeptide, wherein said second polypeptide is: OP-1, OP-2, OP- 20 3, BMP2, BMP3, BMP4, BMP5, BMP6, or BMP9.

In another embodiment, the morphogen in any of the above pharmaceutical composition embodiments comprises a polypeptide having at least 70% homology or 50% identity with an amino acid sequence of a C-terminal seven-cysteine domain of human OP-1 (SEQ ID NO: 2). In another embodiment, the polypeptide has at 25 least 75% homology or 60% identity with an amino acid sequence of a C-terminal seven-cysteine domain of human OP-1 (SEQ ID NO: 2). In yet another embodiment, the polypeptide has at least 80% homology or 70% identity with an amino acid sequence of a C-terminal seven-cysteine domain of human OP-1 (SEQ ID NO: 2). In yet another embodiment, the polypeptide has at least 90% identity

with an amino acid sequence of a C-terminal seven-cysteine domain of human OP-1 (SEQ ID NO: 2).

5 The invention also comprises a package pharmaceutical comprising any of the pharmaceutical compositions described herein, in association with instructions for administering the composition to a mammal for treatment or prevention of chronic renal failure.

10 The invention also comprises a package pharmaceutical comprising any of the pharmaceutical compositions described herein, in association with instructions for administering the composition to a mammal for delaying the need or reducing the frequency of chronic dialysis treatments.

15 In one embodiment, the invention provides a method of treating or preventing chronic renal failure in a mammal, comprising conjointly administering to said mammal (i) an OP/BMP morphogen, an inducer of endogenous OP/BMP morphogen expression, or an agonist of an OP/BMP morphogen receptor; and (ii) an Angiotensin-Converting Enzyme Inhibitor (ACEI).

20 In one embodiment, the invention provides a method of treating or preventing chronic renal failure in a mammal, comprising conjointly administering to said mammal (i) an OP/BMP morphogen, an inducer of endogenous OP/BMP morphogen expression, or an agonist of an OP/BMP morphogen receptor; and (ii) an Angiotensin- II Receptor Antagonist (AIIRA).

25 In another embodiment, the invention provides a method of treating or preventing chronic renal failure in a mammal, comprising introducing into the kidney of said mammal a therapeutically effective amount of renal mesenchymal progenitor cells pre-treated conjointly with an ACEI and an agent that increases the abundance of an OP/BMP morphogen.

In another embodiment, the invention provides a method of treating or preventing chronic renal failure in a mammal, comprising introducing into the kidney of said mammal a therapeutically effective amount of renal mesenchymal progenitor cells pre-treated conjointly with an AIIRA and an agent that increases the

abundance of an OP/BMP morphogen. In one embodiment the agent is an OP/BMP morphogen. In another embodiment the agent is an inducer of an OP/BMP morphogen. In another embodiment the agent is an agonist of an OP/BMP morphogen receptor.

5 In another embodiment the invention provides for a method for delaying the need for, or reducing the frequency of, chronic dialysis treatments, comprising conjointly administering to a mammal: (i) an OP/BMP morphogen, an inducer of endogenous OP/BMP morphogen expression, or an agonist of an OP/BMP morphogen receptor; and (ii) an ACEI.

10 In another embodiment the invention provides for a method for delaying the need for, or reducing the frequency of, chronic dialysis treatments, comprising conjointly administering to a mammal: (i) an OP/BMP morphogen, an inducer of endogenous OP/BMP morphogen expression or an agonist of an OP/BMP morphogen receptor; and (ii) an AIIRA.

15 In any of the above mentioned methods of treatment or prevention, said mammal may be afflicted with a condition selected from: chronic renal failure (CRF), end-stage renal disease (ESRD), chronic diabetic nephropathy, diabetic glomerulopathy, diabetic renal hypertrophy, hypertensive nephrosclerosis, hypertensive glomerulosclerosis, chronic glomerulonephritis, hereditary nephritis, or
20 renal dysplasia.

 In any of the above mentioned methods of treatment or prevention, the examination of a renal biopsy of said mammal may indicate that said mammal is afflicted with a condition selected from: glomerular hypertrophy, tubular hypertrophy, glomerulosclerosis, or tubulo interstitial sclerosis.

25 In any of the above mentioned methods of treatment or prevention, the examination of a renal biopsy of said mammal may indicate renal fibrosis. In one embodiment the examination may be by ultrasound, NMR or CAT scan of said mammal.

The invention also comprises use of: (i) an OP/BMP morphogen, an inducer of endogenous OP/BMP morphogen expression, or an agonist of an OP/BMP morphogen receptor; and (ii) an Angiotensin-Converting Enzyme Inhibitor (ACEI) for the preparation of a medicament for treating or preventing chronic renal failure
5 in a mammal.

The invention also comprises use of: (i) an OP/BMP morphogen, an inducer of endogenous OP/BMP morphogen expression, or an agonist of an OP/BMP morphogen receptor; and (ii) an Angiotensin- II Receptor Antagonist (AIIRA) for the preparation of a medicament for treating or preventing chronic renal failure in a
10 mammal.

In another embodiment the invention provides for use of mesenchymal progenitor cells that have been pretreated with an ACEI and an agent that increases the abundance of an OP/BMP morphogen for the preparation of a medicament to be introduced into the kidney of a mammal for treating or preventing chronic renal
15 failure in a mammal.

In another embodiment the invention provides for use of mesenchymal progenitor cells that have been pretreated with an AIIRA and an agent that increases the abundance of an OP/BMP morphogen for the preparation of a medicament to be introduced into the kidney of a mammal for treating or preventing chronic renal
20 failure in a mammal. In one embodiment the agent is an OP/BMP morphogen. In another embodiment the agent is an inducer of an OP/BMP morphogen. In another embodiment the agent is an agonist of an OP/BMP morphogen receptor.

The invention also comprises use of an (i) OP/BMP morphogen, an inducer of endogenous OP/BMP morphogen expression, or an agonist of an OP/BMP morphogen receptor; and (ii) an ACEI to prepare a medicament for delaying or
25 reducing the frequency of chronic dialysis treatments in a mammal.

The invention also comprises use of an (i) OP/BMP morphogen, an inducer of endogenous OP/BMP morphogen expression, or an agonist of an OP/BMP

morphogen receptor; and (ii) an AIIRA to prepare a medicament for delaying or reducing the frequency of chronic dialysis treatments in a mammal.

5 In any of the above mentioned embodiments, said mammal may possess a number of functional nephron units which is less than about 40% of a number of functional nephron units present in a mammal having intact healthy kidneys. In one embodiment, said mammal possesses a number of functional nephron units which is less than about 20% of a number of functional nephron units present in a mammal having intact healthy kidneys.

10 In any of the above mentioned embodiments, said mammal may be a kidney transplant recipient. In one embodiment, said mammal possesses only one kidney.

In any of the above mentioned embodiments, examination of a urinary sediment of said mammal may indicate a presence of broad casts.

15 In some of the above mentioned embodiments, said mammal may have a GFR which is chronically less than about 40% of a GFR_{exp} for said mammal. In some of the above mentioned embodiments, said mammal may have a GFR which is chronically less than about 20% of a GFR_{exp} for said mammal.

20 In any of the above mentioned embodiments, said mammal may be a human male weighing at least about 50 kg and has a GFR which is chronically less than about 40 ml/min. In any of the above mentioned embodiments, said mammal may be a human female weighing at least about 40 kg and has a GFR which is chronically less than about 30 ml/min.

In any of the above mentioned embodiments, said method of treatment or prevention, or said medicament, may reduce serum creatinine levels in said mammal by at least about 5% over 3 months.

25 In any of the above mentioned embodiments, prior to said treatment or prevention, said mammal may present a chronic decline in a clinical indicator of renal function, and after at least about 3 months of said treatment or prevention, said indicator may stabilize.

In any of the above mentioned embodiments, at least one of said ACEI, said AIIRA or said morphogen may be administered orally, parenterally, intravenously, intraperitoneally, or into a renal capsule, or by an implanted device. In any of the above mentioned embodiments, a stent may be implanted into said mammal for said administration of at least one of said ACEI, said AIIRA or said morphogen.

In any of the above mentioned embodiments, at least one of said ACEI or said AIIRA, and at least one of said morphogen may be conjointly administered at least once a week for a period of at least about one month.

In any of the above mentioned embodiments, at least one of said ACEI or AIIRA, and at least one of said morphogen may be conjointly administered at least once a week for a period of at least about one year.

In any of the above mentioned embodiments, said ACEI or said AIIRA, and said morphogen may be administered: (i) through different routes or (ii) at different frequencies.

In any of the above mentioned embodiments, said morphogen may be administered at a dosage of about 0.01-1000 $\mu\text{g/kg}$ body weight of said mammal.

In any of the above mentioned embodiments, said morphogen may be administered at a dosage of a dosage of about 10-300 $\mu\text{g/kg}$ body weight of said mammal.

In any of the above mentioned embodiments that comprises the administration or use of ACEI, said ACEI may be administered orally at a concentration of about 1-10,000 mg/L, preferably 10-1000 mg/L, 10-100 mg/L, 100-1000 mg/L, most preferably 100 mg/L.

In any of the above mentioned embodiments that comprises the administration or use of AIIRA, said AIIRA may be administered orally at a concentration of about 0.01-100 mg/kg body weight, preferably 0.1-10 mg/kg body weight, 0.2-5 mg/kg body weight, 0.5-2 mg/kg body weight, most preferably 1 mg/kg body weight.

In any of the above mentioned embodiments, said OP/BMP morphogen and, ACEI or AIIRA may be administered in a single pharmaceutical composition. In any of the above mentioned embodiments, said OP/BMP morphogen and, ACEI or AIIRA may be administered in separate pharmaceutical compositions at or around
5 the same time. In any of the above mentioned embodiments, said OP/BMP morphogen and, ACEI or AIIRA may be administered in separate pharmaceutical compositions at different times.

In any of the above mentioned embodiments, said morphogen may: (a) induce chondrogenesis in an ectopic bone assay; (b) prevent, inhibit, delay or
10 alleviate loss of renal function in an animal model of chronic renal failure, or (c) cause a clinically significant improvement in a standard marker of renal function when administered to a mammal in, or at risk of, chronic renal failure.

In any of the above mentioned embodiments, said morphogen may comprise a polypeptide including at least a C-terminal cysteine domain of a protein selected
15 from: a pro form, a mature form, or a soluble form of a polypeptide, wherein said polypeptide is: OP-1, OP-2, OP-3, BMP2, BMP3, BMP4, BMP5, BMP6, or BMP9.

In any of the above mentioned embodiments, said morphogen may comprise a polypeptide including at least a C-terminal cysteine domain of a polypeptide selected from: a pro form, a mature form, or a soluble form of human OP-1.

20 In one embodiment, the morphogen used in any of the above mentioned embodiments may comprise a polypeptide having at least 70% homology or 50% identity with an amino acid sequence of a C-terminal seven-cysteine domain of human OP-1 (SEQ ID NO: 2). In another embodiment, the morphogen used in any of the above mentioned embodiments may comprise a polypeptide having at least
25 75% homology or 60% identity with an amino acid sequence of a C-terminal seven-cysteine domain of human OP-1 (SEQ ID NO: 2). In another embodiment, the morphogen used in any of the above mentioned embodiments may comprise a polypeptide having at least 80% homology or 70% identity with an amino acid sequence of a C-terminal seven-cysteine domain of human OP-1 (SEQ ID NO: 2).
30 In another embodiment, the morphogen used in any of the above mentioned

embodiments may comprise a polypeptide having at least 90% identity with an amino acid sequence of a C-terminal seven-cysteine domain of human OP-1 (SEQ ID NO: 2).

In any of the above mentioned embodiments said ACEI may be: any one
5 compound of the formulas I-XXVIII or their salts thereof; acylmercapto and mercaptoalkanoyl prolines; captopril (1-[(2S)-3-mercapto-2-methylpropionyl]-L-proline); ether or thioether mercaptoacyl prolines; zofenopril; carboxyalkyl dipeptides; enalapril (N-(1-ethoxycarbonyl-3-phenylpropyl)-L-ananyl-L-proline); lisinopril; quinapril; ramipril; carboxyalkyl dipeptide mimics; cilazapril; benazapril;
10 phosphinylalkanoyl prolines; fosinopril;trandolopril; phosphonamidate substituted amino or imino acids; phosphonate substituted amino or imino acids and salts thereof; ceronapril ((S)-1-[6-amino-2-[[hydroxyl(4-phenylbutyl)phosphinyl]oxy]-1-oxohexyl]-L-proline); BRL 36,378; MC-838; CGS 14824 (3-([1-ethoxycarbonyl-3-phenyl-(1S)-propyl]-amino)-2,3,4,5-tetrahydro-2-oxo-1-(3S)-benzazepine-1 acetic
15 acid HCL); CGS 16,617 (3(S)-[[[(1S)-5-amino-1-carboxypentyl]amino]2,3,4,5-tetrahydro-2-oxo-1H-1-benzazepine-1-ethanoic acid); Cetapril (alacepril, Dainippon); Ru 44570; Cilazapril; Ro 31-2201; Lisinopril; Indalapril (delapril); Rentiapril (fentiapril, Santen); Indolapril; Spirapril; Perindopril; Quinapril; CI 925 ([3S-[2[R(*)R(*)]]3R(*)]-2-[2-[[1-(ethoxy-carbonyl)-3-phenylpropyl]amino]-1-oxopropyl]-1,2,3,4-tetrahydro-6,7-dimethoxy-3-isoquinolinecarboxylic acid HCL);
20 WY-44221; mercapto-containing compounds; pivopril; YS980; Omapatrilat; Alacepril; moveltopril; quinaprilat; moexipril; perinodpril (S-9490); pentopril; ancovenin; phenacein; or nicotianamin. In a preferred embodiment the ACEI is Enalapril.

25 In any of the above mentioned embodiments said AIIRA may be: Losartan (Cozaar[®]), Valsartan (Diovan[®]), Irbesartan (Avapro[®]), Candesartan (Atacand[®]), Telmisartan (Micardis[®]), tasosartan, zolarsartan, Teveten (eprosartan mesylate) or olmesartan medoxomil (Benicar).

30 ACE inhibitors, AIIRAs and/or morphogens may be formulated with one or more pharmaceutically acceptable carriers (additives) and/or diluents. As described in detail below, such pharmaceutical compositions may be specially formulated for

administration in solid or liquid form, including those adapted for the following: (1) oral administration, for example, drenches (aqueous or non-aqueous solutions or suspensions), tablets, boluses, powders, granules, pastes for application to the tongue; (2) parenteral administration, for example, by subcutaneous, intrathecal, 5 intracerebroventricular, intramuscular, or intravenous injection as, for example, a sterile solution or suspension, including administration using a minipump or other mechanical-assisted delivery, such as ALZET osmotic pumps that continuously deliver agents at controlled rates; (3) topical application, for example, as a cream, ointment or spray applied to the skin; or (4) intravaginally or intrarectally, for 10 example, as a pessary, cream or foam. However, in certain embodiments the subject compounds may be simply dissolved or suspended in sterile water. In certain embodiments, the pharmaceutical preparation is non-pyrogenic, i.e., does not elevate the body temperature of a patient.

The phrase 'therapeutically effective amount' as used herein means that 15 amount of a compound, material, or composition comprising a compound of the present invention which is effective for producing some desired therapeutic effect in at least a sub-population of cells in an animal and thereby blocking the biological consequences of that pathway in the treated cells, at a reasonable benefit/risk ratio applicable to any medical treatment.

20 The phrase 'pharmaceutically acceptable' is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

25 The phrase 'pharmaceutically acceptable carrier' as used herein means a pharmaceutically acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting the subject compounds from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be 'acceptable' in the sense 30 of being compatible with the other ingredients of the formulation and not injurious to the patient. Some examples of materials which can serve as pharmaceutically

acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) phosphate buffer solutions; and (21) other non-toxic compatible substances employed in pharmaceutical formulations.

As set out above, certain compounds contain a basic functional group, such as amino or alkylamino, and are, thus, capable of forming pharmaceutically acceptable salts with pharmaceutically acceptable acids. The term 'pharmaceutically acceptable salts' in this respect, refers to the relatively non-toxic, inorganic and organic acid addition salts of compounds of the present invention. These salts can be prepared *in situ* during the final isolation and purification of the compounds of the invention, or by separately reacting a purified compound of the invention in its free base form with a suitable organic or inorganic acid, and isolating the salt thus formed. Representative salts include the hydrobromide, hydrochloride, sulfate, bisulfate, phosphate, nitrate, acetate, valerate, oleate, palmitate, stearate, laurate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthylate, mesylate, glucoheptonate, lactobionate, and laurylsulphonate salts and the like. (See, for example, Berge et al. (1977) "Pharmaceutical Salts", *J. Pharm. Sci.* 66:1-19).

The pharmaceutically acceptable salts of the subject compounds include the conventional nontoxic salts or quaternary ammonium salts of the compounds, e.g., from non-toxic organic or inorganic acids. For example, such conventional nontoxic salts include those derived from inorganic acids such as hydrochloride, hydrobromic, sulfuric, sulfamic, phosphoric, nitric, and the like; and the salts

prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, palmitic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isothionic, and the like.

5 In other cases, compounds contain one or more acidic functional groups and, thus, are capable of forming pharmaceutically acceptable salts with pharmaceutically acceptable bases. The term 'pharmaceutically acceptable salts' in these instances refers to the relatively non-toxic, inorganic and organic base addition salts of compounds of the present invention. These salts can likewise be prepared *in*
10 *situ* during the final isolation and purification of the compounds, or by separately reacting the purified compound in its free acid form with a suitable base, such as the hydroxide, carbonate or bicarbonate of a pharmaceutically acceptable metal cation, with ammonia, or with a pharmaceutically acceptable organic primary, secondary or tertiary amine. Representative alkali or alkaline earth salts include the lithium,
15 sodium, potassium, calcium, magnesium, and aluminum salts and the like. Representative organic amines useful for the formation of base addition salts include ethylamine, diethylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine and the like. (See, for example, Berge et al., *supra*).

Wetting agents, emulsifiers and lubricants, such as sodium lauryl sulfate and
20 magnesium stearate, as well as coloring agents, release agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the compositions.

Examples of pharmaceutically acceptable antioxidants include: (1) water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate,
25 sodium metabisulfite, sodium sulfite and the like; (2) oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol, and the like; and (3) metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like.

30 Formulations of the present invention include those suitable for oral, nasal, topical (including buccal and sublingual), rectal, vaginal and/or parenteral

administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. The amount of active ingredient which can be combined with a carrier material to produce a single dosage form will vary depending upon the host being treated, the particular mode of administration. The amount of active ingredient that can be combined with a carrier material to produce a single dosage form will generally be that amount of the compound which produces a therapeutic effect. Generally, out of one hundred per cent, this amount will range from about 1 per cent to about ninety-nine percent of active ingredient, preferably from about 5 per cent to about 70 per cent, most preferably from about 10 per cent to about 30 per cent.

Methods of preparing these formulations or compositions include the step of bringing into association a compound of the present invention with the carrier and, optionally, one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association a compound of the present invention with liquid carriers, or finely divided solid carriers, or both, and then, if necessary, shaping the product.

Formulations of the invention suitable for oral administration may be in the form of capsules, cachets, pills, tablets, lozenges (using a flavored basis, usually sucrose and acacia or tragacanth), powders, granules, or as a solution or a suspension in an aqueous or non-aqueous liquid, or as an oil-in-water or water-in-oil liquid emulsion, or as an elixir or syrup, or as pastilles (using an inert base, such as gelatin and glycerin, or sucrose and acacia) and/or as mouth washes and the like, each containing a predetermined amount of a compound of the present invention as an active ingredient. A compound of the present invention may also be administered as a bolus, electuary or paste.

In solid dosage forms of the invention for oral administration (capsules, tablets, pills, dragees, powders, granules and the like), the active ingredient is mixed with one or more pharmaceutically acceptable carriers, such as sodium citrate or dicalcium phosphate, and/or any of the following: (1) fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; (2) binders, such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone,

5 sucrose and/or acacia; (3) humectants, such as glycerol; (4) disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; (5) solution retarding agents, such as paraffin; (6) absorption accelerators, such as quaternary ammonium compounds; (7) wetting
agents, such as, for example, cetyl alcohol and glycerol monostearate; (8)
absorbents, such as kaolin and bentonite clay; (9) lubricants, such a talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof; and (10) coloring agents. In the case of capsules, tablets and pills, the pharmaceutical compositions may also comprise buffering agents. Solid
10 compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugars, as well as high molecular weight polyethylene glycols and the like.

A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared using binder (for
15 example, gelatin or hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (for example, sodium starch glycolate or cross-linked sodium carboxymethyl cellulose), surface-active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

20 The tablets, and other solid dosage forms of the pharmaceutical compositions of the present invention, such as dragees, capsules, pills and granules, may optionally be scored or prepared with coatings and shells, such as enteric coatings and other coatings well known in the pharmaceutical-formulating art. They may also be formulated so as to provide slow or controlled release of the active ingredient
25 therein using, for example, hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile, other polymer matrices, liposomes and/or microspheres. They may be sterilized by, for example, filtration through a bacteria-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions that can be dissolved in sterile water, or some other sterile injectable
30 medium immediately before use. These compositions may also optionally contain opacifying agents and may be of a composition that they release the active

ingredient(s) only, or preferentially, in a certain portion of the gastrointestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes. The active ingredient can also be in microencapsulated form, if appropriate, with one or more of the above-described
5 excipients.

Liquid dosage forms for oral administration of the compounds of the invention include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active ingredient, the liquid dosage forms may contain inert diluents commonly used in the art, such as,
10 for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and
15 mixtures thereof.

Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, coloring, perfuming and preservative agents.

Suspensions, in addition to the active compounds, may contain suspending
20 agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

Formulations of the pharmaceutical compositions of the invention for rectal or vaginal administration may be presented as a suppository, which may be prepared
25 by mixing one or more compounds of the invention with one or more suitable nonirritating excipients or carriers comprising, for example, cocoa butter, polyethylene glycol, a suppository wax or a salicylate, and which is solid at room temperature, but liquid at body temperature and, therefore, will melt in the rectum or vaginal cavity and release the active compound.

Formulations of the present invention which are suitable for vaginal administration also include pessaries, tampons, creams, gels, pastes, foams or spray formulations containing such carriers as are known in the art to be appropriate.

5 Dosage forms for the topical or transdermal administration of a compound of this invention include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants. The active compound may be mixed under sterile conditions with a pharmaceutically acceptable carrier, and with any preservatives, buffers, or propellants that may be required.

10 The ointments, pastes, creams and gels may contain, in addition to an active compound of this invention, excipients, such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof.

15 Powders and sprays can contain, in addition to a compound of this invention, excipients such as lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder, or mixtures of these substances. Sprays can additionally contain customary propellants, such as chlorofluorohydrocarbons and volatile unsubstituted hydrocarbons, such as butane and propane.

20 Transdermal patches have the added advantage of providing controlled delivery of a compound of the present invention to the body. Such dosage forms can be made by dissolving or dispersing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate of such flux can be controlled by either providing a rate controlling membrane or dispersing the compound in a polymer matrix or gel.

25 Pharmaceutical compositions of this invention suitable for parenteral administration comprise one or more compounds of the invention in combination with one or more pharmaceutically acceptable sterile isotonic aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, or sterile powders which may be reconstituted into sterile injectable solutions or dispersions just prior to use, which may contain antioxidants, buffers, bacteriostats, solutes which render 30 the formulation isotonic with the blood of the intended recipient or suspending or thickening agents.

Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable
5 organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of the action of
10 microorganisms may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents that
15 delay absorption such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of a drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material having poor water solubility. The rate of absorption of the drug then
20 depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are made by forming microencapsulated matrices of
25 the subject compounds in biodegradable polymers such as polylactide-polyglycolide. Depending on the ratio of drug to polymer, and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or
30 microemulsions that are compatible with body tissue.

When the compounds of the present invention are administered as pharmaceuticals, to humans and animals, they can be given per se or as a pharmaceutical composition containing, for example, 0.1 to 99.5% (more preferably, 0.5 to 90%) of active ingredient in combination with a pharmaceutically acceptable carrier.

Morphogens, morphogen inducers, or agonists of morphogen receptors, as well as ACE inhibitors (ACEIs) may be administered by any route which is compatible with the particular morphogen, inducer, agonist, or ACEI employed. Thus, as appropriate, administration may be oral or parenteral, including intravenous and intraperitoneal routes of administration. In addition, administration may be by periodic injections of a bolus of the morphogen, inducer, agonist or ACEI, or may be made more continuous by intravenous or intraperitoneal administration from a reservoir which is external (*e.g.*, an ix. bag) or internal (*e.g.*, a bioerodable implant, or a colony of implanted, morphogen-producing cells).

Therapeutic agents of the invention (*i.e.*, morphogens, morphogen inducers, agonists of morphogen receptors, or ACEI) may be provided to an individual by any suitable means, directly (*e.g.*, locally, as by injection, implantation or topical administration to a tissue locus) or systemically (*e.g.*, parenterally or orally). Where the agent is to be provided parenterally, such as by intravenous, subcutaneous, intramolecular, ophthalmic, intraperitoneal, intramuscular, buccal, rectal, vaginal, intraorbital, intracerebral, intracranial, intraspinal, intraventricular, intrathecal, intracisternal, intracapsular, intranasal or by aerosol administration, the agent preferably comprises part of an aqueous or physiologically compatible fluid suspension or solution. Thus, the carrier or vehicle for the agent(s) is physiologically acceptable so that in addition to delivery of the desired agent(s) to the patient, it does not otherwise adversely affect the patient's electrolyte and/or volume balance. The fluid medium for the agent thus can comprise normal physiologic saline (*e.g.*, 9.85% aqueous NaCl, 0.15 M, pH 7-7.4).

Association of the mature morphogen dimer with a morphogen pro domain results in the pro form of the morphogen which typically is more soluble in physiological solutions than the corresponding mature form. In fact, endogenous

morphogens are thought to be transported (*e.g.*, secreted and circulated) in the mammalian body in this form. This soluble form of the protein can be obtained from culture medium of morphogen-secreting mammalian cells, *e.g.*, cells transfected with nucleic acid encoding and competent to express the morphogen. Alternatively,

5 a soluble species can be formulated by complexing the mature, morphogenically active polypeptide dimer (or an active fragment thereof) with a morphogen pro domain polypeptide or a solubility-enhancing fragment thereof. Solubility-enhancing pro domain fragments can be any N-terminal, C-terminal or internal fragment of the pro region of a member of the morphogen family that complexes

10 with the mature polypeptide dimer to enhance stability and/or dissolubility of the resulting noncovalent or covalent complex. Typically, useful fragments are those cleaved at the proteolytic site Arg-Xaa-Xaa-Arg (SEQ ID NO: 30). A detailed description of soluble complex forms of morphogenic proteins, including how to make, test and use them, is described in WO 94/03600 (PCT US 93/07189). In the

15 case of OP-1, useful pro domain polypeptide fragments include the intact pro domain polypeptide (residues 30-292) and fragments 48-292 and 158-292, all of SEQ ID No. 3. Another molecule capable of enhancing solubility and particularly useful for oral administrations, is casein. For example, addition of 0.2% casein increases solubility of the mature active form of OP-1 by 80%. Other components

20 found in milk and/or various serum proteins may also be useful.

Useful solutions for parenteral administration may be prepared by any of the methods well known in the pharmaceutical art, described, for example, in REMMINGTON'S PHARMACEUTICAL SCIENCES (Gennaro, A., ed.), Mack Pub., 1990. Formulations of the therapeutic agents of the invention may include, for

25 example, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, hydrogenated naphthalenes, and the like. Formulations for direct administration, in particular, may include glycerol and other compositions of high viscosity to help maintain the agent at the desired locus. Biocompatible, preferably bioresorbable, polymers, including, for example, hyaluronic acid, collagen, tricalcium phosphate,

30 polybutyrate, lactide, and glycolide polymers and lactide/glycolide copolymers, may be useful excipients to control the release of the agent in vivo. Other potentially useful parenteral delivery systems for these agents include ethylene-vinyl acetate

copolymer particles, osmotic pumps, implantable infusion systems, and liposomes. Formulations for inhalation administration contain as excipients, for example, lactose, or may be aqueous solutions containing, for example, polyoxyethylene 9-lauryl ether, glycocholate and deoxycholate, or oily solutions for administration in the form of nasal drops, or as a gel to be applied intranasally. Formulations for parenteral administration may also include glycocholate for buccal administration, methoxysalicylate for rectal administration, or citric acid for vaginal administration. Suppositories for rectal administration may also be prepared by mixing the morphogen, inducer or agonist with a non-irritating excipient such as cocoa butter or other compositions which are solid at room temperature and liquid at body temperatures.

Formulations for topical administration to the skin surface may be prepared by dispersing the morphogen, inducer, agonist, or ACEI with a dermatologically acceptable carrier such as a lotion, cream, ointment or soap. Particularly useful are carriers capable of forming a film or layer over the skin to localize application and inhibit removal. For topical administration to internal tissue surfaces, the agent may be dispersed in a liquid tissue adhesive or other substance known to enhance adsorption to a tissue surface. For example, hydroxypropylcellulose or fibrinogen/thrombin solutions may be used to advantage. Alternatively, tissue-coating solutions, such as pectin-containing formulations may be used.

Alternatively, the agents described herein may be administered orally. Oral administration of proteins as therapeutics generally is not practiced, as most proteins are readily degraded by digestive enzymes and acids in the mammalian digestive system before they can be absorbed into the bloodstream. However, the morphogens described herein typically are acid stable and protease-resistant (see, for example, U.S. Pat. No. 4,968,590). In addition, at least one morphogen, OP-1, has been identified in mammary gland extract, colostrum and 57-day milk. Moreover, the OP-1 purified from mammary gland extract is morphogenically active and is also detected in the bloodstream. Maternal administration, via ingested milk, may be a natural delivery route of TGF- β superfamily proteins. Letterio, et al., Science 264: 1936-1938 (1994), report that TGF- β is present in murine milk, and that radio-

labeled TGF- β is absorbed by gastrointestinal mucosa of suckling juveniles. Labeled, ingested TGF- β appears rapidly in intact form in the juveniles' body tissues, including lung, heart and liver. Finally, soluble form morphogen, e.g., mature morphogen associated with the pro domain, is morphogenically active. These findings, as well as those disclosed in the examples below, indicate that oral and parenteral administration are viable means for administering TGF- β superfamily proteins, including the morphogens, to an individual. In addition, while the mature forms of certain morphogens described herein typically are sparingly soluble, the morphogen form found in milk (and mammary gland extract and colostrum) is readily soluble, probably by association of the mature, morphogenically active form with part or all of the pro domain of the expressed, full length polypeptide sequence and/or by association with one or more milk components. Accordingly, the compounds provided herein may also be associated with molecules capable of enhancing their solubility in vitro or in vivo.

Most ACE inhibitors can be orally administered. In a preferred embodiment, ACE inhibitor as a pharmaceutical composition can be orally administered through drinking water or other suitable liquid carrier.

The compounds provided herein may also be associated with molecules capable of targeting the morphogen, inducer, agonist, or ACEI to the desired tissue. For example, an antibody, antibody fragment, or other binding protein that interacts specifically with a surface molecule on cells of the desired tissue, may be used. Useful targeting molecules may be designed, for example, using the single chain binding site technology disclosed in U.S. Pat. No. 5,091,513. Targeting molecules can be covalently or non-covalently associated with the morphogen, inducer, agonist, or ACEI.

As will be appreciated by one of ordinary skill in the art, the formulated compositions contain therapeutically effective amounts of the morphogen, morphogen inducers, agonists of morphogen receptors, or ACEI. That is, they contain an amount which provides appropriate concentrations of the agent to the affected tissue for a time sufficient to stimulate a detectable restoration of impaired renal system function, up to and including a complete restoration thereof. As will be

appreciated by those skilled in the art, these concentrations will vary depending upon a number of factors, including the biological efficacy of the selected agent, the chemical characteristics (e.g., hydrophobicity) of the specific agent, the formulation thereof, including a mixture with one or more excipients, the administration route, and the treatment envisioned, including whether the active ingredient will be administered directly into a tissue site, or whether it will be administered systemically. The preferred dosage to be administered is also likely to depend on variables such as the condition of the diseased or damaged tissues, and the overall health status of the particular mammal. As a general matter, single, daily, biweekly or weekly dosages of 0.00001-1000 mg of a morphogen are sufficient, with 0.0001-100 mg being preferable, and 0.001 to 10 mg being even more preferable. Alternatively, a single, daily, biweekly or weekly dosage of about 0.01-1000 µg/kg body weight, more preferably about 0.01-10 µg/kg body weight, or about 10-300 µg/kg body weight may be advantageously employed. As a general matter, single, daily, biweekly, or weekly dosages of ACEI can be administered orally at an amount of about 0.01 – 300 mg/kg body weight, preferably 0.1-30 mg/kg BW, 0.1-3 mg/kg BW, 1-30 mg/kg BW, most preferably about 1-3 mg/kg BW, in, for example, drinking water, are appropriate for ACE inhibitors. The concentrations can be accordingly adjusted or alternatively expressed as the amount of drug that needs to be administered per day per kg of body weight, if other factors (such as the average body weight of a subject mammalian patient being treated, and the average amount of water consumed per day by said specific mammalian patient) are provided. The present effective dose can be administered in a single dose or in a plurality (two or more) of installment doses, as desired or considered appropriate under the specific circumstances. A bolus injection or diffusible infusion formulation can be used. If desired to facilitate repeated or frequent infusions, implantation of a semi-permanent stent (e.g., intravenous, intraperitoneal, intracisternal or intracapsular) may be advisable. It should also be understood that the dosages of morphogens and/or ACEIs, when conjointly administered, may be different from the dosages of morphogens or ACEIs when they are administered alone (not conjoint administration). It should also be understood that a particular dosage of morphogen or ACEI for treating / preventing chronic renal failure may be different from dosages

used for other non-related (such as bone morphogenesis, etc.) uses of the morphogens and/or ACEIs.

5 The morphogens, inducers, agonists, or ACEI of the invention may, of course, be administered alone or in combination with other molecules known to be beneficial in the treatment of the conditions described herein. For example, various well-known growth factors, hormones, enzymes, therapeutic compositions, antibiotics, or other bioactive agents can also be administered with the morphogen and ACEI. Thus, various known growth factors such as NGF, EGF, PDGF, IGF, FGF, TGF- α , and TGF- β , as well as enzymes, enzyme inhibitors, antioxidants, anti-
10 inflammatory agents, free radical scavenging agents, antibiotics and/or chemoattractant / chemotactic factors, can be included in the present morphogen and ACEI formulation.

Finally, it should be understood that morphogen formulation and ACEI formulations of the present invention can be either formulated together, in a single
15 pharmaceutical composition, or formulated separately, in two or more pharmaceutical compositions. It should also be understood that the same formulation can be administered through different routes, depending on specific needs or appropriate treatment conditions.

C. Types of Chronic Renal Failures

20 Many types of Chronic Renal Failure diseases can be treated according to the instant invention. The following discussions are for illustration purpose only, and should not be construed to be limiting in any respect.

The present invention is directed to methods of prevention and/or treatment, and pharmaceutical preparations for use in the prevention and/or treatment, of
25 vertebrate subjects (preferably mammalian subjects) in, or at risk of, chronic renal failure, or at risk of the need for renal replacement therapy. Such subjects include subjects already afflicted with chronic renal failure, or which have already received renal replacement therapy, as well as any subject reasonably expected to suffer a progressive loss of renal function associated with progressive loss of functioning
30 nephron units. Whether a particular subject is at risk is a determination which may

5 routinely be made by one of ordinary skill in the relevant medical or veterinary art. Subjects in, or at risk of, chronic renal failure, or at risk of the need for renal replacement therapy, include but are not limited to the following: subjects which may be regarded as afflicted with chronic renal failure (CRF), end-stage renal
10 disease (ESRD), chronic diabetic nephropathy, diabetic glomerulopathy, diabetic renal hypertrophy, hypertensive nephrosclerosis, hypertensive glomerulosclerosis, chronic glomerulonephritis, hereditary nephritis, or renal dysplasia; subjects having a biopsy indicating glomerular hypertrophy, tubular hypertrophy, chronic glomerulosclerosis, and/or chronic tubulointerstitial sclerosis; subjects having an-
15 ultrasound, MRI, CAT scan, or other non-invasive examination indicating renal fibrosis; subject shaving an unusual number of broad casts present in urinary sediment; subjects having a GFR which is chronically less than about 50%, and more particularly less than about 40%, 30% or 20%, of the expected GFR for the subject; human male subjects weighing at least about 50 kg and having a GFR which
20 is chronically less than about 50 ml/min, and more particularly less than about 40 ml/min, 30 ml/min or 20 ml/min; human female subjects weighing at least about 40 kg and having a GFR which is chronically less than about 40 mL/min, and more particularly less than about 30 mL/min, 20 ml/min or 10 ml/min; subjects possessing a number of functional nephron units which is less than about 50%, and more
25 particularly less than about 40%, 30% or 20%, of the number of functional nephron units possessed by a healthy but otherwise similar subject; subjects which have a single kidney; and subjects which are kidney transplant recipients.

Chronic renal failure (CRF) can be classified by the site (location) of primary damage: Pre-renal CRF, Post-renal CRF and Renal CRF.

25 Pre-Renal CRF. Some medical conditions cause continuous hypoperfusion (low blood flow) of the kidneys, leading to kidney atrophy (shrinking), loss of nephron function, and chronic renal failure (CRF). These conditions include poor cardiac function, chronic liver failure, and atherosclerosis ("hardening") of the renal arteries. Each of these conditions can induce ischemic nephropathy, which is a result
30 of inadequate blood flow (hypoperfusion) to the kidneys. Hypoperfusion manifests

as a progressive loss of kidney function and kidney atrophy (shrinkage). Renal failure results when this process damages both kidneys.

5 Post-Renal CRF. Interference with the normal flow of urine can produce backpressure within the kidneys, can damage nephrons, and lead to obstructive uropathy, a disease of the urinary tract. Abnormalities that may hamper urine flow and cause post-renal CRF include the following:

- **Bladder outlet obstruction** due to an enlarged prostate gland or bladder stone;
- 10 • **Neurogenic bladder**, an over distended bladder caused by impaired communicator nerve fibers from the bladder to the spinal cord;
- **Kidney stones** in both ureters, the tubes that pass urine from each kidney to the bladder;
- **Obstruction of the tubules**, the end channels of the renal nephrons;
- **Retroperitoneal fibrosis**, the formation of fiber-like tissue behind the
15 peritoneum, the membrane that lines the abdominal cavity;
- **Vesicoureteral reflux (VUR)**, the backward flow of urine from the bladder into a ureter.

Renal CRF. Chronic renal failure caused by changes within the kidneys, is called renal CRF, and is broadly categorized as follows:

- 20 • **Diabetic nephropathy**, kidney disease associated with diabetes; the most common cause of CRF;
- **Hypertension Nephrosclerosis**, a condition that occurs with increased frequency in African Americans; the second leading cause of CRF;
- **Chronic glomerular nephritis**, a condition caused by diseases that affect
25 the glomeruli and bring about progressive dysfunction;
- **Chronic interstitial nephritis**, a condition caused by disorders that ultimately lead to progressive scarring of the interstitium;

- **Renal vascular CRF**, large vessel abnormalities such as renal artery stenosis (narrowing of the large arteries that supply the kidneys);
- **Vasculitis**, inflammation of the small blood vessels;
- **Cystic kidney disease**, kidney disease distinguished by multiple cysts (lined cavities or sacs);
- **Hereditary diseases of the kidney**, such as Alport's syndrome (hereditary nephritis).

More detailed descriptions of a few of the above conditions are provided below.

Diabetic nephropathy is kidney disease that develops as a result of diabetes mellitus (DM). DM, also called simply diabetes, affects approximately 5% of the U.S. population. This disease damages many organs, including the eyes, nerves, blood vessels, heart, and kidneys. DM is the most common cause of kidney failure in the United States and accounts for over one-third of all patients who are on dialysis.

DM patients are unable to metabolize carbohydrates (e.g., food starches, sugars, cellulose) properly. The disease is characterized by excessive amounts of sugar in the blood (hyperglycemia) and urine; inadequate production and/or utilization of insulin; and by thirst, hunger, and loss of weight.

Diabetics who require daily insulin shots to maintain life have *insulin-dependent diabetes mellitus*, or DM 1. In this type of diabetes, the pancreas β cells secrete little or no insulin and the blood sugar level remains high, unless treated. DM 1 usually occurs in children and young adults and is often called juvenile onset diabetes. Onset of the disease is abrupt. The patient becomes very sick and requires immediate insulin therapy. Approximately 1 million people in the United States have DM 1.

Approximately 25% to 40% of patients with DM 1 ultimately develop diabetic nephropathy (DN), which progresses through about five predictable stages. During Stage 1 (very early diabetes), increased demand upon the kidneys is indicated by an above-normal glomerular filtration rate (GFR). During Stage 2

(developing diabetes), the GFR remains elevated or has returned to normal, but glomerular damage has progressed to significant microalbuminuria (small but above-normal level of the protein albumin in the urine). Patients in stage 2 excrete more than 30 mg of albumin in the urine over a 24-hour period. Significant
5 microalbuminuria will progress to end-stage renal disease (ESRD). Therefore, all diabetes patients should be screened for microalbuminuria on a routine (yearly) basis. During Stage 3 (overt, or dipstick-positive diabetes), glomerular damage has progressed to clinical albuminuria. The urine is "dipstick positive," containing more than 300 mg of albumin in a 24-hour period. Hypertension (high blood pressure)
10 typically develops during stage 3. During Stage 4 (late-stage diabetes), glomerular damage continues, with increasing amounts of protein albumin in the urine. The kidneys' filtering ability has begun to decline steadily, and blood urea nitrogen (BUN) and creatinine (Cr) has begun to increase. The glomerular filtration rate (GFR) decreases about 10% annually. Almost all patients have hypertension at stage
15 4. During Stage 5 (end-stage renal disease, ESRD), GFR has fallen to approximately 10 milliliters per minute (<10 mL/min) and renal replacement therapy (i.e., hemodialysis, peritoneal dialysis, kidney transplantation) is needed. Progression through these five stages is rather predictable because the onset of DM 1 can be identified, and most patients are free from age-related medical problems.

20 *Non-insulin-dependent diabetes*, or DM 2, differs from DM 1 in that the main problem is a peripheral resistance to the action of the insulin. DM 2 usually occurs in adults over the age of 40 who are overweight and have a family history of the disease. Some patients can manage their diabetes with weight loss and changes in their diet. Others require medication, and many with DM 2 eventually require
25 insulin. Onset is gradual, and patients may be sick for quite some time without knowing it. Nearly 95% of diabetics are diagnosed with DM 2. An estimated 5% to 15% of DM 2 cases also progress through the five stages of diabetic nephropathy (DN), but the timeline is not as clear. Some patients advance through the stages very quickly.

30 Renal artery stenosis (RAS) is the narrowing of the lining of the main artery that supplies the kidney. Most RAS is caused by atherosclerosis or "hardening of the

arteries." Atherosclerosis is the build up of cholesterol deposits, or plaque, in the lining of the arteries. Depending on the degree of narrowing, patients can develop hypertension called renal vascular hypertension (RVH). This form of hypertension is the most common cause of secondary hypertension. In fact, hypertension is second
5 only to diabetes as the leading cause of kidney failure. In the US, 15%-20% of kidney failure cases are due to hypertension. RVH occurs when RAS produces a critical narrowing of the artery that supplies one of the kidneys. Critical RAS is defined as at least 70% narrowing of the renal artery, based on angiographic (blood vessel x-ray) evaluation. Reduced blood flow through the renal artery causes the
10 kidney to release increased amounts of the hormone renin. Renin, a powerful blood pressure regulator, initiates a series of chemical events that result in hypertension. Renal vascular hypertension can be very severe and difficult to control.

The kidney with RAS suffers from the decreased blood flow and often shrinks in size (atrophy). This process is called ischemic nephropathy. The other
15 kidney is at risk for developing damage from the hypertension, often developing hypertensive nephrosclerosis. The persistent elevated blood pressures in this non-stenotic kidney can cause progressive scarring (sclerosis) leading to progressive loss of filtering function in this kidney as well. Both unilateral RAS and bilateral RAS can ultimately lead to chronic renal failure.

20 There are two types of RAS: Atherosclerotic Renal Artery Stenosis (AS-RAS) and Fibromuscular Dysplasia (FMD). AS-RAS is due to the build-up of cholesterol on the inner lining of the renal artery. It is exceedingly more common than the unusual case of FMD-RAS. FMD-RAS occurs almost exclusively in women aged 30 to 40 and rarely affects African Americans or Asians. FMD-RAS is
25 due to an abnormality in the muscular lining of the renal artery. There is often a familial history of FMD RAS.

Cystic kidney disease describes several conditions in which fluid-filled cysts form in the kidneys. Cysts generally develop in weak segments of the tubules that carry urine from the glomeruli. The cyst's growth displaces healthy kidney tissue.
30 The kidneys expand to accommodate the cyst, which can weigh as much as 20 pounds. Three factors determine cyst classification: its cause (acquired, inherited),

its features (complicated, simple, multiple, single), and its location (outer [cortical] or inner [medullary] kidney tissue).

Polycystic kidney disease (PKD; common, with several cysts in the kidney) is a primary cystic kidney disease. PKD type 1 and PKD type 2 are caused by
5 autosomal dominant mutations on chromosomes 16 and 4, respectively, and run in families. PKD autosomal recessive has been linked to chromosome 6. Polycystic kidney disease (PKD) is the most frequently inherited disease; it affects approximately 600,000 people in the United States and over 12,000,000 worldwide. Most suffer from the autosomal dominant type. It is the fourth leading cause of
10 kidney failure and causes 10% of all end-stage renal disease (ESRD), usually between the ages of 40 and 60. It affects men, women, and races equally.

Secondary cystic kidney disease include Acquired cystic kidney disease (ACKD); Medullary cystic disease (inner kidney), which includes Juvenile nephronophthisis (during adolescence) and Medullary sponge kidney (deterioration
15 of kidney with cysts); and Renal cell cancer associated cysts.

Autosomal dominant medullary cystic kidney disease (MCK) causes cysts to form in the inner tissue of the kidney and can develop at a very early age. Recessive juvenile nephronophthisis usually occurs later than MCK, but is associated with similar symptoms, including chronic renal failure and growth problems. Small cysts
20 in the collecting ducts of the inner kidney characterize medullary sponge kidney (MSK), which is associated with hematuria and kidney stones, but not chronic renal failure (CFR).

Acquired cystic kidney disease (ACKD) affects patients with chronic renal failure and causes hematuria, erythrocytosis (increase in red blood cells), and is
25 associated with the development of cancer. Causes of acquired cystic kidney disease (ACKD) are long-term disease (glomerulonephritis) and the scarring that often results from dialysis. ACKD is common among patients with chronic renal failure (CRF). Nearly all of those who use dialysis for more than 5 years develop ACKD.

Proteinuria is an abnormally high amount of protein in the urine. Proteins in
30 the blood, like albumin and immunoglobulin, help coagulation (clotting), balance bodily fluids, and fight infection. The kidneys remove wastes from protein-rich

blood through millions of tiny filtering screens called glomeruli. Most proteins are too large to pass through the glomeruli into the urine. The glomeruli are negatively charged, so they repel the negatively charged proteins. Thus, a size and charge barrier keeps protein molecules from entering the urine. But when the glomeruli are
5 damaged, proteins of various sizes pass through them and are excreted in the urine.

The following five types of proteinuria are distinguished by milligrams (mg) of protein measured during a 24-hour urine collection:

1. Microalbuminuria	30 - 150 mg
2. Mild	150 – 500 mg
3. Moderate	500 – 1000 mg
4. Heavy	1000 – 3000 mg
5. Nephrotic range	more than 3500 mg

As kidney disease progresses, more protein enters the urine. People with
10 nephrotic-range proteinuria typically have extensive glomeruli damage and usually develop nephrotic syndrome (see below).

Hypertension and diabetes are the two biggest risk factors for proteinuria. Old age and weight gain also increase the risk. The following conditions cause proteinuria:

- 15
 - Acute glomerulonephritis;
 - Amyloidosis (protein deposits associated with chronic disease);
 - Focal glomerulonephritis;
 - Hypertension;
 - IgA nephropathy;
- 20
 - Mesangial proliferation
 - Minimal change disease;

Foamy urine and swelling (edema) are two signs of proteinuria that become more evident as the disease progresses. Excess protein can cause the urine to foam in water. This occurs because protein changes the surface tension between urine and water. Edema usually only occurs in nephrotic range proteinuria.

5 Albumin is particularly useful in absorbing bodily fluid into the blood. Because the albumin molecule is relatively small, it is often among the first proteins to enter the urine after glomeruli are damaged. Therefore, even minor kidney dysfunction is detectable with proper diagnosis of microralbuminuria. Reduced albumin level in the blood causes fluid retention and swelling that is first noticeable
10 in the hands, lower legs, and feet. In more serious cases, the abdomen and face may swell.

 Orthostatic proteinuria is a disorder seen occasionally in children and young adults who leak significant amounts of urine when they are upright (orthostatic). Presumably, standing increases the pressure on the glomeruli and causes more
15 protein to enter the urine, while lying down relieves pressure and causes less protein leakage. This is a benign disorder that most young people outgrow.

 Hypertensive people who develop proteinuria stand a significant chance for kidney failure. African Americans are 20 times more likely than Caucasians to develop hypertensive-related kidney failure. Proteinuria in people with diabetes may
20 be a sign that kidney disease is worsening. Microalbuminuria is often cited as a risk for coronary artery disease (CAD) and is often diagnostic of it and related cardiovascular conditions.

Nephrotic syndrome (NS) is a condition that is often caused by any of a group of diseases that damage the kidneys' filtering system, the glomeruli. The
25 structure of the glomeruli prevents most protein from getting filtered through into the urine. Normally, a person loses less than 150 mg of protein in the urine in a 24-hour period. Nephrotic-range proteinuria, the urination of more than 3.5 grams of protein during a 24-hour period, or 25 times the normal amount, is the primary indicator of NS.

30 About two in every 10,000 people experience nephrotic syndrome. Nephrotic syndrome prevalence is difficult to establish in adults because the condition is

usually a result of an underlying disease. In children, it is diagnosed in more boys than girls, usually between 2 and 3 years of age.

In addition to proteinuria, there are three main symptoms of nephrotic syndrome associated with protein leaking into the urine:

- 5 • Hypoalbuminemia (low level of albumin in the blood);
- Edema (swelling);
- Hypercholesterolemia (high level of cholesterol in the blood);

10 *Hypoalbuminemia* is a low level of albumin (a protein) in the blood due to proteinuria. Low albumin in the blood causes fluid to move from the blood into the tissue, causing swelling. The kidney perceives the decrease of fluid in the blood and aggressively retains as much fluid and salt as it can. This contributes to the body's fluid-overload state.

15 Nephrotic-related swelling makes tissue puffy, soft, and impressionable to the touch. Edema is most common in the legs and feet, especially after standing all day. It can cause feelings of tightness in the extremities and may affect mobility. In later stages, swelling may occur in the abdomen (ascites), hands, and around the eyes in the morning (called periorbital edema). In later stages, the whole body may swell (anasarca). Some people gain weight after fluid builds up in their bodies for a long time.

20 *Hypercholesterolemia*, high blood cholesterol, is common in nephrotic syndrome. In addition to albumin, other important enzymes involved in cholesterol metabolism slip through the glomeruli, which contribute to high blood cholesterol.

25 Nephrotic syndrome is associated with renal failure. The disease that causes NS can damage the glomeruli and can interfere with their ability to clean the blood. The edema that is present in the legs may also be occurring in the kidney tissue itself and can interfere with the kidneys' ability to clean the blood. Renal failure can either be gradual (CRF) or acute (ARF).

A hypercoaguable state, in which the blood abnormally overclots, is also seen in some patients with NS. This means that they are at risk for developing a

blood clot in the legs or in the renal veins that transport blood from the kidney.
Some patients take blood thinners to prevent this complication.

There are a number of different disorders that can cause NS. Diabetes and, to a lesser extent, hypertension can cause diffuse damage to the glomeruli and can ultimately lead to NS. The following diseases can cause specific damage to the glomeruli and often result in the development of heavy proteinuria and in many instances NS: Amyloidosis (the stiffening and subsequent malfunction of the kidney due to fibrous protein deposit in the tissue); Congenital nephrosis; Focal segmental glomerular sclerosis (FSGS) (creates scar tissue in the glomerulus, damaging its protein-repellant membrane); Glomerulonephritis (GN), including Diffuse mesangial proliferative GN (affecting the messangium), Membranous (damages the protein-repellant membrane), and Post infectious (occurs after an infection); IgA nephropathy (Berger's disease) (deposit of specific immunoglobulin A causing an inflammatory reaction and leading to glomerulonephritis); Minimal change disease (Nil's disease); and Pre-eclampsia (rarely associated with NS, more often associated with heavy proteinuria).

Many of these diseases tend to occur more often in certain age groups. Less than 1yr old: Congenital nephrosis. Less than 15 years old: Min change, FSGS, and Other. Age 15 to 40: Min change, FSGS, and Other. Over age 40: Membranous GN, and Diabetic nephropathy. Over 60: Amyloidosis may account for up to 20% of cases.

In addition, physical injury to the kidneys (loss of one kidney plus damage to the other, etc.) and other equivalent situations, such as complete or partial loss of kidney function due to diseases affecting the total number of functional nephrons (filtering units that consist of a glomerulus and corresponding tubule), may cause the remaining functional nephrons to "attempt" compensating for the renal damage by hyperfiltration (excessive straining of the blood). Over time, hyperfiltration causes further loss of renal function, leading to chronic renal failure.

IV. Exemplification

Practice of the invention, including additional preferred aspects and embodiments thereof, will be still more fully understood from the following examples, which are presented herein for illustration only and should not be construed as limiting the invention in any way.

- 5 For the following experiments, "OP-1" is used interchangeably with "sBMP-7."

Example 1 Nephrectomy Chronic Renal Failure Injury Model

A rat partial (5/6) nephrectomy or rat remnant kidney model (RRKM) model was employed essentially as described (Vukicevic, et al. (1987) J. Bone Mineral
10 Res. 2: 533, the entire content of which is incorporated herein by reference). Male Munich-Wistar rats (2-3 months old, weighing about 200-250 g) were subjected to renal mass ablation. Specifically, the right kidney is excised in conjunction with selective ligation of the left renal artery branches such that one third of the kidney remains perfused (5/6 NPX, see Figure 12). Immediately following surgery, plasma
15 creatinine and BUN levels rise dramatically due to the loss of renal mass and function. Over the next several weeks of this "acute" failure phase, plasma creatinine and BUN levels of surviving animals decline somewhat toward normal values but remain elevated. Renal function then appears to remain relatively constant or stable for a period of variable duration. After this point, the animals enter a period of
20 chronic renal failure in which there is an essentially linear decline in renal function ending in death. For these reasons, animals are given 4 weeks of recovery prior to initiation of treatments.

Four weeks after nephrectomy, rats were divided into four treatment groups (see Table III), namely OP-1 (150 µg/kg body weight, 3X/week, by i.p. injection),
25 Vehicle (20 mM arginine/150 mM NaCl, 0.1% Tween-80, pH 9.0, 1 ml/kg body weight, 3X/week, by i.p. injection), enalapril (100 mg/L in drinking water, 8-16 mg/kg body weight), and OP-1 with enalapril. For the first two groups (OP-1 and Vehicle), animals were sacrificed 14 weeks after nephrectomy. For the last two groups (enalapril, and OP-1 with enalapril), animals were sacrificed 26 weeks after
30 nephrectomy. As surgical controls, rats were subjected to a "sham" operation in which the kidneys were decapsulated but no renal tissue was removed. Sham-treated

animals were sacrificed 26 weeks after operation. None of the rats died in any group during this study. Systolic blood pressure, urine protein and/or glomerulosclerosis were monitored at intervals between 4 and 26 weeks post surgery.

Table III. Experimental design in the Nephrectomy Chronic Renal Failure Injury Model.

Group	Animals (N)	Treatment	Duration
I	5	Sham	26 weeks
II	10	OP-1	Initiate treatment 4 weeks after nephrectomy
III	10	Vehicle	Terminate animals 14 weeks after nephrectomy
IV	10	OP-1 + enalapril	Initiate treatment 4 weeks after nephrectomy
V	10	Vehicle + enalapril	Terminate animals 26 weeks after nephrectomy

As compared to sham-operated controls, animals after Nephrectomy exhibited symptoms of higher blood pressure and higher proteinuria. While OP-1 did not dramatically lower the blood pressure level (see Figure 13), OP-1 significantly reduced the proteinuria level, from 91 mg/day to 71 mg/day ($P < 0.05$), after 14 weeks post-Nephrectomy (see Figure 14). When animals were co-treated with OP-1 and enalapril, there was no additional benefit in reducing the blood pressure of nephrectomized animals to normal level as compared to animals treated by the ACE inhibitor enalapril alone (see Figure 15). However, the combination of OP-1 and enalapril was more effective in reducing the proteinuria level than enalapril alone, 62 mg/day vs. 105 mg/day (see Figure 16).

In summary, the results from the Nephrectomy Chronic Renal Failure Injury Model demonstrate that OP-1 improves glomerular filtration rate, reduces glomerulosclerosis, and reduces proteinuria. Co-treatment of OP-1 and enalapril reduces late-stage proteinuria more than enalapril alone, thus may have a better effect on renal functions.

Example 2 Unilateral Ureteral Obstruction (UUO) Renal Fibrosis Model

This UUO model was employed essentially as described (Moller, et al. (1984) Virchows Arch 402: 209-237, the entire contents of which are incorporated herein by reference). Sprague-Dawley rats (about 250 g) underwent either a sham operation (ureter manipulated but not ligated) or unilateral ureteral ligation. Two
5 ligatures, 5 mm apart, were placed in the upper two-thirds of the ureter over a section of polyethylene tubing placed around the ureter (see Figure 17). The suture tied to obstruct the ureter was removed along with the tubing at day 5, relieving the obstruction. In this model, hypertension, proteinuria, and lipid dysregulation do not contribute to progressive nephron destruction, and glomerular injury is not prominent
10 early in the course of the injury produced. Uremia is avoided by the function of the contralateral kidney, which undergoes hypertrophy and hyperplasia as the obstructed kidney is destroyed. The renal injury of UUO is mediated in part through stimulation of renal angiotensin II production, which activates type-1 A-II Receptor and the downstream TGF- β in a cascade of events culminating in tubulointerstitial
15 inflammation and fibrosis. Inhibition of angiotensin II production by ACE inhibitors (or inhibition of A-II receptors) decreases expansion of the renal interstitium associated with fibrosis.

The effect of OP-1 on Renal fibrosis has been published (Hruska, et al. (2000) Am J Physiol Renal Physiol 279:F130-43, incorporated herein by reference),
20 and a summarization is as follows. Administration of OP-1 (100 or 300 μ g/kg body weight) prevented interstitial inflammation and fibrogenesis during the first 5 days after obstruction. Compared with ACE inhibition (by enalapril treatment), OP-1 was more effective in preventing tubulointerstitial fibrosis and in preserving renal function (see Figure 18). The mechanism of OP-1- induced renal protection was
25 associated with prevention of tubular atrophy, an effect not shared with enalapril (see Figure 19). OP-1 blocked the stimulation of epithelial cell apoptosis produced by UUO, which promoted maintenance of tubular epithelial integrity. OP-1 preserved renal blood flow (RBF) during UUO, but enalapril also stimulated RBF. OP-1 was more efficacious than enalapril in improving the glomerular filtration rate
30 as evidenced by the inulin clearance rate (see Figure 20). OP-1 also inhibited tubular epithelial disruption stimulated by the renal injury of UUO. Additional effects of OP-1 have been observed in this rat UUO model. For example, OP-1, but not ACE

inhibitor, significantly reduced the loss of medullary tissue in the kidney (see Figure 21), from about 24% to about 16% (OP-1 at 100 µg/kg) or about 13% (OP-1 at 300 µg/kg) ($P < 0.01$).

5 In summary, the results from the UUO Renal Fibrosis Model demonstrate that OP-1 reduces renal fibrosis, increases the glomerular filtration rate, and reduces tubular atrophy and medullary necrosis. OP-1 is more effective than enalapril. However, the combination of OP-1 and enalapril may have additive benefits on renal functions.

Example 3 Streptozotocin-Induced Diabetic Nephropathy Model

10 Nephropathy is one of the most common and most serious complications in type 1 diabetes mellitus. Renal involvement usually starts with renal hypertrophy and glomerular hyperfiltration, which can be observed soon after diabetes onset (Mogensen, et al. (1994) Diabetes Care 17:770-775). Glomerular hyperfiltration is often accompanied by a loss of renal functional reserve. After some years,
15 microalbuminuria (30 to 300 mg/day) may occur as well as morphological changes such as thickening of the glomerular basement membrane and mesangial expansion. The albumin leakage may subsequently become aggravated and overt nephropathy with albuminuria (>300 mg/day) may develop, usually 10 to 20 years after the onset of diabetes. At this time, hypertension becomes more common. Nephrotic syndrome
20 may occur, and glomerular filtration rate declines. The most important therapeutic measures undertaken to avoid, or retard, the progress of nephropathy aim to improve glycemic control and normalize blood pressure. ACE inhibitors have proven effective in the latter respect.

Streptozotocin kills pancreatic β cells and induces type I diabetes (for
25 review, see Cheta et al. (1998) J Pediatr Endocrinol Metab 11:11-9). It is widely used to induce experimental diabetic nephropathy in animals (see Figure 22). Adult female Sprague-Dawley rats (weighing 200-250 g) were intraperitoneally injected with streptozotocin (60 mg/kg body weight) to induce hyperglycemia. Hyperglycemic rats then received daily injections of insulin to maintain blood
30 glucose between 200-400 mg/dL. At week 16 when renal function declined, animals were treated weekly with OP-1 (10, 30 or 100 µg/kg body weight), enalapril (50 or

100 mg/L in drinking water) or a combination of OP-1 and enalapril. Control animals without streptozotocin treatment were handled in all other ways like treated animals. Animals were sacrificed at week 32 post-streptozotocin treatment.

5 Animals treated with streptozotocin exhibited symptoms of lower glomerular
filtration rate and higher proteinuria level. As seen in Figure 23, OP-1, but not
enalapril, significantly increased the glomerular filtration rate (GFR). While the
mean GFR in the 32-week diabetic animals was about 0.38 ml/min/100 g body
weight, the mean GFR in the animals treated with 100 µg/kg of OP-1 was about 0.7
ml/min/100 g body weight ($P<0.05$). The mean GFR in the animals co-treated with
10 OP-1 and enalapril was even higher, about 0.75 ml/min/100 g body weight ($P<0.05$).
As seen in Figure 24, OP-1 (30 and 100 µg/kg), enalapril (50 and 100 mg/L) or the
combination of OP-1 and enalapril, significantly reduced the proteinuria level from
about 140 mg/day to almost normal level.

15 In summary, the results from Streptozotocin-Induced Diabetic Nephropathy
Model demonstrate that OP-1 increases the glomerular filtration rate and reduces
proteinuria. The combinatorial treatment of OP-1 and enalapril may have better
effects on renal functions.

Example 4 Alloxan-Induced Diabetic Nephropathy Model

20 Similar to streptozotocin, alloxan kills pancreatic β cells and induces type I
diabetes (for review, see Cheta et al. (1998) J Pediatr Endocrinol Metab 11:11-9).
Alloxan is also widely used to induce experimental diabetic nephropathy in animals
(Figure 25). Adult female Sprague-Dawley rats (weighing 200-250 g) were
intraperitoneally injected with alloxan (70 mg/kg body weight) to induce
hyperglycemia. Renal arteries were clamped just prior to injection to prevent direct
25 nephrotoxicity, and then clamps were removed 5 minutes after injection. At week 16
when renal function declined, animals were treated with OP-1 (10 µg/kg body
weight, 3X/week, or 30 µg/kg body weight, 1X/week), enalapril (100 mg/L in
drinking water) or the combination of OP-1 and enalapril. Control animals without
alloxan treatment were handled in all other ways like treated animals. The treatment
30 duration was 12 weeks. All rats were sacrificed after 26 weeks.

Animals treated with alloxan showed higher serum creatinine level and higher proteinuria after 12-weeks of treatment. As seen in Figure 26, OP-1 (10 or 30 $\mu\text{g/kg}$) dramatically reduced the serum creatinine level, from about 115 $\mu\text{mole/L}$ to about 65 $\mu\text{mole/L}$ or 55 $\mu\text{mole/L}$ ($P < 0.01$). Compared with OP-1, elanapril reduced the serum creatinine level at a lesser degree. The combination of OP-1 and elanapril also significantly reduced the serum creatinine level. As seen Figure 27, OP-1 (10 or 30 $\mu\text{g/kg}$) or enalapril reduced the proteinuria level, from about 180 mg/dL/24 hr to about 80 (or 110) or 140 mg/dL/24 hr , respectively. In contrast, the combination of OP-1 and elanapril dramatically reduced the proteinuria level to as low as about 30 mg/dL/24 hr ($P < 0.01$).

In summary, the results from Alloxan-Induced Diabetic Nephropathy Model demonstrate that OP-1 or enalapril decreases the serum creatinine level and reduces the proteinuria level. The combinatorial treatment of OP-1 and enalapril shows better effects on renal functions.

Example 4 Bone Morphogenic Protein-7 (BMP-7), a Novel Effective Therapy For Diabetic Nephropathy

Overview

A long-term streptozotocin model of diabetic nephropathy was used to test and compare the therapeutic actions of BMP-7 with those of Enalapril. The study design was a treatment protocol beginning at 16 weeks when glomerular hypertrophy and proteinuria were established. The effects of therapy with BMP-7 (10, 30, or 100 $\mu\text{g/kg}$ iv, biw) were compared to a maximal dose of Enalapril (20 mg/kg) and to a vehicle control. The highest dose of BMP-7 and Enalapril were equal in partially reversing kidney hypertrophy. In the diabetic rats treated with BMP-7, 100 $\mu\text{g/kg}$, GFR at 32 weeks was significantly higher than in the diabetic vehicle treated rats, 0.7 ± 0.08 vs 0.34 ± 0.02 ml/min/100g bw ($P < 0.05$). The GFR of 32 week diabetic Enalapril treated rats was 0.58 ± 0.06 (not significant compared to vehicle treated and sham injected rats 0.55 ± 0.02). Albuminuria was reversed to normal by BMP-7 in a dose dependent manner.

The reduction in proteinuria by the intermediate dose of BMP-7 was similar to the effect of Enalapril therapy. Glomerular area and interstitial volume were significantly decreased in the BMP-7 and Enalapril treated animals. Glomerular sclerosis was prevented by BMP-7 therapy more effectively than by Enalapril. The first insight into the mechanism of BMP-7 therapy, was produced by analysis of blood pressures. Enalapril controlled hypertension throughout the course of therapy while BMP-7 did not effect blood pressure until the final four weeks of therapy. Important mechanistic insight derived from the demonstration that lost epithelial cell differentiation marked by loss at hyperglycemic vehicle treated diabetic rats BMP-7 expression in the kidney. At the same time another developmental morphogen, Wnt4, was widely expressed. BMP-7 and Enalapril therapy restored BMP-7 expression at high levels in the collecting duct without affecting Wnt4 expression. We conclude that BMP-7 reversed diabetic and hyperglycemia induced glomerular hypertrophy and injury, restoring GFR, protein excretion in glomerular histology towards normal and generally outperforming Enalapril. Restoration of BMP-7 expression, representing preservation of the collecting duct phenotype, restored the normal developmental Wnt4 interacting partner. This was associated with a successful repair reaction and a reversal of the ill-fated injury response.

Introduction

Recent studies have made the surprising observation that a renal tubular developmental morphogen, bone morphogenetic protein-7 (BMP-7), was effective in preventing the tubulointerstitial nephritis stimulated by obstructive uropathy (1). The mechanism of action appeared to be preservation of epithelial cell phenotype, inhibition of epithelial-mesenchymal transdifferentiation and inhibition of injury induced epithelial cell apoptosis. These actions of BMP-7 are reminiscent of the effects that the morphogen exercises during development.

During vertebrate development, the permanent kidney is generated by the interactions of the ureteric bud and the metanephric mesenchyme (2,3). At day 11 post-coitum (dpc) in the mouse, the ureteric bud branches out and invades the metanephric mesenchyme. Thereafter, nephrogenesis derives from reciprocal inductive interaction between these two tissues. The metanephric mesenchyme

induces the ureteric bud to grow and bifurcate to form the collecting ducts. At the same time permissive survival signals from the ureteric bud interact with mesenchymal signals to induce the conversion of metanephric mesenchyme into an epithelial structure. Epithelialization begins at day 11.5 pc with condensation around
5 the ureteric bud, and it progresses with the condensed mesenchyme segregating into pretubular aggregates. Epithelialization of these aggregates leads to development of the comma shaped bodies, S-shaped bodies and eventually the epithelial component of the nephron including glomerular podocytes.

BMP-7 has been shown to be a required factor leading to the condensation
10 and epithelialization of the metanephric mesenchyme, and the reciprocal induction of collecting duct differentiation (4-6). It is expressed in the ureteric bud, and in the condensing mesenchyme at day 11.5. BMP-7 is a survival factor for the condensing mesenchymal cells which die between 12.5 and 14.5 dpc in its absence (4). At day 12 pc, in the absence of BMP-7, glomerulus formation ceases as the mesenchymal
15 cells apoptose. BMP-7 interacts with another critical tubular inductive morphogen, Wnt4 (7) (8). Wnt4 expression is initiated at day 12.5 in the aggregating mesenchyme and pretubular aggregates. It is required as an inductive signal for epithelization (7). At birth, BMP-7 deficient kidneys are dysgenic, hypoplastic and cystic with severely dilated collecting ducts separated by areas of stromal cells an
20 extracellular matrix. The kidneys are hydronephrotic, and they do not have metanephric mesenchyme or evidence of glomerulus formation in the cortical nephrogenic zone (4,5). Cysts appear to originate from derivatives of the ureteric bud suggesting abnormal activity of the cell cycle and disordered polarity in these cells. Glomerular density is less than 3/ section compared to greater than 100/section
25 in wild type kidneys (8). Wnt4 deficient kidneys, on the other hand are also dysplastic and hydronephrotic, but they are totally devoid of glomeruli (8).

Compared to many other morphogens, expression of BMP-7 in the tubular epithelial segments derived from the ureteric bud does not cease following its developmental inductive actions, rather its expression persists and it likely functions
30 physiologically as a collecting duct epithelial cell differentiation factor. As such it

inhibits proliferation by blocking progress of the cell cycle at the G1 checkpoint and it prevents apoptosis (1,9).

5 Tissue injuries frequently stimulate attempts at repair that recapitulate development, including entry of surviving tissue cells into the cell cycle. This would require the actions of differentiation factors to be overcome perhaps by decreasing their levels. In turn, the absence of key factors could cause these attempted repairs to default into a fibrotic process resulting in permanent loss of the starting tissue. Unsuccessful injury repair characterizes many renal diseases that lead to loss of excretory function. Recently, renal injuries, including that caused by high glucose
10 levels, have been shown to decrease BMP-7 levels (1,10-12). Treatment of one of these injuries with BMP-7 prevented renal tubular atrophy and epithelial cell apoptosis. Failure of tubular development and mesenchymal apoptosis are features of development in the BMP-7 deficient state.

Here we report in a proof of concept study that BMP-7 is an effective
15 therapy for diabetic nephropathy. We report the discovery of a new mechanism of renal injury stimulated by diabetes, that of re-expression of a critical tubular epithelial inductive signal, Wnt4, that interacts with BMP-7 during development. Wnt4 is known to stimulate tubulointerstitial fibrosis, and its re-expression during diabetic hyperglycemic injury is a mechanism of synergism with TGF β resulting in
20 a failed injury repair reaction and promotion of disease. Therapy of diabetic nephropathy with BMP-7 partially reversed the renal injury induced by diabetes and hyperglycemia, and it prevented the development of glomerulosclerosis. The actions of BMP-7 were compared to the known therapeutic agent for diabetic nephropathy – angiotensin converting enzyme inhibition. While both agents were efficacious,
25 BMP-7 was more effective in reversing proteinuria and preventing glomerulosclerosis. Diabetic injury resulted in the loss of tubular epithelial and glomerular podocyte phenotype manifested by loss of BMP-7 expression, and therapy with BMP-7 restored the phenotype of the collecting duct manifested by restoration of BMP-7 expression. BMP-7, in turn, stimulated a successful repair
30 reaction possibly by interacting with Wnt4 and inhibiting the actions of TGF β .

Materials and Methods

Animals: Female Sprague-Dawley rats 10 weeks of age weighing 190 -220g were used. Animals were allowed free access to standard rat chow and to tap water. Diabetes mellitus was induced by a single tail-vein injection of Streptozotocin (STZ 62 mg/kg of body weight, Sigma chemical company, St. Louis, MO) dissolved in normal saline at day 0. The diabetic state was confirmed 72 hours later by the determination of blood glucose concentration >300 mg/dl. From day 3, all the diabetic rats received daily subcutaneous injections of 1.0-7.0 u human recombinant long-acting insulin injection (Eli Lilly & Co., Indianapolis, ID) as required to maintain the blood glucose concentration between 300-500 mg/dl. Tail blood glucose levels were measured twice a week with an Accu – Chek™ Advantage meter (Roche Diagnostic Corporation, Indianapolis, IN). Body weights were taken once a week. Food and water intake were monitored daily.

Renal hypertrophy was well developed at 16 weeks prior to treatment with vehicle, BMP-7, or enalapril. At 16 weeks, animals were divided into 8 groups. Group 1 were 16 weeks of DM. Group 2 were 16 weeks normal controls. Group 3 DM animals were treated with tail-vein injection of vehicle twice a week for 16 weeks. Group 4, Group 5, and Group 6 were DM animals treated with tail-vein injections of BMP-7 100, 30, and 10 µg/kg body weight respectively twice a week for 16 weeks. Group 7 were treated with Enalapril 20 mg/kg through the drinking water for 16 weeks. Group 8 animals were normal control showed for 32 weeks alongside the diabetic animals. Group 1 and Group 2 animals were sacrificed at 16 weeks. The others were sacrificed at 32 weeks. All treatments began at 16 weeks and continued through 32 weeks. Glomerular filtration rate (GFR), urine albumin excretion, kidney weight, glomerular area, mesangial matrix area, interstitial volume, monocyte and macrophage infiltration, thickness of glomerular basement membrane (GBM) and glomerulosclerosis were measured.

Renal Function: In all animals, glomerular filtration rate (GFR) was measured as the clearance of inulin. Rats were anesthetized with a Ketamine/Xylozine cocktail. A catheter was inserted into the femoral vein under a dissecting microscope for infusion. Another catheter was placed into the femoral artery for collecting blood samples. Urine was collected by bladder cannulation.

After the completion of surgery, a bolus of 2 ml/kg of 3% inulin (Cypros Pharmaceutical Corp, W. Carlsbad, CA) and 0.2% p-aminohippurate (PAH) in normal saline was infused as a priming load, followed by a sustaining infusion of the same solution at the rate of 8 ml/h.kg. Urine collection was initiated for three 20-minute collection periods after an hour of equilibration. An arterial blood sample was obtained during each clearance period. Plasma and urine were analyzed for inulin.

Urine Protein Excretion: Before the clearance study, the rats were placed in individual metabolic cages for two 24 hour urine collections. Urine volume was measured and urinary protein concentration was determined with a Bio-Rad protein assay.

Preparation of Kidneys: After the clearance studies, the rats were euthanized. Both kidneys were rapidly removed and placed in ice-cold phosphate buffered saline (PBS). Kidneys were weighed and then sliced on a cold glass plate. Two 2 mm coronal sections were immersed in Histochoice and in 10% buffered formalin. Kidney sections were embedded in paraffin and cut at 3 μ m and stained with hematoxylin and eosin, Gomori's Trichrome and periodic acid Schiff (PAS).

Renal Morphology: The morphometric analysis was done in a blind manner. OsteomeasureTM was used for morphometric analysis. Glomeruli were traced at x 400 magnification. Tissue sections stained by PAS were used. The measured glomerular parameters were as follows: (a) glomerular area, determined out of 60 glomeruli per group, which had vascular pole on it from randomly selected sections. (b) mesangial matrix area (defined as PAS-positive material in the mesangium), (c) ratio of the mesangial matrix area to the glomerular area, and (d) focal segmental glomerulosclerosis (glomerular sclerosis was defined as global sclerosis or segments of glomerular tufts demonstrating collapsed, obliterated capillaries with sparseness of normal cellular elements). The percent of sclerotic glomeruli was determined was determined out of 150 glomeruli per animal from randomly selected sections (13).

Interstitial volume was determined by a point-counting technique on tissue sections stained by the Gomori's Trichrome method, and was expressed as the mean

percentage of grid points laying within the interstitial area in up to 5 fields in the cortex (1) (14).

Quantitation of monocyte/macrophage infiltration was determined by counting ED-1 antigen positive cells in tissue sections (1) (14). Kidneys were fixed
5 in Histochoice, paraffin embedded, and sectioned. Sections were dewaxed and rehydrated prior to incubation with the ED-1 antibody obtained from Harlan (Indianapolis, IN). The location of the primary antibodies was visualized using alkaline phosphatase-linked second antibody. To obtain numbers of infiltrating monocyte/macrophage in the glomeruli, 50 consecutive cross-sections of glomeruli
10 of each animal were evaluated. To obtain numbers of infiltrating monocyte/macrophage in the renal cortical tubulointerstitium, 10 consecutive non-overlapping fields were counted in each section and viewed at 400x magnification.

Electron microscopy: Fresh tissue was fixed in 3% (wt/vol) glutaraldehyde buffer and post-fixed in OsO₄. Tissue was then dehydrated in ethanol and embedded
15 in Poly/bed 812 resin (Poly Science Inc, Warrington, PA). Thin sections were stained with uranyl acetate and lead citrate and examined using a transmission electron microscope.

Blood pressure determinations: The tail artery cuff method was used to follow blood pressures.

In situ hybridizations: ³⁵S-UPT labeled sense and antisense constructs were
20 prepared as previously described (15). Briefly frozen sections (4-6 microns) were fixed in 4% formaldehyde in PBA for 20 min at room temperature. Sections were washed once in PBS at three times the normal concentration of salt. Sections were then washed three times in PBS for 5 min and in water for 2 min and once in 0.1 M
25 triethanolamine, pH 8.0 for 10 min. Additional washings for 10 min in 0.25% acetic anhydride in 0.1 M triethanolamine, pH 8.0 and twice for 2 min in 2x standard sodium citrate were performed. The sections were then dehydrated through graded ethanols followed by air drying for 5 min. This was followed by vacuum drying for 1 h at room temperature. Each slide was then hybridized for 18 h at 60°C with 10⁶
30 counts/min of ³³P radiolabeled riboprobe in 80 µl of hybridization mixture (50% formamide; 2% Denhardt's solution; 10% dextran sulfate; 0.3 M NaCl; 10 mM Tris,

pH = 8.0; 2 mM EDTA, 0.25 g/ml tRNA). The sense and antisense probes for BMP-7 were a kind gift of Kuber Sampath. The sense and antisense probes for Wnt4 were previously described (15). Slides were then treated for 20 min in 4x SSC at room temperature; 4x SSC for 5 min at room temperature; Rnase A (20 µg/ml) for 30 min
5 at 37°C in 0.5 M NaCl, 0.01 M Tris, pH = 8.0 and 1 mM EDTA; twice 2 x SSC for 5 min at room temperature; 1x SSC for 10 min at room temperature; 0.5 x SSC for 10 min at room temperature; and 0.1 x SSC for 30 min at 60°C. The sections were then dehydrated using increasing concentrations of ethanol followed by vacuum desiccation for 30 min. Slides were dipped in liquid photographic emulsion (Kodak
10 NTB2) and exposed for 1 wk at 4°C. After development, slides were counterstained with hematoxylin and eosin.

Statistical Analysis: Results were expressed as mean \pm SEM. In situ method statistical Analysis was carried out by using a one-way ANOVA, or a nonparametric ANOVA. Statistical significance was achieved if the $p < 0.05$. Data were analyzed
15 using the InStat software.

Results

General Group Comparisons: The number of animals in each group, body weight at the beginning and the study end, kidney weight, blood glucose concentration, insulin dose, urine protein excretion, GFR and the mortality data of
20 each group are summarized in Table 1. There were no significant differences in body weight gain between the different groups during the 16 weeks of treatment. Also there were no significant differences in blood glucose levels and insulin doses between the different groups during the 16 weeks of treatment. The blood glucose levels were significantly higher in the 16 week DM group than those of all the 32
25 weeks treatment groups, and insulin doses in the 16 week DM were significantly lower than those of all the 32 weeks treatment groups ($p < 0.01$). We increased the insulin doses during the study to increase blood sugar control, in order to decrease the mortality we were encountering. Specifically, the blood sugar range was decreased from 400-600 to 300-500. As can be seen from Table 1, these actions
30 were equally applied to all groups. The mortality observed in the various treatment groups was similar. The causes of death were hypoglycemia, ketoacidosis, and

anesthesia. Some animals died of unknown reasons because the biochemical data on the day of death were lacking, and autopsies were not informative, suggesting ketoacidosis or hypoglycemia as the causes.

Table 1 Characterization of Animals in Various Groups

Group	N	Body Weight (g)			kidney weight (g)	Blood sugar (mg/dl)	Insulin Dose (u/wk)	Protein Ex (mg/day)	GFR (ml/min/100g bw)	Mortality	Cause-Mortality			
		Beginning	End	Weight Gain							keto-acidosis	hypo-glycemia	anes-thesia	un-known
NI 16	7		266.7±6.7		0.81±0.02			3.76±0.39	0.49±0.04	0/7				
NI 32	14		296.1±5.1		0.85±0.02			8.24±1.28	0.55±0.02	0/14				
DM 16	4				1.42±0.02	508.1±5.7*	8±1*	35.63±13.35	1.56±0.27	0/4				
DM 32	18	219.8±6.3	302.1±6.0	82.3±8.8	1.44±0.04	472.9±5.2	11±1	174.44±52.50	0.34±0.02	4/18		1/18	1/18	2/18
BMP 10	9	213.4±4.1	292.6±7.0	79.2±5.5	1.19±0.03	460.0±5.5	12±1	59.46±21.84	0.59±0.07	2/9				2/9
BMP 30	9	219.1±3.6	280.2±5.9	65.6±3.6	1.19±0.03	455.8±5.7	13±1	33.02±9.11	0.62±0.09	1/9			1/9	
BMP 100	10	204.8±4.9	290.3±6.6	85.5±6.9	1.10±0.03	461.6±5.5	16±1	14.27±3.50	0.70±0.08	1/10		1/10		
E 100	10	216.1±4.0	294.9±4.4	78.7±5.9	1.09±0.04	459.7±5.9	14±1	33.59±7.58	0.58±0.06	3/10	1/10	1/10		1/10
*p<0.01 vs the 32 wk groups; Other significant differences demonstrated in figures														
Values are means ± SEM														

5 *Characterization of the Streptozotocin Diabetic Nephropathy Model:* The effects of diabetes on kidney weight and GFR are shown in Figure 1. DM was induced at week 0. At 16 weeks of DM, kidney weight had increased 1.8 fold compared to normal (1.42 ± 0.02 versus 0.81 ± 0.02 g, $p<0.01$), while GFR increased 3.2-fold compared to normal (1.56 ± 0.27 versus 0.49 ± 0.04 ml/min/100g body wt, $p<0.01$). These data demonstrated that we had induced significant renal hypertrophy in this animal model. All treatments began at 16 weeks and continued through 32 weeks (or for an additional 16 weeks).

After 16 weeks of vehicle treatment, the kidney weights were not changed (1.44 ± 0.04 g versus 1.42 ± 0.02 g) (Figure 1). But the GFR of vehicle treated diabetic rats was decreased 75% to significantly lower than normal rats maintained for 32 weeks (0.34 ± 0.02 versus 0.55 ± 0.02 ml/min/100gbw, $p<0.05$). This demonstrated that without treatment there was a change from renal hyperfiltration to renal failure in control rats during the treatment period.

On the other hand, as shown in Figure 2, BMP-7 and enalapril treated animals had significantly decreased kidney weights (1.10 ± 0.03 in the BMP-7 high dose group, 1.09 ± 0.04 in enalapril treated group) compared to the diabetic vehicle-

treated rats ($p < 0.01$). There was a dose-dependent ordering of kidney weights in the BMP-7 treated group (1.19 ± 0.03 , 1.19 ± 0.03 , 1.10 ± 0.03 , respectively). These data demonstrated that BMP-7 and enalapril treatment partially reversed diabetic renal hypertrophy.

5 *Effects of Therapy on Kidney Hypertrophy:* While kidney weight was not affected by 16 weeks of vehicle treatment, GFR was reduced from 1.56 ± 0.27 ml/min/100 g bw to 0.34 ± 0.02 , which was 40% reduced from normal of 0.55 ± 0.02 ($p < 0.05$) (Figure 1 and 3). The GFR in the BMP-7 and enalapril treatment groups was normal or greater (Figure 3). The GFR of the BMP-7 high dose group was
10 significantly greater than the GFR of the DM group (0.70 ± 0.08 versus 0.34 ± 0.02 , $p < 0.01$). There was a dose-dependent ordering of GFR in the BMP-7 treated groups (0.59 ± 0.07 , 0.61 ± 0.09 , and 0.70 ± 0.08 , respectively). These data indicated that the vehicle control group had developed progressive renal failure during the 16 weeks of vehicle, while the treatment groups had partial reversal of renal
15 hypertrophy and prevention of further injury resulting in GFR significantly greater than normal at the high dose BMP-7 and dose dependently decreasing to normal with decreasing BMP-7 doses.

Urine Protein Excretion: The effects of diabetes and treatment on urine protein excretion are shown in Figure 4. Diabetic rats exhibited a pronounced
20 increase in protein excretion rate compared with nondiabetic rats at both 16 weeks (35.63 ± 13.35 versus 3.76 ± 0.39 mg/day) and 32 weeks (174.44 ± 52.50 versus 8.24 ± 1.28 , $p < 0.01$). This response to diabetes was reversed by BMP-7 and enalapril treatment ($p < 0.01$, DM versus BMP10; $p < 0.001$, DM versus BMP30, BMP100 and Enalapril). There was a significant dose-dependent ordering of protein
25 excretion in the BMP-7 treated groups (59.46 ± 21.84 , 33.02 ± 9.11 , and 14.27 ± 3.50 , respectively, $p < 0.05$ comparing low dose to high dose BMP-7).

Treatment Effects on Pathology: Compared to normal, kidneys of 16 week diabetic rats had massive glomerular hypertrophy to go along with the increase in kidney weights (Figure 5). The 16 week diabetic kidneys also had mild increases in
30 mesangial matrix and thickening of glomerular and tubular basement membranes (Figure 5) in agreement with previous studies (16). By 32 weeks of diabetes, the

kidneys of vehicle treated rats continued to be hypertrophied, but a significant component of glomerular area was sclerotic in a focal segmental pattern (Figures 5 and 23). The sclerotic areas demonstrated increased matrix, obliteration of capillaries and sparseness of cellular elements. Electron microscopy confirmed the increase in GBM thickness produced by diabetes by 16 weeks as shown in Figure 5 (data not shown). The focal-segmental nature of glomerular sclerosis in streptozotocin diabetic nephropathy is a known deviation from the human disease pathology of a diffuse global sclerosis with intermittent Kimmelstiel-Wilson nodular glomerular sclerosis lesions(17). This difference has been previously documented (13) (17).

Treatment with BMP-7 prevented the development of glomerulosclerosis along with a dose dependent reduction in the accumulation of mesangial matrix such that the intermediate and high dose BMP-7 treatments were associated with near normal glomerular histology except for incomplete resolution of glomerular hypertrophy (Figures 5-7). Enalapril therapy was less effective in preventing glomerulosclerosis, and it was less effective than intermediate or high dose BMP-7 in eliminating accumulation of mesangial matrix (Figure 5).

Morphometric Parameters: The effect of diabetes and the various treatments on glomerular area are shown in Figure 7A. Diabetic rats had a massively larger glomerular area than normal control rats (1.28 ± 0.03 versus $0.90 \pm 0.02 \times 10^4 \mu\text{m}^2$, $p < 0.001$) concordant with their increased kidney weights. All the treatments except vehicle partially reversed the glomerular hypertrophy ($p < 0.001$). The mesangial matrix area was also increased in diabetic rats compared to all the BMP-7 and enalapril treatment groups, but the ratio of mesangial matrix area to glomerular area were not different between different groups (data not shown).

As shown in Figure 7B, the cortical interstitial volume was increased from $9.0 \pm 0.6\%$ in nondiabetic rats to $13.1 \pm 0.7\%$ in the DM rats. High dose BMP-7 and Enalapril treatments significantly reduced the increase in interstitial volume to $10.7 \pm 0.3\%$; and $10.3 \pm 0.4\%$, respectively, $p < 0.01$. DM significantly increased the monocyte/macrophage infiltration both in the glomeruli (2.08 ± 0.24 vs. 1.01 ± 0.17 , $p < 0.01$) and in the cortical tubulointerstitium (7.25 ± 0.51 vs. 4.30 ± 0.35 , $p < 0.001$)

compared to normal rats. BMP-7 and Enalapril treatment decreased the tubulointerstitial infiltration.

The effect of diabetes, BMP-7, or enalapril treatment on glomerular pathology is shown in Fig. 8. Focal segmental glomerulosclerosis (FSGS) developed in diabetic rats compared with nondiabetic rats at 32 weeks ($10.7 \pm 4.0\%$ versus $0.7 \pm 0.2\%$, $p < 0.001$). This degree and type of glomerulosclerosis is similar to that reported in previous studies with STZ DM (13) (17). Glomerulosclerosis was markedly reduced by BMP-7 and less so by Enalapril treatment. There was a dose-dependent ordering of the reduction in glomerulosclerosis in the BMP-7 treated groups ($4.1 \pm 1.4\%$, $p < 0.05$; $3.5 \pm 0.8\%$, $p < 0.01$; and $2.1 \pm 0.3\%$, $p < 0.001$, respectively) (Fig. 8).

Mechanism of BMP-7 action: A significant component of the renoprotective actions of the positive control in these studies, Enalapril, is attributed to control of hypertension in diabetes (18) (19) (20,21). Therefore, we compared blood pressures between the various treatment groups in this study. As shown in Figure 9, the 16 week diabetic rats were mildly hypertensive in agreement with previous studies. Mean systolic blood pressures were 160 ± 3 compared to 141 ± 4 in normal control rats (17). By 20 weeks, Enalapril therapy had reversed the hypertension of the diabetic rats, and Enalapril therapy maintained normotension throughout the rest of the 32 weeks. In contrast, BMP-7 therapy, had no effect on blood pressure until after 28 weeks when it began to cause reductions in hypertension until, at 32 weeks, blood pressures were normalized in the BMP-7 high dose treated animals. At 32 weeks the diabetic vehicle treated rats exhibited worsening systolic hypertension (Fig. 9) and widening of their pulse pressures consistent with increasing loss of vascular pliability.

In further pursuit of the mechanism of the renoprotective actions of BMP-7 therapy, we reasoned that a critical pathogenetic action of diabetes might be induced loss of expression of a key differentiation factor/s, such as BMP-7, enabling the ill-fated injury response to be initiated. This turned out to be so as shown in Figure 10. In the normal rat kidney BMP-7 is expressed in the medulla, the cortical collecting ducts and glomerular podocytes (22). By 16 weeks of diabetes, BMP-7 expression

was absent from the kidney, and this absence was maintained during therapy with vehicle (Figure 10). Treatment with either BMP-7 or Enalapril restored BMP-7 expression in its normal pattern at high levels (Figure 10). The significance of the changes in BMP-7 expression is related to the expression of another critical kidney developmental morphogen, Wnt4, during renal injury. Wnt4 is normally expressed during kidney development following BMP-7 at the time of epithelialization of the condensing metanephric mesenchyme around the tip of the ureteric bud (7,8). Wnt4 along with a reciprocal signal from the ureteric bud (BMP-7) is required for epithelial differentiation of the mesenchyme and tubule formation of the condensing mesenchyme leading to formation of the comma and S-shaped bodies and glomerular development. However, in tubulointerstitial injuries, Wnt4 is re-expressed but in the absence of key factors (probably the reciprocal developmental differentiation factor) it promotes interstitial fibrosis (15). As shown in Figure 11, diabetic injury is associated with Wnt4 expression throughout the kidney, and BMP-7 and Enalapril therapy do not affect Wnt4 expression. During development Wnt4 and BMP-7 interact in the development of the nephron. In the absence of BMP-7 such as in renal injuries, Wnt4 re-expression promotes TGF β induced signaling(23), and this is the likely mechanism by which Wnt4 re-expression appears to promote renal fibrogenesis (15). As shown above, BMP-7 and Enalapril therapy stimulate a successful repair reaction and reinduction of BMP-7 expression. This may indicate that the Wnt4 actions were channeled into the successful repair reaction similar to its role in nephrogenesis.

Discussion

The data presented here demonstrate that the long-term model of streptozotocin induced diabetes is associated with the development of nephropathy and renal failure that resembles the clinical course of human diabetic nephropathy associated with Type I diabetes (24,25). By 16 weeks massive renal hypertrophy, hyperfiltration and proteinuria were established in our model. At that time, the earliest changes of glomerular mesangial matrix accumulation were detectable as previously reported (16). Over the ensuing 16 weeks during the course of the various therapies utilized here, diabetic renal injury either progressed or was partially

reversed. In the vehicle treated group, diabetic injury progressed leading to glomerular sclerosis, worsening of the proteinuria, the development of nephrotic syndrome and the development of renal insufficiency. The progression of the nephropathy to severe proteinuria and renal insufficiency was due to progression of the diabetic glomerular pathology to the level of 10% of the glomeruli being sclerotic. This severity of disease is similar to previous reports of glomerulosclerosis in STZ induced DM nephropathy (13,17,26-28).

In three groups of diabetic rats, BMP-7 was administered twice a week intravenously at (10, 30, and 100 µg/kg/bw). The effects of BMP-7 therapy were profound. BMP-7 partially restored kidney weights towards normal, restored glomerular filtration rate to normal, dose dependently eliminated proteinuria, partially reversed glomerular hypertrophy, reversed the increase in the expanded interstitial volume and prevented the development of glomerular sclerosis. The reversal of renal injury was the mechanism by which kidney weights, glomerular area, and glomerular filtration rate were reduced. All of the parameters of nephropathy that were assessed demonstrated dose ordering in their response that was small though present. Only the differences in protein excretion between the low dose and high dose groups achieved statistical significance.

The therapeutic effects of BMP-7 were further characterized by direct comparison to the known therapeutic actions of Enalapril, an angiotensin converting enzyme (ACE) inhibitor. ACE inhibition was first shown to be a renal disease therapeutic agent in STZ diabetes (18,19) (16) (20), and it is especially effective in human diabetic nephropathy (21,29) (30). Subsequently, along with AT-1 receptor blockade, it has become the main renal disease therapeutic for slowing progression of disease (29) (30). In the design of the studies reported here, a treatment of established disease was used. High dose BMP-7 was more effective than Enalapril in reversing proteinuria, maintaining GFR, decreasing mesangial matrix expansion and preventing the development of glomerular sclerosis. BMP-7 and Enalapril therapies were equally effective in reducing kidney weights, reversing glomerular hypertrophy and decreasing interstitial volume expansion. Enalapril therapy normalized blood pressure during the entire 16 week course of therapy. An important component of

the therapeutic effectiveness of Enalapril is estimated to be due to the control of systemic and glomerular hemodynamics (18-20). BMP-7 did not affect blood pressure until late in the course of the study, at a time when systolic hypertension was becoming prominent and vascular calcification was present (Davies and Hruska, data not published). We interpret these data to correlate with the actions of BMP-7 to prevent vascular calcification (31). The doses of Enalapril used here (20 mg/kg) were maximal (1,14,32). They were 10-50 fold greater than the doses used in clinical medicine. Preliminary studies demonstrated that there was no difference in the Enalapril response between 5 and 20 mg/kg. Furthermore, we have demonstrated that addition of Enalapril to the drinking water is equally effective as more arduous means of administering Enalapril such as interperitoneal injections and oral/esophageal gavage (Hruska, Wang unpublished data). Therefore, we conclude that BMP-7 therapy of streptozotocin induced diabetic nephropathy in the rat is at least equally effective to Enalapril therapy, and in the area of preventing progression to glomerular sclerosis clearly more effective in this study.

BMP-7 is a critical renal morphogen that is expressed in the normal adult kidney and lost through the influence of renal injury as shown here in diabetic nephropathy. Its therapeutic actions in renal disease appear to be related to recapitulation of its developmental actions. BMP-7 stimulates epithelial differentiation and provides metanephric mesenchymal survival signal during morphogenesis (4) (5) (6). This is similar to the actions of BMP-7 during renal tubulointerstitial injury (1). BMP-7 decreases expression of markers of injury response such as vimentin and $\alpha 1(I)$ procollagen, $\alpha 1(IV)$ collagen, and it increases expression of epithelial phenotype markers such as E-cadherin (1,5,9,12,22,33) (Hruska, unpublished). It prevents tubular epithelial apoptosis leading to injury prevention and disease resistance (1) (12). BMP-7 is expressed in the Wolffian duct at the time of ureteric bud development (4). It continues to be expressed in the ureteric bud and the developing collecting duct throughout development (34). In addition, BMP-7 is expressed in and required for the condensing metanephric mesenchyme at the tip of the ureteric bud and in the pretubular aggregates beginning at day 12.5(4). BMP-7 expression subsequently disappears from the comma and S-shaped bodies. In the adult kidney BMP-7 is expressed in glomerular podocytes, the

thick ascending limb, distal tubule, and collecting duct (22). The strongest expression is in the collecting duct especially in the medullary segments. In the absence of BMP-7, condensation of metanephric mesenchyme is diminished and mesenchymal cells apoptose between days 12 and 14 dpc (4). This is despite

5 expression of Wnt 4, another critical tubular developmental morphogen expressed in the condensing metanephric mesenchyme at the time of formation of the pretubular aggregates (8). Wnt 4 is required and sufficient for induction of tubulogenesis from day 14dpc through development of the early nephrons (8). Wnt 4 expression is diminished and absent from the Sshaped bodies following its inductive actions.

10 From their overlapping expression patterns between day 12.5 and 14dpc, the phenotype of their knockouts, and the overlapping times of their required presence, it is clear that Wnt4 and BMP-7 interact during renal development (4) (5) (7,8). In the adult kidney, Wnt4 expression is limited to the terminal medullary collecting duct (15). However, increased Wnt 4 expression is observed throughout the

15 collecting duct in response to renal injuries while BMP-7 expression is lost, as shown here and elsewhere (1,7,15). Here we show that diabetic nephropathy causes an even more widespread expression of Wnt4 than ureteral obstruction. Wnt4 expression in response to renal injury promotes tubulointerstitial fibrosis (15), and it promotes epithelial to mesenchymal transdifferentiation (EMT) similar to TGF β

20 (15,35). This is a critical mechanism of producing interstitial myofibroblasts and promoting fibrogenesis (36). These actions of WNT4 are similar to those of TGF β in renal injury. The mechanism of the relationship between Wnt4 and TGF β in renal injury is that Wnt 4 signaling stabilizes β -catenin and increases nuclear β -catenin levels (37,38). In the nucleus β -catenin binds in a transcriptional complex with

25 SMAD 4 (23). SMAD 4 in the nucleus exists as a dimer containing a regulatory SMAD. TGF- β induces the regulatory SMAD's 2 and 3, and the SMAD 2/4 and 3/4 dimers in the transcriptional complex associated with β -catenin and the TCF/LEF family regulate gene transcription. The Wnt4/ TGF β interaction leads to promotion of renal fibrosis and EMT (15) (23). Since TGF- β is increased in response to

30 diabetic injury (26,39) (27) (28), and BMP-7 is decreased (data presented here) (40), activation of Wnt4 leads to synergism with TGF- β (23). In the presence of BMP-7,

TGF- β signaling is inhibited (9). BMP-7 signaling interacts with Wnt 4 expression leading to the formation of transcriptional complexes containing the BMP-7 stimulated regulatory SMAD's 1,5, and 8 as in development. Thus, the transcriptional signaling induced by TGF- β is competitively inhibited. In addition,
5 BMP-7 stimulates the inhibitory SMAD, SMAD 6 (9), and in the proximal tubule, SMAD 7 (41), which further inhibit TGF- β induced signaling.

The STZ induced DM rat model and type I human DM overexpress TGF β 1,2, and 3 and the TGF β RII in renal tubular and glomerular cells(26,27,27,28,39,42). TGF β and the receptor together constitute a major biologic
10 signal inducing a switch toward a profibrotic cellular phenotype (43). Here we demonstrate that another profibrotic cytokine, Wnt4, that synergizes with TGF β is upregulated in STZ DM. We propose that Wnt4 upregulation may be a common response to renal injury, which, in the absence of BMP-7, stimulates a failed injury response through its interactions with TGF β .

15 In summary, the proof of concept studies reported here demonstrate successful treatment of streptozotocin induced diabetic nephropathy in renal insufficiency with BMP-7. BMP-7 effects were equal to and in some respects more potent than a positive control, Enalapril therapy. Diabetic injury induced diffuse Wnt-4 expression representing an injury response and reexpression of a
20 developmental morphogen. In the absence of BMP-7, Wnt4 may synergize with TGF β in stimulating diabetic injury. BMP-7 therapy restored tubular epithelial phenotype, BMP-7 expression, and inhibited TGF β actions. As a result a significant positive repair reaction was produced by BMP-7 and glomerular sclerosis was prevented as well as the development of renal insufficiency.

25 **Reference List**

1. Hruska,K.A., Guo,G., Wozniak,M., Martin,D., Miller,S., Liapis,H., Loveday,K., Klahr,S., Sampath,T.K., and Morrissey,J. 2000. Osteogenic protein-1 (OP-1) prevents renal fibrogenesis associated with ureteral obstruction. Am J Phys (Renal) 279:F130-F143.

2. Grobstein,C. 1953. Morphogenetic interactions between embryonic mouse tissues separated by a membrane filter. *Nature* 172:869-871.
3. Saxen,L. 1987. *Organogenesis of the Kidney*. Cambridge University Press, Cambridge, UK.
- 5 4. Luo,G., Hofmann,C., Bronckers,A.L., Sohocki,M., Bradley,A., and Karsenty,G. 1995. BMP-7 is an inducer of nephrogenesis, and is also required for eye development and skeletal patterning. *Genes Dev.* 9:2808-2820.
- 10 5. Dudley,A.T., Lyons,K.M., and Robertson,E.J. 1995. A requirement for bone morphogenetic protein-7 during development of the mammalian kidney and eye. *Genes Dev* 9:2795-2807.
6. Vukicevic,S., Kopp,J.B., Luyten,F.P., and Sampath,T.K. 1996. Induction of nephrogenic mesenchyme by osteogenic protein 1 (bone morphogenetic protein 7). *Proc. Natl. Acad. Sci. U.S.A.* 93:9021-9026.
- 15 7. Stark,K., Vainio,S., Vassileva,G., and McMahon,A.P. 1994. Epithelial transformation of metanephric mesenchyme in the developing kidney regulated by Wnt-4. *Nature* 372 :679-683.
8. Kispert A., Vainio S., and McMahon,A.P. 1998. Wnt-4 is a mesenchymal signal for epithelial transformation of metanephric mesenchyme in the
20 developing kidney. *Development* 125:4225-4234.
9. Dorai,H., Vukicevic,S., and Sampath,T.K. 2000. Bone morphogenetic protein-7 (osteogenic protein-1) inhibits smooth muscle cell proliferation and stimulates the expression of markers that are characteristic of SMC phenotype in vitro. *J Cellular Physiol* 184:37-45.
- 25 10. Simon,M., Maresh,J.G., Harris,S.E., Hernandez,J.D., Arar,M., Olson,M.S., and Abboud,H.E. 1999. Expression of bone morphogenetic protein-7 mRNA in normal and ischemic adult rat kidney. *Amer.J.Physiol.* 276:F382-F389.

11. Wang,S. and Hirschberg,R. 2000. Loss of renal tubular BMP7 during the evolution of experimental diabetic nephropathy. *J.Am.Soc.Nephrol.* 11:655A (Abstr.)
12. Vukicevic,S., Basic,V., Rogic,D., Basic,N., Shih,M.-S., Shepard,A., Jin,D.,
5 Dattatreymurthy,B., Jones,W., Dorai,H. et al. 1998. Osteogenic protein-1 (Bone 30 morphogenetic protein-7) reduces severity of injury after ischemic acute renal failure in rat. *J.Clin.Invest.* 102:202-214.
13. Lal,M.A., Körner,A., Matsuo,Y., Zelenin,S., Cheng,S.X.J., Jaremko,G.,
DiBona,G.F., klöf,A.-C., and Aperia,A. 2000. Combined antioxidant and
10 COMT inhibitor treatment reverses renal abnormalities in diabetic rats. *Diabetes* 49:1381-1389.
14. Ishidoya,S., Morrissey,J., McCracken,R., Reyes,A., and Klahr,S. 1995. Angiotensin II receptor antagonist ameliorates renal tubulointerstitial fibrosis caused by unilateral ureteral ligation. *Kidney Int.* 47:1285-1294.
- 15 15. Surendran,K., McCaul,S.P., and Simon,T.C. 2002. A role for Wnt-4 in renal fibrosis. *Am J Physiol Renal Physiol* 282:F431-F441.
16. Sassy-Prigent,C., Heudes,D., Jouquey,S., Auberval,D., Belair,M.-F.,
Michel,O., Hamon,G., Bariety,J., and Bruneval,P. 1995. Morphometric
detection of incipient glomerular lesions in diabetic nephropathy in rats. *Lab*
20 *Invest* 73:64-71.
17. Remuzzi,A., Perico,N., Amuchastegui,c.s., Malanchini,B., Mazerska,M.,
Battaglia,C., Bertani,T., and Remuzzi,G. 1993. Short- and long-term effect of angiotensin II receptor blockade in rats with experimental diabetes. *J Am Soc Neph* 4:40-49.
- 25 18. Zatz,R., Dunn,B.R., Meyer,T.W., Anderson,S., Rennke,H.G., and Brenner,B.M. 1986. Prevention of diabetic glomerulopathy by pharmacological amelioration of glomerular capillary hypertension. *J.Clin.Invest.* 77:1925-1930.

19. Anderson,S., Rennke,H.G., Garcia,D.L., and Brenner,B.M. 1989. Short and long-term effects of antihypertensive therapy in the diabetic rat. *Kidney International* 36:526-536.
20. Katoh,M., Ohmachi,Y., Kurosawa,Y., Yoneda,H., Tanaka,N., and Narita,H.
5 2000. Effects of imidapril and captopril on streptozotocin-induced diabetic nephropathy in mice. *Eur J Pharmacol* 398:381-387.
21. Lewis,E.J., Hunsicker,L.G., Bain,R.P., and Rohde,R.D. 1993. The effect of angiotensin converting-enzyme inhibition on diabetic nephropathy. *NEJM* 329:1456-1462.
- 10 22. Gould,S.E., Day,M., Jones,S., and Dorai,H. 2002. BMP-7 regulates chemokine, cytokine, and hemodynamic gene expression in proximal tubule cells. *Kidney International* 61:51-60.
23. Nishita,M., Hashimoto,M.K., Ogata,S., Laurent,M.N., Ueno,N., Shibuya,H., and Cho,K.W.Y. 2000. Interaction between Wnt and TGF- β signalling
15 pathways during formation of Spemann's organizer. *Nature* 403:781-785.
24. Sharma,K., Yulin,J., Guo,J., and Ziyadeh,J. 1996. Neutralization of TGF-B by anti-TGF- β antibody attenuates kidney hypertrophy and the enhanced extracellular matrix gene expression in streptozotocin-diabetic mice. *Diabetes* 45:522-530. 31
- 20 25. Wilson,G.L. and Letter,E.H. 1990. Streptozotocin interactions with pancreatic beta cells andthe inductionof insulin-dependent diabetes. *Curr Top Microbiol Immunol* 156:27-54.
26. Yamamoto,T., Nakamura,T., Noble,N.A., Ruoslahti,E., and Border,W. 1993. Expression of transforming growth factor β is elevated in human and
25 experimental diabetic nephropathy. *Proc.Natl.Acad.Sci.U.S.A.* 90:1814-1818.
27. Nakamura,T., Fukui,M., Ebihara,I., Osada,S., Nagaoka,I., Tomino,Y., and Koide,H. 1993. mRNA expression of growth factors in glomeruli from diabetic rats. *Diabetes* 42:450-456.

28. Shankland,S.J. and Scholey,J.W. 1994. Expression of transforming growth factor B1 during diabetic renal hypertrophy. *Kidney International* 46:430-442.
29. Mathiesen,E.R., Hommel,E., Giese,J., and Parving,H.H. 1991. Efficacy of captopril in postponing nephropathy in normotensive insulin dependent diabetic patients with microalbuminuria. *British Medical Journal* 303:81-87.
30. Wiegmann,T.B., Herron,K.G., Chonko,A.M., MacDougall,M.L., and Moore,W.V. 1992. Effect of angiotensin-converting enzyme inhibition on renal function and albuminuria in normotensive type I diabetic patients. *Diabetes* 41:62-67.
31. Davies,M.R., Petrosova,T., and Hruska,K.A. 2001. BMP-7 is an effective treatment for vascular calcification in chronic kidney disease (CKD). *J.Am.Soc.Nephrol.* (Abstr.)
32. Morrissey,J., Hruska,K., Guo,G., Wang,S., Chen,Q., and Klahr,S. 2002. Bone morphogenetic protein-7 improves renal fibrosis and accelerates the return of renal function. *J.Am.Soc.Nephrol.* 13:S14-S21.
33. Zeisberg,M., Maeshima,Y., Hanai,J.-I., Strutz,F., Dorai,H., and Kalluri,R. 2001. Bone morphogenic protein-7, an inducer of tubulogenesis in kidney development, is an inhibitor of chronic progressive renal disease. *J Am Soc Neph* 12:830A (Abstr.)
34. Lyons,K.M., Hogan,B.L., and Robertson,E.J. 1995. Colocalization of BMP 7 and BMP 2 RNAs suggests that these factors cooperatively mediate tissue interactions during murine development. *Mech.Dev.* 50:71-83.
35. Hay,E.D. and Zuk,A. 1995. Transformations between epithelium and mesenchyme: normal, pathological, and experimentally induced. *Am J Kidney Dis* 26:678-690.
36. Iwano,M., Plieth,D., Danoff,T.M., Xue,C., Okada,H., and Neilson,E.G. 2002. Evidence that fibroblasts derive from epithelium during tissue fibrosis. *J.Clin.Invest.* 110:341-350.

37. Cox,R.T. and Peifer,M. 1998. Wingless signaling: the inconvenient complexities of life. *Current Biology* 8:R140-R144.
38. Hartmann,C. and Tabin,C.J. 2000. Dual roles of Wnt signaling during chondrogenesis in the chicken limb. *Development* 127:3141-3159. 32
- 5 39. Hill,C., Flyvbjerg,A., Grønbæk,H., Petrik,J., Hill,D.J., Thomas,C.R., Sheppard,M.C., and Logan,A. 2000. The renal expression of transforming growth factor- β isoforms and their receptors in acute and chronic experimental diabetes in rats. *Endocrinology* 141:1196-1208.
40. Wang,S.-N., Lapage,J., and Hirschberg,R. 2001. Loss of tubular bone morphogenetic protein-7 in diabetic nephropathy. *J.Am.Soc.Nephrol.* 12:2392-2399. 10
41. Hill-Kapturczak,N., Truong,L., Thamilselvan,V., Visner,G.A., Nick,H.S., and Agarwall,A. 2000. Smad7-dependent regulation of heme oxygenase-1 by transforming growth factor- β in human renal epithelial cells. *J.Biol.Chem.* 275:40904-40909. 15
42. Sharma,V.K., Bologa,RM., Xu,G.-P., Li,B., Mouradian,J., Wang,J., Serur,D., Rao,V., and Suthanthiran,M. 1996. Intragraft TGF- β 1 mRNA: A correlate of interstitial fibrosis and chronic allograft nephropathy. *Kidney International* 49:1297-1303.
- 20 43. Sharma,K. and Ziyadeh,F.N. 1995. Hyperglycemia and diabetic kidney disease. The case for transforming growth factor-beta as a key mediator. *Diabetes* 44:1139-1146.

Equivalents

- 25 The invention may be embodied in other specific forms without departing from the spirit or essential characteristics thereof. The foregoing embodiments are therefore to be considered in all respects illustrative rather than limiting on the invention described herein. Scope of the invention is thus indicated by the appended claims rather than by the foregoing description, and all changes which come within

the meaning and range of equivalency of the claims are intended to be embraced therein.

SEQUENCE LISTING

SEQ ID NO:1

Ser Thr Gly Ser Lys Gln Arg Ser Gln Asn Arg Ser Lys Thr Pro Lys
1 5 10 15

Asn Gln Glu Ala Leu Arg Met Ala Asn Val Ala Glu Asn Ser Ser Ser
20 25 30

Asp Gln Arg Gln Ala Cys Lys Lys His Glu Leu Tyr Val Ser Phe Arg
35 40 45

Asp Leu Gly Trp Gln Asp Trp Ile Ile Ala Pro Glu Gly Tyr Ala Ala
50 55 60

Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asn Ser Tyr Met Asn
65 70 75 80

Ala Thr Asn His Ala Ile Val Gln Thr Leu Val His Phe Ile Asn Pro
85 90 95

Glu Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln Leu Asn Ala Ile
100 105 110

Ser Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Ile Leu Lys Lys Tyr
115 120 125

Arg Asn Met Val Val Arg Ala Cys Gly Cys His
130 135

SEQ ID NO:2

His Arg Arg Leu Arg Ser Gln Glu Arg Arg Glu Met Gln Arg Glu Ile
1 5 10 15

Leu Ser Ile Leu Gly Leu Pro His Arg Pro Arg Pro His Leu Gln Gly
20 25 30

Lys His Asn Ser Ala Pro Met Phe Met Leu Asp Leu Tyr Asn Ala Met
35 40 45

Ala Val Glu Glu Gly Gly Gly Pro Gly Gly Gln Gly Phe Ser Tyr Pro
50 55 60

Tyr Lys Ala Val Phe Ser Thr Gln Gly Pro Pro Leu Ala Ser Leu Gln
65 70 75 80

Asp Ser His Phe Leu Thr Asp Ala Asp Met Val Met Ser Phe Val Asn
85 90 95

Leu

SEQ ID NO:3

Met His Val Arg Ser Leu Arg Ala Ala Ala Pro His Ser Phe Val Ala
1 5 10 15

Leu Trp Ala Pro Leu Phe Leu Leu Arg Ser Ala Leu Ala Asp Phe Ser
20 25 30

Leu Asp Asn Glu Val His Ser Ser Phe Ile His Arg Arg Leu Arg Ser
35 40 45

Gln Glu Arg Arg Glu Met Gln Arg Glu Ile Leu Ser Ile Leu Gly Leu
50 55 60

Pro His Arg Pro Arg Pro His Leu Gln Gly Lys His Asn Ser Ala Pro
65 70 75 80

Met Phe Met Leu Asp Leu Tyr Asn Ala Met Ala Val Glu Glu Gly Gly
85 90 95

Gly Pro Gly Gly Gln Gly Phe Ser Tyr Pro Tyr Lys Ala Val Phe Ser
100 105 110

Thr Gln Gly Pro Pro Leu Ala Ser Leu Gln Asp Ser His Phe Leu Thr
115 120 125

Asp Ala Asp Met Val Met Ser Phe Val Asn Leu Val Glu His Asp Lys
130 135 140

Glu Phe Phe His Pro Arg Tyr His His Arg Glu Phe Arg Phe Asp Leu
145 150 155 160

Ser Lys Ile Pro Glu Gly Glu Ala Val Thr Ala Ala Glu Phe Arg Ile
165 170 175

Tyr Lys Asp Tyr Ile Arg Glu Arg Phe Asp Asn Glu Thr Phe Arg Ile
180 185 190

Ser Val Tyr Gln Val Leu Gln Glu His Leu Gly Arg Glu Ser Asp Leu
195 200 205

Phe Leu Leu Asp Ser Arg Thr Leu Trp Ala Ser Glu Glu Gly Trp Leu
210 215 220

Val Phe Asp Ile Thr Ala Thr Ser Asn His Trp Val Val Asn Pro Arg
225 230 235 240

His Asn Leu Gly Leu Gln Leu Ser Val Glu Thr Leu Asp Gly Gln Ser
245 250 255

Ile Asn Pro Lys Leu Ala Gly Leu Ile Gly Arg His Gly Pro Gln Asn
260 265 270

Lys Gln Pro Phe Met Val Ala Phe Phe Lys Ala Thr Glu Val His Phe
275 280 285

Arg Ser Ile Arg Ser Thr Gly Ser Lys Gln Arg Ser Gln Asn Arg Ser
290 295 300

Lys Thr Pro Lys Asn Gln Glu Ala Leu Arg Met Ala Asn Val Ala Glu
305 310 315 320

Asn Ser Ser Ser Asp Gln Arg Gln Ala Cys Lys Lys His Glu Leu Tyr
325 330 335

Val Ser Phe Arg Asp Leu Gly Trp Gln Asp Trp Ile Ile Ala Pro Glu
340 345 350

Gly Tyr Ala Ala Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asn
355 360 365

Ser Tyr Met Asn Ala Thr Asn His Ala Ile Val Gln Thr Leu Val His
370 375 380

Phe Ile Asn Pro Glu Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln
385 390 395 400

Leu Asn Ala Ile Ser Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Ile
405 410 415

Leu Lys Lys Tyr Arg Asn Met Val Val Arg Ala Cys Gly Cys His
420 425 430

SEQ ID NO :4

Ser Thr Gly Gly Lys Gln Arg Ser Gln Asn Arg Ser Lys Thr Pro Lys
1 5 10 15

Asn Gln Glu Ala Leu Arg Met Ala Ser Val Ala Glu Asn Ser Ser Ser
20 25 30

Asp Gln Arg Gln Ala Cys Lys Lys His Glu Leu Tyr Val Ser Phe Arg
35 40 45

Asp Leu Gly Trp Gln Asp Trp Ile Ile Ala Pro Glu Gly Tyr Ala Ala
50 55 60

Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asn Ser Tyr Met Asn
65 70 75 80
Ala Thr Asn His Ala Ile Val Gln Thr Leu Val His Phe Ile Asn Pro
85 90 95
Asp Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln Leu Asn Ala Ile
100 105 110
Ser Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Ile Leu Lys Lys Tyr
115 120 125
Arg Asn Met Val Val Arg Ala Cys Gly Cys His
130 135

SEQ ID NO:5

Ala Val Arg Pro Leu Arg Arg Arg Gln Pro Lys Lys Ser Asn Glu Leu
1 5 10 15
Pro Gln Ala Asn Arg Leu Pro Gly Ile Phe Asp Asp Val His Gly Ser
20 25 30
His Gly Arg Gln Val Cys Arg Arg His Glu Leu Tyr Val Ser Phe Gln
35 40 45
Asp Leu Gly Trp Leu Asp Trp Val Ile Ala Pro Gln Gly Tyr Ser Ala
50 55 60
Tyr Tyr Cys Glu Gly Glu Cys Ser Phe Pro Leu Asp Ser Cys Met Asn
65 70 75 80
Ala Thr Asn His Ala Ile Leu Gln Ser Leu Val His Leu Met Lys Pro
85 90 95
Asn Ala Val Pro Lys Ala Cys Cys Ala Pro Thr Lys Leu Ser Ala Thr
100 105 110
Ser Val Leu Tyr Tyr Asp Ser Ser Asn Asn Val Ile Leu Arg Lys His
115 120 125
Arg Asn Met Val Val Lys Ala Cys Gly Cys His
130 135

SEQ ID NO:6

Ala Ala Arg Pro Leu Lys Arg Arg Gln Pro Lys Lys Thr Asn Glu Leu
1 5 10 15
Pro His Pro Asn Lys Leu Pro Gly Ile Phe Asp Asp Gly His Gly Ser
20 25 30
Arg Gly Arg Glu Val Cys Arg Arg His Glu Leu Tyr Val Ser Phe Arg
35 40 45

Asp Leu Gly Trp Leu Asp Trp Val Ile Ala Pro Gln Gly Tyr Ser Ala
50 55 60
Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asp Ser Cys Met Asn
65 70 75 80
Ala Thr Asn His Ala Ile Leu Gln Ser Leu Val His Leu Met Lys Pro
85 90 95
Asp Val Val Pro Lys Ala Cys Cys Ala Pro Thr Lys Leu Ser Ala Thr
100 105 110
Ser Val Leu Tyr Tyr Asp Ser Ser Asn Asn Val Ile Leu Arg Lys His
115 120 125
Arg Asn Met Val Val Lys Ala Cys Gly Cys His
130 135

SEQ ID NO :7

Met Arg Ala Trp Leu Leu Leu Leu Ala Val Leu Ala Thr Phe Gln Thr
1 5 10 15
Ile Val Arg Val Ala Ser Thr Glu Asp Ile Ser Gln Arg Phe Ile Ala
20 25 30
Ala Ile Ala Pro Val Ala Ala His Ile Pro Leu Ala Ser Ala Ser Gly
35 40 45
Ser Gly Ser Gly Arg Ser Gly Ser Arg Ser Gly Gly Ala Ser Thr Ser
50 55 60
Thr Ala Leu Ala Lys Ala Phe Asn Pro Phe Ser Glu Pro Ala Ser Phe
65 70 75 80
Ser Asp Ser Asp Lys Ser His Arg Ser Lys Thr Asn Lys Lys Pro Ser
85 90 95
Lys Ser Asp Ala Asn Arg Gln Phe Asn Glu Val His Lys Pro Arg Thr
100 105 110
Asp Gln Leu Glu Asn Ser Lys Asn Met Ser Lys Gln Leu Val Asn Lys
115 120 125
Pro Asn His Asn Lys Met Ala Val Lys Glu Gln Arg Ser His His Lys
130 135 140
Lys Ser His His His Arg Ser His Gln Pro Lys Gln Ala Ser Ala Ser

145		150		155		160									
Thr	Glu	Ser	His	Gln	Ser	Ser	Ser	Ile	Glu	Ser	Ile	Phe	Val	Glu	Glu
				165					170					175	
Pro	Thr	Leu	Val	Leu	Asp	Arg	Glu	Val	Ala	Ser	Ile	Asn	Val	Pro	Ala
			180					185					190		
Asn	Ala	Lys	Ala	Ile	Ile	Ala	Glu	Gln	Gly	Pro	Ser	Thr	Tyr	Ser	Lys
		195					200					205			
Glu	Ala	Leu	Ile	Lys	Asp	Lys	Leu	Lys	Pro	Asp	Pro	Ser	Thr	Leu	Val
	210					215					220				
Glu	Ile	Glu	Lys	Ser	Leu	Leu	Ser	Leu	Phe	Asn	Met	Lys	Arg	Pro	Pro
225					230					235					240
Lys	Ile	Asp	Arg	Ser	Lys	Ile	Ile	Ile	Pro	Glu	Pro	Met	Lys	Lys	Leu
			245						250					255	
Tyr	Ala	Glu	Ile	Met	Gly	His	Glu	Leu	Asp	Ser	Val	Asn	Ile	Pro	Lys
			260					265					270		
Pro	Gly	Leu	Leu	Thr	Lys	Ser	Ala	Asn	Thr	Val	Arg	Ser	Phe	Thr	His
		275					280					285			
Lys	Asp	Ser	Lys	Ile	Asp	Asp	Arg	Phe	Pro	His	His	His	Arg	Phe	Arg
	290					295					300				
Leu	His	Phe	Asp	Val	Lys	Ser	Ile	Pro	Ala	Asp	Glu	Lys	Leu	Lys	Ala
305					310					315					320
Ala	Glu	Leu	Gln	Leu	Thr	Arg	Asp	Ala	Leu	Ser	Gln	Gln	Val	Val	Ala
				325					330					335	
Ser	Arg	Ser	Ser	Ala	Asn	Arg	Thr	Arg	Tyr	Gln	Val	Leu	Val	Tyr	Asp
			340					345					350		
Ile	Thr	Arg	Val	Gly	Val	Arg	Gly	Gln	Arg	Glu	Pro	Ser	Tyr	Leu	Leu
		355					360					365			
Leu	Asp	Thr	Lys	Thr	Val	Arg	Leu	Asn	Ser	Thr	Asp	Thr	Val	Ser	Leu
	370					375					380				

Asp Val Gln Pro Ala Val Asp Arg Trp Leu Ala Ser Pro Gln Arg Asn
385 390 395 400

Tyr Gly Leu Leu Val Glu Val Arg Thr Val Arg Ser Leu Lys Pro Ala
405 410 415

Pro His His His Val Arg Leu Arg Arg Ser Ala Asp Glu Ala His Glu
420 425 430

Arg Trp Gln His Lys Gln Pro Leu Leu Phe Thr Tyr Thr Asp Asp Gly
435 440 445

Arg His Lys Ala Arg Ser Ile Arg Asp Val Ser Gly Gly Glu Gly Gly
450 455 460

Gly Lys Gly Gly Arg Asn Lys Arg Gln Pro Arg Arg Pro Thr Arg Arg
465 470 475 480

Lys Asn His Asp Asp Thr Cys Arg Arg His Ser Leu Tyr Val Asp Phe
485 490 495

Ser Asp Val Gly Trp Asp Asp Trp Ile Val Ala Pro Leu Gly Tyr Asp
500 505 510

Ala Tyr Tyr Cys His Gly Lys Cys Pro Phe Pro Leu Ala Asp His Phe
515 520 525

Asn Ser Thr Asn His Ala Val Val Gln Thr Leu Val Asn Asn Met Asn
530 535 540

Pro Gly Lys Val Pro Lys Ala Cys Cys Val Pro Thr Gln Leu Asp Ser
545 550 555 560

Val Ala Met Leu Tyr Leu Asn Asp Gln Ser Thr Val Val Leu Lys Asn
565 570 575

Tyr Gln Glu Met Thr Val Val Gly Cys Gly Cys Arg
580 585

SEQ ID NO :8

Met Val Trp Leu Arg Leu Trp Ala Phe Leu His Ile Leu Ala Ile Val
1 5 10 15

Thr Leu Asp Pro Glu Leu Lys Arg Arg Glu Glu Leu Phe Leu Arg Ser
20 25 30

Leu Gly Phe Ser Ser Lys Pro Asn Pro Val Ser Pro Pro Val Pro
35 40 45

Ser Ile Leu Trp Arg Ile Phe Asn Gln Arg Met Gly Ser Ser Ile Gln
50 55 60

Lys Lys Lys Pro Asp Leu Cys Phe Val Glu Glu Phe Asn Val Pro Gly
65 70 75 80

Ser Val Ile Arg Val Phe Pro Asp Gln Gly Arg Phe Ile Ile Pro Tyr
85 90 95

Ser Asp Asp Ile His Pro Thr Gln Cys Leu Glu Lys Arg Leu Phe Phe
100 105 110

Asn Ile Ser Ala Ile Glu Lys Glu Glu Arg Val Thr Met Gly Ser Gly
115 120 125

Ile Glu Val Gln Pro Glu His Leu Leu Arg Lys Gly Ile Asp Leu Arg
130 135 140

Leu Tyr Arg Thr Leu Gln Ile Thr Leu Lys Gly Met Gly Arg Ser Lys
145 150 155 160

Thr Ser Arg Lys Leu Leu Val Ala Gln Thr Phe Arg Leu Leu His Lys
165 170 175

Ser Leu Phe Phe Asn Leu Thr Glu Ile Cys Gln Ser Trp Gln Asp Pro
180 185 190

Leu Lys Asn Leu Gly Leu Val Leu Glu Ile Phe Pro Lys Lys Glu Ser
195 200 205

Ser Trp Met Ser Thr Ala Asn Asp Glu Cys Lys Asp Ile Gln Thr Phe
210 215 220

Leu Tyr Thr Ser Leu Leu Thr Val Thr Leu Asn Pro Leu Arg Cys Lys
225 230 235 240

Arg Pro Arg Arg Lys Arg Ser Tyr Ser Lys Leu Pro Phe Thr Ala Ser

245	250	255
Asn Ile Cys Lys Lys Arg His Leu Tyr Val Glu Phe Lys Asp Val Gly		
260	265	270
Trp Gln Asn Trp Val Ile Ala Pro Gln Gly Tyr Met Ala Asn Tyr Cys		
275	280	285
Tyr Gly Glu Cys Pro Tyr Pro Leu Thr Glu Ile Leu Asn Gly Ser Asn		
290	295	300
His Ala Ile Leu Gln Thr Leu Val His Ser Ile Glu Pro Glu Asp Ile		
305	310	315
		320
Pro Leu Pro Cys Cys Val Pro Thr Lys Met Ser Pro Ile Ser Met Leu		
325	330	335
Phe Tyr Asp Asn Asn Asp Asn Val Val Leu Arg His Tyr Glu Asn Met		
340	345	350
Ala Val Asp Glu Cys Gly Cys Arg		
355	360	

SEQ ID NO:9

Met Arg Lys Met Gln Lys Glu Ile Leu Ser Val Leu Gly Pro Pro His		
1	5	10
		15
Arg Pro Arg Pro Leu His Gly Leu Gln Gln Pro Gln Pro Pro Val Leu		
20	25	30
Pro Pro Gln Gln Gln Gln Gln Gln Gln Gln Gln Thr Ala Arg Glu		
35	40	45
Glu Pro Pro Pro Gly Arg Leu Lys Ser Ala Pro Leu Phe Met Leu Asp		
50	55	60
Leu Tyr Asn Ala Leu Ser Asn Asp Asp Glu Glu Asp Gly Ala Ser Glu		
65	70	75
		80
Gly Val Gly Gln Glu Pro Gly Ser His Gly Gly Ala Ser Ser Ser Gln		
85	90	95
Leu Arg Gln Pro Ser Pro Gly Ala Ala His Ser Leu Asn Arg Lys Ser		

100	105	110
Leu Leu Ala Pro Gly Pro Gly Gly Gly Ala Ser Pro Leu Thr Ser Ala		
115	120	125
Gln Asp Ser Ala Phe Leu Asn Asp Ala Asp Met Val Met Ser Phe Val		
130	135	140
Asn Leu Val Glu Tyr Asp Lys Glu Phe Ser Pro His Gln Arg His His		
145	150	155
		160
Lys Glu Phe Lys Phe Asn Leu Ser Gln Ile Pro Glu Gly Glu Ala Val		
165	170	175
Thr Ala Ala Glu Phe Arg Val Tyr Lys Asp Cys Val Val Gly Ser Phe		
180	185	190
Lys Asn Gln Thr Phe Leu Ile Ser Ile Tyr Gln Val Leu Gln Glu His		
195	200	205
Gln His Arg Asp Ser Asp Leu Phe Leu Leu Asp Thr Arg Val Val Trp		
210	215	220
Ala Ser Glu Glu Gly Trp Leu Glu Phe Asp Ile Thr Ala Thr Ser Asn		
225	230	235
		240
Leu Trp Val Val Thr Pro Gln His Asn Met Gly Leu Gln Leu Ser Val		
245	250	255
Val Thr Arg Asp Gly Leu His Val Asn Pro Arg Ala Ala Gly Leu Val		
260	265	270
Gly Arg Asp Gly Pro Tyr Asp Lys Gln Pro Phe Met Val Ala Phe Phe		
275	280	285
Lys Val Ser Glu Val His Val Arg Thr Thr Arg Ser Ala Ser Ser Arg		
290	295	300
Arg Arg Gln Gln Ser Arg Asn Arg Ser Thr Gln Ser Gln Asp Val Ser		
305	310	315
		320
Arg Gly Ser Gly Ser Ser Asp Tyr Asn Gly Ser Glu Leu Lys Thr Ala		
325	330	335

Cys Lys Lys His Glu Leu Tyr Val Ser Phe Gln Asp Leu Gly Trp Gln
340 345 350

Asp Trp Ile Ile Ala Pro Lys Gly Tyr Ala Ala Asn Tyr Cys Asp Gly
355 360 365

Glu Cys Ser Phe Pro Leu Asn Ala His Met Asn Ala Thr Asn His Ala
370 375 380

Ile Val Gln Thr Leu Val His Leu Met Asn Pro Glu Tyr Val Pro Lys
385 390 395 400

Pro Cys Cys Ala Pro Thr Lys Leu Asn Ala Ile Ser Val Leu Tyr Phe
405 410 415

Asp Asp Asn Ser Asn Val Ile Leu Lys Lys Tyr Arg Asn Met Val Val
420 425 430

Arg Ala Cys Gly Cys His
435

SEQ ID NO:10

ggggacttct tgaacttgca gggagaataa cttgcgcacc ccactttgcg ccggtgcctt 60
tgccccagcg gagcctgctt cgccatctcc gagccccacc gccctccac tcctcggcct 20
tgcccgacac tgagacgctg ttcccgagct gaaaagagag actgcgcggc cggcaccg 180
gagaaggagg aggcaaagaa aaggaaacgga cattcggtcc ttgcgccagg tcctttgacc 240
agagtttttc catgtggacg ctctttcaat ggacgtgtcc ccgctgctt cttagacgga 300
ctgcggtctc cttaaaggtcg accatggtgg ccgggacccg ctgtcttcta gcgttgctgc 360
ttccccaggt ctcctgggc ggcgcggtg gcctcggtcc ggagctgggc cgcaggaagt 420
tcgcgcgggc gtcgtcgggc cgccctcat ccagccctc tgacgaggtc ctgagcgagt 480
tcgagttgcg gctgctcagc atgttcggcc tgaaacagag acccaccacc agcagggacg 540
ccgtggtgcc cccctacatg ctagacctgt atcgaggca ctcaggtcag ccgggctcac 600
ccgccccaga ccaccggttg gagagggcag ccagccgagc caaactgtg cgcagcttcc 660
accatgaaga atctttggaa gaactaccag aaacgagtgg gaaaacaacc cggagattct 720
tctttaattt aagttctatc cccacggagg agtttatcac ctcagcagag cttcaggttt 780
tccgagaaca gatgcaagat gctttaggaa acaatagcag tttccatcac cgaattaata 840

tttatgaaat cataaaacct gcaacagcca actcgaaatt ccccgtagacc agacttttgg 900
acaccagggtt ggtgaatcag aatgcaagca ggtgggaaag ttttgatgtc acccccgctg 960
tgatgcggtg gactgcacag ggacacgcca accatggatt cgtgggtgaa gtggccact 1020
tggaggagaa acaaggtgtc tccaagagac atgttaggat aagcaggtct ttgcaccaag 1080
atgaacacag ctggtcacag ataaggccat tgctagtaac ttttgccat gatggaaaag 1140
ggcatcctct ccacaaaaga gaaaaacgtc aagccaaaca caaacagcgg aaacgcctta 1200
agtccagctg taagagacac cctttgtacg tggacttcag tgacgtggg tggaatgact 1260
ggattgtggc tccccgggg tatcacgct tttactgcca cggagaatgc ctttttctc 1320
tggctgatca tctgaactcc actaatcatg ccattgttca gacgttggtc aactctgtta 1380
actctaagat tcctaaggca tgctgtgtcc cgacagaact cagtgtctatc tcgatgtgt 1440
accttgacga gaatgaaaag gttgtattaa agaactatca ggacatgggt gtggagggtt 1500
gtgggtgtcg ctagtacagc aaaattaaat acataaatat atatata 1547

SEQ ID NO:11

ggcagaggag gagggaggga gggaaggagc gcggagcccg gcccggaagc taggtgagt 60
tggcatccga gctgaggagc gcgagcctga gacgccgtg ctgctccggc tgagtatcta 20
gcttgtctcc ccgatgggat tcccgccaa gctatctcga gcctgcagcg ccacagtccc 80
cggccctcgc ccaggttcac tgcaaccgtt cagaggtccc caggagctgc tgctggcgag 40
cccgtactg cagggacctg tggagccatt ccgtagtgc atcccgagca acgcactgct 300
gcagcttccc tgagcctttc cagcaagttt gttcaagatt ggctgtcaag aatcatggac 360
tgttattata tgccttgttt tctgtcaaga caccatgatt cctggtaacc gaatgctgat 420
ggtcgtttta ttatgccaag tcctgctagg aggcgcgagc catgctagtt tgatacctga 480
gacggggaag aaaaaagtcg ccgagattca gggccacgcg ggaggacgcc gctcagggca 540
gagccatgag ctctgcggg acttcgaggc gacacttctg cagatgtttg ggctgcgccg 600
ccgcccgcag cctagcaaga gtgccgtcat tccggactac atgcgggatc tttaccggct 660
tcagtctggg gaggaggagg aagagcagat ccacagcact ggtcttgagt atcctgagcg 720
cccggccagc cgggccaaca ccgtgaggag ctccaccac gaagaacatc tggagaacat 780
cccagggacc agtgaaaact ctgcttttcg tttcctcttt aacctcagca gcatccctga 840
gaacgaggtg atctcctctg cagagcttcg gctcttccgg gagcaggtgg accagggcc 900

tgattgggaa aggggcttcc accgtataaa catttatgag gttatgaagc ccccagcaga 960
 agtgggtgcct gggcacctca tcacacgact actggacacg agactgggtcc accacaatgt 1020
 gacacgggtgg gaaacttttg atgtgagccc tgcggtcctt cgctggaccc gggagaagca 1080
 gccaaactat gggctagcca ttgaggtgac tcacctccat cagactcgga cccaccaggg 1140
 ccagcatgtc aggattagcc gatcggtacc tcaagggagt gggaattggg cccagctccg 1200
 gccctcctg gtcacctttg gccatgatgg ccggggccat gccttgaccc gacgccggag 1260
 ggccaagcgt agccctaagc atcactcaca gcggggccagg aagaagaata agaactgccg 1320
 gcgccactcg ctctatgtgg acttcagcga tgtgggctgg aatgactgga ttgtggcccc 1380
 accaggctac caggccttct actgccatgg ggactgcccc tttccactgg ctgaccacct 1440
 caactcaacc aaccatgcca ttgtgcagac cctgggtcaat tctgtcaatt ccagtatccc 1500
 caaagcctgt tgtgtgcca ctgaactgag tgccatctcc atgctgtacc tggatgagta 1560
 tgataagggtg gtactgaaaa attatcagga gatggtagta gagggatgtg ggtgccgctg 1620
 agatcaggca gtccttgagg atagacagat atacacacca cacacacaca ccacatacac 1680
 cacacacaca cgttcccatc cactcaccca cacactacac agactgcttc cttatagctg 1740
 gactttttatt t 1751

SEQ ID NO:12

Met Ala Gly Ala Ser Arg Leu Leu Phe Leu Trp Leu Gly Cys Phe Cys
 1 5 10 15
 Val Ser Leu Ala Gln Gly Glu Arg Pro Lys Pro Pro Phe Pro Glu Leu
 20 25 30
 Arg Lys Ala Val Pro Gly Asp Arg Thr Ala Gly Gly Gly Pro Asp Ser
 35 40 45
 Glu Leu Gln Pro Gln Asp Lys Val Ser Glu His Met Leu Arg Leu Tyr
 50 55 60
 Asp Arg Tyr Ser Thr Val Gln Ala Ala Arg Thr Pro Gly Ser Leu Glu
 65 70 75 80
 Gly Gly Ser Gln Pro Trp Arg Pro Arg Leu Leu Arg Glu Gly Asn Thr
 85 90 95
 Val Arg Ser Phe Arg Ala Ala Ala Ala Glu Thr Leu Glu Arg Lys Gly

100	105	110
Leu Tyr Ile Phe Asn Leu Thr Ser Leu Thr Lys Ser Glu Asn Ile Leu		
115	120	125
Ser Ala Thr Leu Tyr Phe Cys Ile Gly Glu Leu Gly Asn Ile Ser Leu		
130	135	140
Ser Cys Pro Val Ser Gly Gly Cys Ser His His Ala Gln Arg Lys His		
145	150	155
		160
Ile Gln Ile Asp Leu Ser Ala Trp Thr Leu Lys Phe Ser Arg Asn Gln		
165	170	175
Ser Gln Leu Leu Gly His Leu Ser Val Asp Met Ala Lys Ser His Arg		
180	185	190
Asp Ile Met Ser Trp Leu Ser Lys Asp Ile Thr Gln Phe Leu Arg Lys		
195	200	205
Ala Lys Glu Asn Glu Glu Phe Leu Ile Gly Phe Asn Ile Thr Ser Lys		
210	215	220
Gly Arg Gln Leu Pro Lys Arg Arg Leu Pro Phe Pro Glu Pro Tyr Ile		
225	230	235
		240
Leu Val Tyr Ala Asn Asp Ala Ala Ile Ser Glu Pro Glu Ser Val Val		
245	250	255
Ser Ser Leu Gln Gly His Arg Asn Phe Pro Thr Gly Thr Val Pro Lys		
260	265	270
Trp Asp Ser His Ile Arg Ala Ala Leu Ser Ile Glu Arg Arg Lys Lys		
275	280	285
Arg Ser Thr Gly Val Leu Leu Pro Leu Gln Asn Asn Glu Leu Pro Gly		
290	295	300
Ala Glu Tyr Gln Tyr Lys Lys Asp Glu Val Trp Glu Glu Arg Lys Pro		
305	310	315
		320
Tyr Lys Thr Leu Gln Ala Gln Ala Pro Glu Lys Ser Lys Asn Lys Lys		
325	330	335

Lys Gln Arg Lys Gly Pro His Arg Lys Ser Gln Thr Leu Gln Phe Asp
 340 345 350

Glu Gln Thr Leu Lys Lys Ala Arg Arg Lys Gln Trp Ile Glu Pro Arg
 355 360 365

Asn Cys Ala Arg Arg Tyr Leu Lys Val Asp Phe Ala Asp Ile Gly Trp
 370 375 380

Ser Glu Trp Ile Ile Ser Pro Lys Ser Phe Asp Ala Tyr Tyr Cys Ser
 385 390 395 400

Gly Ala Cys Gln Phe Pro Met Pro Lys Ser Leu Lys Pro Ser Asn His
 405 410 415

Ala Thr Ile Gln Ser Ile Val Arg Ala Val Gly Val Val Pro Gly Ile
 420 425 430

Pro Glu Pro Cys Cys Val Pro Glu Lys Met Ser Ser Leu Ser Ile Leu
 435 440 445

Phe Phe Asp Glu Asn Lys Asn Val Val Leu Lys Val Tyr Pro Asn Met
 450 455 460

Thr Val Glu Ser Cys Ala Cys Arg
 465 470

SEQ ID NO:13

Met Pro Pro Pro Gln Gln Gly Pro Cys Gly His His Leu Leu Leu Leu
 1 5 10 15

Leu Ala Leu Leu Leu Pro Ser Leu Pro Leu Thr Arg Ala Pro Val Pro
 20 25 30

Pro Gly Pro Ala Ala Ala Leu Leu Gln Ala Leu Gly Leu Arg Asp Glu
 35 40 45

Pro Gln Gly Ala Pro Arg Leu Arg Pro Val Pro Pro Val Met Trp Arg
 50 55 60

Leu Phe Arg Arg Arg Asp Pro Gln Glu Thr Arg Ser Gly Ser Arg Arg
 65 70 75 80

Thr Ser Pro Gly Val Thr Leu Gln Pro Cys His Val Glu Glu Leu Gly
85 90 95

Val Ala Gly Asn Ile Val Arg His Ile Pro Asp Arg Gly Ala Pro Thr
100 105 110

Arg Ala Ser Glu Pro Val Ser Ala Ala Gly His Cys Pro Glu Trp Thr
115 120 125

Val Val Phe Asp Leu Ser Ala Val Glu Pro Ala Glu Arg Pro Ser Arg
130 135 140

Ala Arg Leu Glu Leu Arg Phe Ala Ala Ala Ala Ala Ala Pro Glu
145 150 155 160

Gly Gly Trp Glu Leu Ser Val Ala Gln Ala Gly Gln Gly Ala Gly Ala
165 170 175

Asp Pro Gly Pro Val Leu Leu Arg Gln Leu Val Pro Ala Leu Gly Pro
180 185 190

Pro Val Arg Ala Glu Leu Leu Gly Ala Ala Trp Ala Arg Asn Ala Ser
195 200 205

Trp Pro Arg Ser Leu Arg Leu Ala Leu Ala Leu Arg Pro Arg Ala Pro
210 215 220

Ala Ala Cys Ala Arg Leu Ala Glu Ala Ser Leu Leu Leu Val Thr Leu
225 230 235 240

Asp Pro Arg Leu Cys His Pro Leu Ala Arg Pro Arg Arg Asp Ala Glu
245 250 255

Pro Val Leu Gly Gly Gly Pro Gly Gly Ala Cys Arg Ala Arg Arg Leu
260 265 270

Tyr Val Ser Phe Arg Glu Val Gly Trp His Arg Trp Val Ile Ala Pro
275 280 285

Arg Gly Phe Leu Ala Asn Tyr Cys Gln Gly Gln Cys Ala Leu Pro Val
290 295 300

Ala Leu Ser Gly Ser Gly Gly Pro Pro Ala Leu Asn His Ala Val Leu

305 310 315 320

Arg Ala Leu Met His Ala Ala Ala Pro Gly Ala Ala Asp Leu Pro Cys
325 330 335

Cys Val Pro Ala Arg Leu Ser Pro Ile Ser Val Leu Phe Phe Asp Asn
340 345 350

Ser Asp Asn Val Val Leu Arg Gln Tyr Glu Asp Met Val Val Asp Glu
355 360 365

Cys Gly Cys Arg
370

SEQ ID NO:14

Met Ser Gly Leu Arg Asn Thr Ser Glu Ala Val Ala Val Leu Ala Ser
1 5 10 15

Leu Gly Leu Gly Met Val Leu Leu Met Phe Val Ala Thr Thr Pro Pro
20 25 30

Ala Val Glu Ala Thr Gln Ser Gly Ile Tyr Ile Asp Asn Gly Lys Asp
35 40 45

Gln Thr Ile Met His Arg Val Leu Ser Glu Asp Asp Lys Leu Asp Val
50 55 60

Ser Tyr Glu Ile Leu Glu Phe Leu Gly Ile Ala Glu Arg Pro Thr His
65 70 75 80

Leu Ser Ser His Gln Leu Ser Leu Arg Lys Ser Ala Pro Lys Phe Leu
85 90 95

Leu Asp Val Tyr His Arg Ile Thr Ala Glu Glu Gly Leu Ser Asp Gln
100 105 110

Asp Glu Asp Asp Asp Tyr Glu Arg Gly His Arg Ser Arg Arg Ser Ala
115 120 125

Asp Leu Glu Glu Asp Glu Gly Glu Gln Gln Lys Asn Phe Ile Thr Asp
130 135 140

Leu Asp Lys Arg Ala Ile Asp Glu Ser Asp Ile Ile Met Thr Phe Leu

145		150		155		160
Asn Lys Arg His	His Asn Val Asp Glu Leu Arg His Glu His Gly Arg	165		170		175
Arg Leu Trp Phe Asp Val Ser Asn Val Pro Asn Asp Asn Tyr Leu Val		180		185		190
Met Ala Glu Leu Arg Ile Tyr Gln Asn Ala Asn Glu Gly Lys Trp Leu		195		200		205
Thr Ala Asn Arg Glu Phe Thr Ile Thr Val Tyr Ala Ile Gly Thr Gly		210		215		220
Thr Leu Gly Gln His Thr Met Glu Pro Leu Ser Ser Val Asn Thr Thr		225		230		235
Gly Asp Tyr Val Gly Trp Leu Glu Leu Asn Val Thr Glu Gly Leu His		245		250		255
Glu Trp Leu Val Lys Ser Lys Asp Asn His Gly Ile Tyr Ile Gly Ala		260		265		270
His Ala Val Asn Arg Pro Asp Arg Glu Val Lys Leu Asp Asp Ile Gly		275		280		285
Leu Ile His Arg Lys Val Asp Asp Glu Phe Gln Pro Phe Met Ile Gly		290		295		300
Phe Phe Arg Gly Pro Glu Leu Ile Lys Ala Thr Ala His Ser Ser His		305		310		315
His Arg Ser Lys Arg Ser Ala Ser His Pro Arg Lys Arg Lys Lys Ser		325		330		335
Val Ser Pro Asn Asn Val Pro Leu Leu Glu Pro Met Glu Ser Thr Arg		340		345		350
Ser Cys Gln Met Gln Thr Leu Tyr Ile Asp Phe Lys Asp Leu Gly Trp		355		360		365
His Asp Trp Ile Ile Ala Pro Glu Gly Tyr Gly Ala Phe Tyr Cys Ser		370		375		380

Gly Glu Cys Asn Phe Pro Leu Asn Ala His Met Asn Ala Thr Asn His
385 390 395 400

Ala Ile Val Gln Thr Leu Val His Leu Leu Glu Pro Lys Lys Val Pro
405 410 415

Lys Pro Cys Cys Ala Pro Thr Arg Leu Gly Ala Leu Pro Val Leu Tyr
420 425 430

His Leu Asn Asp Glu Asn Val Asn Leu Lys Lys Tyr Arg Asn Met Ile
435 440 445

Val Lys Ser Cys Gly Cys His
450 455

SEQ ID NO:15

1 mhltvflkkg ivgflwscwv lvgyakgglg dnhvhssfiy rrlrnherre igreilsilg
61 lphrprpfsp gkqassaplf mldlynamtn eenpeeseys vraslaetr garkgypasp
121 ngyprrigls rtthlttqsp plaslhdtmf lndadmvmf vnlverdkdf shqrrhykef
181 rfdltqiphg eavtaafri ykdrsnrfe netikisiyq iiketytnrda dlflldtrka
241 qaldvgwlvf ditvtsnhwv inpqnnlglq lcaetgdgrs invksaglv rggpgskqpf
301 mvaffkasev llrsvraank rknqnrnkss shqdssrmss vgdyntseqk qackkhelyv
361 sfrdlgwqdw iiapegyaaf ycdgecsfpl nahmnatnha ivqtlvhlmf pdhvpkpcca
421 ptklnaisvl yfddssnvil kkyrnmvrs cgch

SEQ ID NO:16

Met Pro Gly Leu Gly Arg Arg Ala Gln Trp Leu Cys Trp Trp Trp Gly
1 5 10 15

Leu Leu Cys Ser Cys Cys Gly Pro Pro Pro Leu Arg Pro Pro Leu Pro
20 25 30

Ala Ala Ala Ala Ala Ala Ala Gly Gly Gln Leu Leu Gly Asp Gly Gly
35 40 45

Ser Pro Gly Arg Thr Glu Gln Pro Pro Pro Ser Pro Gln Ser Ser Ser
50 55 60

Gly Phe Leu Tyr Arg Arg Leu Lys Thr Gln Glu Lys Arg Glu Met Gln
65 70 75 80

Lys Glu Ile Leu Ser Val Leu Gly Leu Pro His Arg Pro Arg Pro Leu
85 90 95

His Gly Leu Gln Gln Pro Gln Pro Pro Ala Leu Arg Gln Gln Glu Glu
100 105 110

Gln Gln Gln Gln Gln Gln Leu Pro Arg Gly Glu Pro Pro Pro Gly Arg
115 120 125

Leu Lys Ser Ala Pro Leu Phe Met Leu Asp Leu Tyr Asn Ala Leu Ser
130 135 140

Ala Asp Asn Asp Glu Asp Gly Ala Ser Glu Gly Glu Arg Gln Gln Ser
145 150 155 160

Trp Pro His Glu Ala Ala Ser Ser Ser Gln Arg Arg Gln Pro Pro Pro
165 170 175

Gly Ala Ala His Pro Leu Asn Arg Lys Ser Leu Leu Ala Pro Gly Ser
180 185 190

Gly Ser Gly Gly Ala Ser Pro Leu Thr Ser Ala Gln Asp Ser Ala Phe
195 200 205

Leu Asn Asp Ala Asp Met Val Met Ser Phe Val Asn Leu Val Glu Tyr
210 215 220

Asp Lys Glu Phe Ser Pro Arg Gln Arg His His Lys Glu Phe Lys Phe
225 230 235 240

Asn Leu Ser Gln Ile Pro Glu Gly Glu Val Val Thr Ala Ala Glu Phe
245 250 255

Arg Ile Tyr Lys Asp Cys Val Met Gly Ser Phe Lys Asn Gln Thr Phe
260 265 270

Leu Ile Ser Ile Tyr Gln Val Leu Gln Glu His Gln His Arg Asp Ser
275 280 285

Asp Leu Phe Leu Leu Asp Thr Arg Val Val Trp Ala Ser Glu Glu Gly
290 295 300

Trp Leu Glu Phe Asp Ile Thr Ala Thr Ser Asn Leu Trp Val Val Thr
305 310 315 320

Pro Gln His Asn Met Gly Leu Gln Leu Ser Val Val Thr Arg Asp Gly

	325		330		335
Val His Val His Pro Arg Ala Ala Gly Leu Val Gly Arg Asp Gly Pro					
	340		345		350
Tyr Asp Lys Gln Pro Phe Met Val Ala Phe Phe Lys Val Ser Glu Val					
	355		360		365
His Val Arg Thr Thr Arg Ser Ala Ser Ser Arg Arg Arg Gln Gln Ser					
	370		375		380
Arg Asn Arg Ser Thr Gln Ser Gln Asp Val Ala Arg Val Ser Ser Ala					
	385		390		400
Ser Asp Tyr Asn Ser Ser Glu Leu Lys Thr Ala Cys Arg Lys His Glu					
	405		410		415
Leu Tyr Val Ser Phe Gln Asp Leu Gly Trp Gln Asp Trp Ile Ile Ala					
	420		425		430
Pro Lys Gly Tyr Ala Ala Asn Tyr Cys Asp Gly Glu Cys Ser Phe Pro					
	435		440		445
Leu Asn Ala His Met Asn Ala Thr Asn His Ala Ile Val Gln Thr Leu					
	450		455		460
Val His Leu Met Asn Pro Glu Tyr Val Pro Lys Pro Cys Cys Ala Pro					
	465		470		475
Thr Lys Leu Asn Ala Ile Ser Val Leu Tyr Phe Asp Asp Asn Ser Asn					
	485		490		495
Val Ile Leu Lys Lys Tyr Arg Asn Met Val Val Arg Ala Cys Gly Cys					
	500		505		510
His					

SEQ ID NO:17

GGTGCGGGCC CGGAGCCCGG AGCCCGGGTA GCGCGTAGAG CCGGCGCG ATG CAC GTG	57
Met His Val	
1	
CGC TCA CTG CGA GCT GCG GCG CCG CAC AGC TTC GTG GCG CTC TGG GCA	105
Arg Ser Leu Arg Ala Ala Ala Pro His Ser Phe Val Ala Leu Trp Ala	

5	10	15	
CCC CTG TTC CTG CTG CGC TCC GCC CTG GCC GAC TTC AGC CTG GAC AAC			153
Pro Leu Phe Leu Leu Arg Ser Ala Leu Ala Asp Phe Ser Leu Asp Asn			
20	25	30	35
GAG GTG CAC TCG AGC TTC ATC CAC CGG CGC CTC CGC AGC CAG GAG CGG			201
Glu Val His Ser Ser Phe Ile His Arg Arg Leu Arg Ser Gln Glu Arg			
	40	45	50
CGG GAG ATG CAG CGC GAG ATC CTC TCC ATT TTG GGC TTG CCC CAC CGC			249
Arg Glu Met Gln Arg Glu Ile lieu Ser Ile Leu Gly Leu Pro His Arg			
	55	60	65
CCG CGC CCG CAC CTC CAG GGC AAG CAC AAC TCG GCA CCC ATG TTC ATG			297
Pro Arg Pro His Leu Gln Gly Lys His Asn Ser Ala Pro Met Phe Met			
	70	75	80
CTG GAC CTG TAC AAC GCC ATG GCG GTG GAG GAG GGC GGC GGG CCC GGC			345
Leu Asp Leu Tyr Asn Ala Met Ala Val Glu Glu Gly Gly Gly Pro Gly			
	85	90	95
GGC CAG GGC TTC TCC TAC CCC TAC AAG GCC GTC TTC AGT ACC CAG GGC			393
Gly Gln Gly Phe Ser Tyr Pro Tyr Lys Ala Val Phe Ser Thr Gln Gly			
100	105	110	115
CCC CCT CTG GCC AGC CTG CAA GAT AGC CAT TTC CTC ACC GAC GCC GAC			441
Pro Pro Leu Ala Ser Leu Gln Asp Ser His Phe Leu Thr Asp Ala Asp			
	120	125	130
ATG GTC ATG AGC TTC GTC AAC CTC GTG GAA CAT GAC AAG GAA TTC TTC			489
Met Val Met Ser Phe Val Asn Leu Val Glu His Asp Lys Glu Phe Phe			
	135	140	145
CAC CCA CGC TAC CAC CAT CGA GAG TTC CGG TTT GAT CTT TCC AAG ATC			537
His Pro Arg Tyr His His Arg Glu Phe Arg Phe Asp Leu Ser Lys Ile			
	150	155	160
CCA GAA GGG GAA GCT GTC ACG (;CA GCC GAA TTC CGG ATC TAC AAG GAC			
Pro Glu Gly Glu Ala Val Thr Ala Ala Glu Phe Arg Ile Tyr Lys Asp			
	165	170	175
TAC ATC CGG GAA CGC TTC GAC AAT GAG ACG TTC CGG ATC AGC GTT TAT			633
Tyr Ile Arg Glu Arg Phe Asp Asn Glu Thr Phe Arg Ile Ser Val Tyr			
180	185	190	195
CAG GTG CTC CAG GAG CAC TTG GGC AGG GAA TCG GAT CTC TTC CTG CTC			681
Gln Val Leu Gln Glu His Leu Gly Arg Glu Ser Asp Leu Phe Leu Leu			
	200	205	210
GAC AGC CGT ACC CTC TGG GCC TCG GAG GAG GGC TGG CTG GTG TTT GAC			729
Asp Ser Arg Thr Leu Trp Ala Ser Glu Glu Gly Trp Leu Val Phe Asp			
	215	220	225

ATC ACA GCC ACC AGC AAC CAC TGG GTG GTC AAT CCG CGG CAC AAC CTG Ile Thr Ala Thr Ser Asn His Trp Val Val Asn Pro Arg His Asn Leu 230 235 240	777
GGC CTG CAG CTC TCG GTG GAG ACG CTG GAT GGG CAG AGC ATC AAC CCC Gly Leu Gln Leu Ser Val Glu Thr Leu Asp Gly Gln Ser Ile Asn Pro 245 250 255	825
AAG TTG GCG GGC CTG ATT GGG CGG CAC GGG CCC CAG AAC AAG CAG CCC Lys Leu Ala Gly Leu Ile Gly Arg His Gly Pro Gln Asn Lys Gln Pro 260 265 270	873
TTC ATG GTG GCT TTC TTC AAG GCC ACG GAG GTC CAC TTC CGC AGC ATC Phe Met Val Ala Phe Phe Lys Ala Thr Glu Val His Phe Arg Ser Ile 280 285 290	921
CGG TCC ACG GGG AGC AAA CAG CGC AGC CAG AAC CGC TCC AAG ACG CCC Arg Ser Thr Gly Ser Lys Gln Arg Ser Gln Asn Arg Ser Lys Thr Pro 295 300 305	969
AAG AAC CAG GAA GCC CTG CGG ATG GCC AAC GTG GCA GAG AAC AGC AGC Lys Asn Gln Glu Ala Leu Arg Met Ala Asn Val Ala Glu Asn Ser Ser 310 315 320	1017
AGC GAC CAG AGG CAG GCC TGT AAG AAG CAC GAG CTG TAT GTC AGC TTC Ser Asp Gln Arg Gln Ala Cys Lys Lys His Glu Leu Tyr Val Ser Phe 325 330 335	1065
CGA GAC CTG GGC TGG CAG GAC TGG ATC ATC GCG CCT GAA GGC TAC GCC Arg Asp Leu Gly Trp Gln Asp Trp Ile Ile Ala Pro Glu Gly Tyr Ala 340 345 350 355	1113
GCC TAC TAC TGT GAG GGG GAG TGT GCC TTC CCT CTG AAC TCC TAC ATG Ala Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asn Ser Tyr Met 360 365 370	1161
AAC GCC ACC AAC CAC GCC ATC GTG CAG ACG CTG GTC CAC TTC ATC AAC Asn Ala Thr Asn His Ala Ile Val Gln Thr Leu Val His Phe Ile Asn 375 380 385	1209
CCG GAA ACG GTG CCC AAG CCC TGC TGT GCG CCC ACG CAG CTC AAT GCC Pro Glu Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln Leu Asn Ala 390 395 400	1257
ATC TCC GTC CTC TAC TTC GAT GAC AGC TCC AAC GTC ATC CTG AAG AAA Ile Ser Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Ile Leu Lys Lys 405 410 415	1305
TAC AGA AAC ATG GTG GTC CGG GCC TGT GGC TGC CAC TAGCTCCTCC Tyr Arg Asn Met Val Val Arg Ala Cys Gly Cys His 420 425 430	1351
GAGAATTCAG ACCCTTTGGG GCCAAGTTTT TCTGGATCCT CCATTGCTCG CCTTGCCAG	1411
GAACCAGCAG ACCAACTGCC TTTTGTGAGA CCTTCCCCTC CCTATCCCCA ACTTTAAAGG	1471

CTGCAGCAAG	TGACCTCGGG	TCGTGGACCG	CTGCCCTGCC	CCCTCCGCTG	CCACCTGGGG	60
CGGCGCGGGC	CCGGTGCCCC	GGATCGCGCG	TAGAGCCGGC	GCG	ATG CAC GTG CGC Met His Val Arg 1	115
TCG CTG CGC GCT GCG GCG CCA CAC AGC TTC GTG GCG CTC TGG GCG CCT Ser Leu Arg Ala Ala Ala Pro His Ser Phe Val Ala Leu Trp Ala Pro 5 10 15 20	163					
CTG TTC TTG CTG CGC TCC GCC CTG GCC GAT TTC AGC CTG GAC AAC GAG Leu Phe Leu Leu Arg Ser Ala Leu Ala Asp Phe Ser Leu Asp Asn Glu 25 30 35	211					
GTG CAC TCC AGC TTC ATC CAC CGG CGC CTC CGC AGC CAG GAG CGG CGG Val His Ser Ser Phe Ile His Arg Arg Leu Arg Ser Gln Glu Arg Arg 40 45 50	259					
GAG ATG CAG CGG GAG ATC CTG TCC ATC TTA GGG TTG CCC CAT CGC CCG Glu Met Gln Arg Glu Ile Leu Ser Ile Leu Gly Leu Pro His Arg Pro 55 60 65	307					
CGC CCG CAC CTC CAG GGA AAG CAT AAT TCG GCG CCC ATG TTC ATG TTG Arg Pro His Leu Gln Gly Lys His Asn Ser Ala Pro Met Phe Met Leu 70 75 80	355					
GAC CTG TAC AAC GCC ATG GCG GTG GAG GAG AGC GGG CCG GAC GGA CAG Asp Leu Tyr Asn Ala Met Ala Val Glu Glu Ser Gly Pro Asp Gly Gln 85 90 95 100	403					
GGC TTC TCC TAC CCC TAC AAG GCC GTC TTC AGT ACC CAG GGC CCC CCT Gly Phe Ser Tyr Pro Tyr Lys Ala Val Phe Ser Thr Gln Gly Pro Pro 105 110 115	451					
TTA GCC AGC CTG CAG GAC AGC CAT TTC CTC ACT GAC GCC GAC ATG GTC Leu Ala Ser Leu Gln Asp Ser His Phe Leu Thr Asp Ala Asp Met Val 120 125 130	499					
ATG AGC TTC GTC AAC CTA GTG GAA CAT GAC AAA GAA TTC CAC CCT Met Ser Phe Val Asn Leu Val Glu His Asp Lys Glu Phe His Pro 135 140 145	547					

CGA	TAC	CAC	CAT	CGG	GAG	TTC	CGG	TTT	GAT	CTT	TCC	AAG	ATC	CCC	GAG	595
Arg	Tyr	His	His	Arg	Glu	Phe	Arg	Phe	Asp	Leu	Ser	Lys	Ile	Pro	Glu	
150						155					160					
GGC	GAA	CGG	GTG	ACC	GCA	GCC	GAA	TTC	AGG	ATC	TAT	AAG	GAC	TAC	ATC	643
Gly	Glu	Arg	Val	Thr	Ala	Ala	Glu	Phe	Arg	Ile	Tyr	Lys	Asp	Tyr	Ile	
165					170					175					180	
CGG	GAG	CGA	TTT	GAC	AAC	GAG	ACC	TTC	CAG	ATC	ACA	GTC	TAT	CAG	GTG	691
Arg	Glu	Arg	Phe	Asp	Asn	Glu	Thr	Phe	Gln	Ile	Thr	Val	Tyr	Gln	Val	
				185					190					195		
CTC	CAG	GAG	CAC	TCA	GGC	AGG	GAG	TCG	GAC	CTC	TTC	TTG	CTG	GAC	AGC	739
Leu	Gln	Glu	His	Ser	Gly	Arg	Glu	Ser	Asp	Leu	Phe	Leu	Leu	Asp	Ser	
			200					205				210				
CGC	ACC	ATC	TGG	GCT	TCT	GAG	GAG	GGC	TGG	TTG	GTG	TTT	GAT	ATC	ACA	787
Arg	Thr	Ile	Trp	Ala	Ser	Glu	Glu	Gly	Trp	Leu	Val	Phe	Asp	Ile	Thr	
	215					220					225					
GCC	ACC	AGC	AAC	CAC	TGG	GTG	GTC	AAC	CCT	CGG	CAC	AAC	CTG	GGC	TTA	835
Ala	Thr	Ser	Asn	His	Trp	Val	Val	Asn	Pro	Arg	His	Asn	Leu	Gly	Leu	
230						235					240					
CAG	CTC	TCT	GTG	GAG	ACC	CTG	GAT	GGG	CAG	AGC	ATC	AAC	CCC	AAG	TTG	883
Gln	Leu	Ser	Val	Glu	Thr	Leu	Asp	Gly	Gln	Ser	Ile	Asn	Pro	Lys	Leu	
245				250						255					260	
GCA	GGC	CTG	ATT	GGA	CGG	CAT	GGA	CCC	CAG	AAC	AAG	CAA	CCC	TTC	ATG	931
Ala	Gly	Leu	Ile	Gly	Arg	His	Gly	Pro	Gln	Asn	Lys	Gln	Pro	Phe	Met	
				265				270						275		
GTG	GCC	TTC	TTC	AAG	GCC	ACG	GAA	GTC	CAT	CTC	CGT	AGT	ATC	CGG	TCC	979
Val	Ala	Phe	Phe	Lys	Ala	Thr	Glu	Val	His	Leu	Arg	Ser	Ile	Arg	Ser	
			280					285					290			
ACG	GGG	GGC	AAG	CAG	CGC	AGC	CAG	AAT	CGC	TCC	AAG	ACG	CCA	AAG	AAC	1027
Thr	Gly	Gly	Lys	Gln	Arg	Ser	Gln	Asn	Arg	Ser	Lys	Thr	Pro	Lys	Asn	
	295					300					305					
CAA	GAG	GCC	CTG	AGG	ATG	GCC	AGT	GTG	GCA	GAA	AAC	AGC	AGC	AGT	GAC	1075
Gln	Glu	Ala	Leu	Arg	Met	Ala	Ser	Val	Ala	Glu	Asn	Ser	Ser	Ser	Asp	
310						315					320					
CAG	AGG	CAG	GCC	TGC	AAG	AAA	CAT	GAG	CTG	TAC	GTC	AGC	TTC	CGA	GAC	1123
Gln	Arg	Gln	Ala	Cys	Lys	Lys	His	Glu	Leu	Tyr	Val	Ser	Phe	Arg	Asp	
325				330					335						340	
CTT	GGC	TGG	CAG	GAC	TGG	ATC	ATT	GCA	CCT	GAA	GGC	TAT	GCT	GCC	TAC	1171
Leu	Gly	Trp	Gln	Asp	Trp	Ile	Ile	Ala	Pro	Glu	Gly	Tyr	Ala	Ala	Tyr	
			345						350					355		
TAC	TGT	GAG	GGA	GAG	TGC	GCC	TTC	CCT	CTG	AAC	TCC	TAC	ATG	AAC	GCC	1219
Tyr	Cys	Glu	Gly	Glu	Cys	Ala	Phe	Pro	Leu	Asn	Ser	Tyr	Met	Asn	Ala	
			360					365					370			
ACC	AAC	CAC	GCC	ATC	GTC	CAG	ACA	CTG	GTT	CAC	TTC	ATC	AAC	CCA	GAC	1267
Thr	Asn	His	Ala	Ile	Val	Gln	Thr	Leu	Val	His	Phe	Ile	Asn	Pro	Asp	

375	360	385	
ACA GTA CCC AAG CCC TGC TGT GCG CCC ACC CAG CTC AAC GCC ATC TCT			1315
Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln Leu Asn Ala Ile Ser			
390	395	400	
GTC CTC TAC TTC GAC GAC AGC TCT AAT GTC ATC CTG AAG AAG TAC AGA			1363
Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Ile Leu Lys Lys Tyr Arg			
405	410	415	420
AAC ATG GTG GTC CGG GCC TGT GGC TGC CAC TAGCTCTTCC TGAGACCCTG			1413
Asn Met Val Val Arg Ala Cys Gly Cys His			
425	430		
ACCTTTGCGG GGCCACACCT TTCCAAATCT TCGATGTCTC ACCATCTAAG TCTCTCACTG			1473
CCCACCTTGG CGAGGAGAAC AGACCAACCT CTCCTGAGCC TTCCCTCACC TCCCAACCGG			1533
AAGCATGTAA GGGTTCCAGA AACCTGAGCG TGCAGCAGCT GATGAGCGCC CTTTCCTTCT			1593
GGCACGTGAC GGACAAGATC CTACCAGCTA CCACAGCAAA CGCCTAAGAG CAGGAAAAAT			1653
GTCTGCCAGG AAAGTGTCCA GTGTCCACAT GGCCCCTGGC GCTCTGAGTC TTTGAGGAGT			1713
AATCGCAAGC CTCGTTTCAGC TGCAGCAGAA GGAAGGGCTT AGCCAGGGTG GGCGCTGGCG			1773
TCTGTGTTGA AGGGAAACCA AGCAGAAGCC ACTGTAATGA TATGTCACAA TAAAACCCAT			1833
GAATGAAAAA AAAAAAAAAA AAAAAAAAAA AAAAGAATTC			1873

SEQ ID NO:19

Met His Val Arg Ser Leu Arg Ala Ala Ala Pro His Ser Phe Val Ala	
1	15
Leu Trp Ala Pro Leu Phe Leu Leu Arg Ser Ala Leu Ala Asp Phe Ser	
20	30
Leu Asp Asn Glu Val His Ser Ser Phe Ile His Arg Arg Leu Arg Ser	
35	45
Gln Glu Arg Arg Glu Met Gln Arg Glu Ile Leu Ser Ile Leu Gly Leu	
50	60
Pro His Arg Pro Arg Pro His Leu Gln Gly Lys His Asn Ser Ala Pro	
65	80
Met Phe Met Leu Asp Leu Tyr Asn Ala Met Ala Val Glu Glu Ser Gly	
85	95
Pro Asp Gly Gln Gly Phe Ser Tyr Pro Tyr Lys Ala Val Phe Ser Thr	
100	110
Gln Gly Pro Pro Leu Ala Ser Leu Gln Asp Ser His Phe Leu Thr Asp	
115	125

Ala	Asp	Met	Val	Met	Ser	Phe	Val	Asn	Leu	Val	Glu	His	Asp	Lys	Glu	130	135	140
Phe	Phe	His	Pro	Arg	Tyr	His	His	Arg	Glu	Phe	Arg	Phe	Asp	Leu	Ser	145	150	155
Lys	Ile	Pro	Glu	Gly	Glu	Arg	Val	Thr	Ala	Ala	Glu	Phe	Arg	Ile	Tyr	165	170	175
Lys	Asp	Tyr	Ile	Arg	Glu	Arg	Phe	Asp	Asn	Glu	Thr	Phe	Gln	Ile	Thr	180	185	190
Val	Tyr	Gln	Val	Lau	Gln	Glu	His	Ser	Gly	Arg	Glu	Ser	Asp	Leu	Phe	195	200	205
Leu	Leu	Asp	Ser	Arg	Thr	Ile	Trp	Ala	Ser	Glu	Glu	Gly	Trp	Leu	Val	210	215	220
Phe	Asp	Ile	Thr	Ala	Thr	Ser	Asn	His	Trp	Val	Val	Asn	Pro	Arg	His	225	230	235
Asn	Leu	Gly	Lau	Gln	Leu	Ser	Val	Glu	Thr	Leu	Asp	Gly	Gln	Ser	Ile	245	250	255
Asn	Pro	Lys	Leu	Ala	Gly	Leu	Ile	Gly	Arg	His	Gly	Pro	Gln	Asn	Lys	260	265	270
Gln	Pro	Phe	Met	Val	Ala	Phe	Phe	Lys	Ala	Thr	Glu	Val	His	Leu	Arg	275	280	285
Ser	Ile	Arg	Ser	Thr	Gly	Gly	Lys	Gln	Arg	Ser	Gln	Asn	Arg	Ser	Lys	290	295	300
Thr	Pro	Lys	Asn	Gln	Glu	Ala	Leu	Arg	Met	Ala	Ser	Val	Ala	Glu	Asn	305	310	315
Ser	Ser	Ser	Asp	Gln	Arg	Gln	Ala	Cys	Lys	Lys	His	Glu	Leu	Tyr	Val	325	330	335
Ser	Phe	Arg	Asp	Leu	Gly	Trp	Gln	Asp	Trp	Ile	Ile	Ala	Pro	Glu	Gly	340	345	350
Tyr	Ala	Ala	Tyr	Tyr	Cys	Glu	Gly	Glu	Cys	Ala	Phe	Pro	Leu	Asn	Ser	355	360	365
Tyr	Met	Asn	Ala	Thr	Asn	His	Ala	Ile	Val	Gln	Thr	Leu	Val	His	Phe	370	375	380
Ile	Asn	Pro	Asp	Thr	Val	Pro	Lys	Pro	Cys	Cys	Ala	Pro	Thr	Gln	Leu	385	390	395
Asn	Ala	Ile	Ser	Val	Leu	Tyr	Phe	Asp	Asp	Ser	Ser	Asn	Val	Ile	Leu	405	410	415
Lys	Lys	Tyr	Arg	Asn	Met	Val	Val	Arg	Ala	Cys	Gly	Cys	His			420	425	430

SEQ ID NO:20

GGCGCCGCA GAGCAGGAGT GGCTGGAGGA GCTGTGGTTG GAGCAGGAGG TGGCACGGCA	60
GGGCTGGAGG GCTCCCTATG AGTGGCGGAG ACGGCCCAGG AGGCGCTGGA GCAACAGCTC	120
CCACACCGCA CCAAGCGGTG GCTGCAGGAG CTCGCCCATC GCCCCTGCGC TGCTCGGACC	180
GCGGCCACAG CCGGACTGGC GGGTACGGCG GCGACAGAGG CATTGGCCGA GAGTCCCAGT	240
CCGCAGAGTA GCCCCGGCCT CGAGGCGGTG GCGTCCCGGT CCTCTCCGTC CAGGAGCCAG	300
GACAGGTGTC GCGCGGCGGG GCTCCAGGGA CCGCGCCTGA GGCCGGCTGC CCGCCCGTCC	360
CGCCCCGCCC CGCCGCCCGC CGCCCGCCGA GCCCAGCCTC CTTGCCGTCG GGGCGTCCCC	420
AGGCCCTGGG TCGGCCCGG AGCCGATGCG CGCCCGCTGA GCGCCCCAGC TGAGCGCCCC	480
CGGCCTGCC ATG ACC GCG CTC CCC GGC CCG CTC TGG CTC CTG GGC CTG	528
Met Thr Ala Leu Pro Gly Pro Leu Trp Leu Leu Gly Leu	
1 5 10	
GCG CTA TGC GCG CTG GGC GGG GGC GGC CCC GGC CTG CGA CCC CCG CCC	576
Ala Leu Cys Ala Leu Gly Gly Gly Gly Pro Gly Leu Arg Pro Pro Pro	
15 20 25	
GGC TGT CCC CAG CGA CGT CTG GGC GCG CGC GAG CGC CGG GAC GTG CAG	624
Gly Cys Pro Gln Arg Leu Gly Ala Arg Glu Arg Arg Asp Val Gln	
30 35 40 45	
CGC GAG ATC CTG GCG GTG CTC GGG CTG CCT GGG CGG CCC CGG CCC CGC	672
Arg Glu Ile Leu Ala Val Leu Gly Leu Pro Gly Arg Pro Arg Pro Arg	
50 55 60	
GCG CCA CCC GCC GCC TCC CGG CTG CCC GCG TCC GCG CCG CTC TTC ATG	720
Ala Pro Pro Ala Ala Ser Arg Leu Pro Ala Ser Ala Pro Leu Phe Met	
65 70 75	
CTG GAC CTG TAC CAC GCC ATG GCC GGC GAC GAC GAC GAG GAC GGC GCG	768
Leu Asp Leu Tyr His Ala Met Ala Gly Asp Asp Asp Glu Asp Gly Ala	
80 85 90	
CCC GCG GAG CGG CGC CTG GGC CGC GCC GAC CTG GTC ATG AGC TTC GTT	816
Pro Ala Glu Arg Arg Leu Gly Arg Ala Asp Leu Val Met Ser Phe Val	
95 100 105	
AAC ATG GTG GAG CGA GAC CGT GCC CTG GGC CAC CAG GAG CCC CAT TGG	864
Asn Met Val Glu Arg Asp Arg Ala Leu Gly His Gln Glu Pro His Trp	
110 115 120 125	
AAG GAG TTC CGC TTT GAC CTG ACC CAG ATC CCG GCT GGG GAG GCG GTC	912
Lys Glu Phe Arg Phe Asp Leu Thr Gln Ile Pro Ala Gly Glu Ala Val	
130 135 140	
ACA GCT GCG GAG TTC CGG ATT TAC AAG GTG CCC AGC ATC CAC CTG CTC	960
Thr Ala Ala Glu Phe Arg Ile Tyr Lys Val Pro Ser Ile His Leu Leu	
145 150 155	

AAC AGG ACC CTC CAC GTC AGC ATG TIC CAG GTG GTC CAG GAG CAG TCC Asn Arg Thr Leu His Val Ser Met Phe Gln Val Val Gln Glu Gln Ser 160 165 170	1008
AAC AGG GAG TCT GAC TTG TTC TTT TTG GAT CTT CAG ACG CTC CGA GCT Asn Arg Glu Ser Asp Leu Phe Phe Leu Asp Leu Gln Thr Leu Arg Ala 175 180 185	1056
GGA GAC GAG GGC TGG CTG GTG CTG GAT GTC ACA GCA GCC AGT GAC TGC Gly Asp Glu Gly Trp Leu Val Leu Asp Val Thr Ala Ala Ser Asp Cys 190 195 200 205	1104
TGG TTG CTG AAG CGT CAC AAG GAC CTG GGA CTC CGC CTC TAT GTG GAG Trp Leu Leu Lys Arg His Lys Asp Leu Gly Leu Arg Leu Tyr Val Glu 210 215 220	1152
ACT GAG GAC GGG CAC AGC GTG GAT CCT GGC CTG GCC GGC CTG CTG GGT Thr Glu Asp Gly His Ser Val Asp Pro Gly Leu Ala Gly Leu Leu Gly 225 230 235	1200
CAA CGG GCC CCA CGC TCC CAA CAG CCT TTC GTG GTC ACT TTC TTC AGG Gln Arg Ala Pro Arg Ser Gln Gln Pro Phe Val Val Thr Phe Phe Arg 240 245 250	1248
GCC AGT CCG AGT CCC ATC CGC ACC CCT CGG GCA GTG AGG CCA CTG AGG Ala Ser Pro Ser Pro Ile Arg Thr Pro Arg Ala Val Arg Pro Leu Arg 255 260 265	1296
AGG AGG CAG CCG AAG AAA AGC AAC GAG CTG CCG CAG GCC AAC CGA CTC Arg Arg Gln Pro Lys Lys Ser Asn Glu Leu Pro Gln Ala Asn Arg Leu 270 275 280 285	1344
CCA GGG ATC TTT GAT GAC GTC CAC GGC TCC CAC GGC CGG CAG GTC TGC Pro Gly Ile Phe Asp Asp Val His Gly Ser His Gly Arg Gln Val Cys 290 295 300	1392
CGT CGG CAC GAG CTC TAC GTC AGC TTC CAG GAC CTC GGC TGG CTG GAC Arg Arg His Glu Leu Tyr Val Ser Phe Gln Asp Leu Gly Trp Leu Asp 305 310 315	1440
TGG GTC ATC GCT CCC CAA GGC TAC TCG GCC TAT TAC TGT GAG GGG GAG Trp Val Ile Ala Pro Gln Gly Tyr Ser Ala Tyr Tyr Cys Glu Gly Glu 320 325 330	1488
TGC TCC TTC CCA CTG GAC TCC TGC ATG AAT GCC ACC AAC CAC GCC ATC Cys Ser Phe Pro Leu Asp Ser Cys Met Asn Ala Thr Asn His Ala Ile 335 340 345	1536
CTG CAG TCC CTG GTG CAC CTG ATG AAG CCA AAC GCA GTC CCC AAG GCG Leu Gln Ser Leu Val His Leu Met Lys Pro Asn Ala Val Pro Lys Ala 350 355 360 365	1584
TGC TGT GCA CCC ACC AAG CTG AGC GCC ACC TCT GTG CTC TAC TAT GAC Cys Cys Ala Pro Thr Lys Leu Ser Ala Thr Ser Val Leu Tyr Tyr Asp 370 375 380	1632
AGC AGC AAC AAC GTC ATC CTG CGC AAA CAC CGC AAC ATG GTG GTC AAG Ser Ser Asn Asn Val Ile Leu Arg Lys His Arg Asn Met Val Val Lys 385 390 395	1680

GCC TGC GGC TGC CAC T GAGTCAGCCC GCCCAGCCCT ACTGCAG
 Ala Cys Gly Cys His
 400

1723

SEQ ID NO:21

Met Thr Ala Leu Pro Gly Pro Leu Trp Leu Leu Gly Leu Ala Leu Cys
 1 5 10 15
 Ala Leu Gly Gly Gly Gly Pro Gly Leu Arg Pro Pro Pro Gly Cys Pro
 20 25 30
 Gln Arg Arg Leu Gly Ala Arg Glu Arg Arg Asp Val Gln Arg Glu Ile
 35 40 45
 Leu Ala Val Leu Gly Leu Pro Gly Arg Pro Arg Pro Arg Ala Pro Pro
 50 55 60
 Ala Ala Ser Arg Leu Pro Ala Ser Ala Pro Leu Phe Met Leu Asp Leu
 65 70 75 80
 Tyr His Ala Met Ala Gly Asp Asp Asp Glu Asp Gly Ala Pro Ala Glu
 85 90 95
 Arg Arg Leu Gly Arg Ala Asp Leu Val Met Ser Phe Val Asn Met Val
 100 105 110
 Glu Arg Asp Arg Ala Leu Gly His Gln Glu Pro His Trp Lys Glu Phe
 115 120 125
 Arg Phe Asp Leu Thr Gln Ile Pro Ala Gly Glu Ala Val Thr Ala Ala
 130 135 140
 Glu Phe Arg Ile Tyr Lys Val Pro Ser Ile His Leu Leu Asn Arg Thr
 145 150 155 160
 Leu His Val Ser Met Phe Gln Val Val Gln Glu Gln Ser Asn Arg Glu
 165 170 175
 Ser Asp Leu Phe Phe Leu Asp Leu Gln Thr Leu Arg Ala Gly Asp Glu
 180 185 190
 Gly Trp Leu Val Leu Asp Val Thr Ala Ala Ser Asp Cys Trp Leu Leu
 195 200 205
 Lys Arg His Lys Asp Leu Gly Leu Arg Leu Tyr Val Glu Thr Glu Asp
 210 215 220
 Gly His Ser Val Asp Pro Gly Leu Ala Gly Leu Leu Gly Gln Arg Ala
 225 230 235 240
 Pro Arg Ser Gln Gln Pro Phe Val Val Thr Phe Phe Arg Ala Ser Pro
 245 250 255
 Ser Pro Ile Arg Thr Pro Arg Ala Val Arg Pro Leu Arg Arg Arg Gln
 260 265 270

Pro Lys Lys Ser Asn Glu Leu Pro Gln Ala Asn Arg Leu Pro Gly Ile
 275 280 285

Phe Asp Asp Val His Gly Ser His Gly Arg Gln Val Cys Arg Arg His
 290 295 300

Glu Leu Tyr Val Ser Phe Gln Asp Leu Gly Trp Leu Asp Trp Val Ile
 305 310 315 320

Ala Pro Gln Gly Tyr Ser Ala Tyr Tyr Cys Glu Gly Glu Cys Ser Phe
 325 330 335

Pro Leu Asp Ser Cys Met Asn Ala Thr Asn His Ala Ile Leu Gln Ser
 340 345 350

Leu Val His Leu Met Lys Pro Asn Ala Val Pro Lys Ala Cys Cys Ala
 355 360 365

Pro Thr Lys Leu Ser Ala Thr Ser Val Leu Tyr Tyr Asp Ser Ser Asn
 370 375 380

Asn Val Ile Leu Arg Lys His Arg Asn Met Val Val Lys Ala Cys Gly
 385 390 395 400

Cys His

SEQ ID NO:22

GCCAGGCACA GGTGCGCCGT CTGGTCCTCC CCGTCTGGCG TCAGCCGAGC CCGACCAGCT 60

ACCA GTGGAT GCGCGCCGGC TGAAAGTCCG AG ATG GCT ATG CGT CCC GGG CCA 113
 Met Ala Met Arg Pro Gly Pro
 1 5

CTC TGG CTA TTG GGC CTT GCT CTG TGC GCG CTG GGA GGC CAC GGT 161
 Leu Trp Leu Leu Gly Leu Ala Leu Cys Ala Leu Gly Gly His Gly
 10 15 20

CCG CGT CCC CCG CAC ACC TGT CCC CAG CGT CGC CTG GGA GCG CGC GAG 209
 Pro Arg Pro Pro His Thr Cys Pro Gln Arg Arg Leu Gly Ala Arg Glu
 25 30 35

CGC CGC GAC ATG CAG CGT GAA ATC CTG GCG GTG CTC GGG CTA CCG GGA 257
 Arg Arg Asp Met Gln Arg Glu Ile Leu Ala Val Leu Gly Leu Pro Gly
 40 45 50 55

CGG CCC CGA CCC CGT GCA CAA CCC GCC GCT GCC CGG CAG CCA GCG TCC 305
 Arg Pro Arg Pro Arg Ala Gln Pro Ala Ala Arg Gln Pro Ala Ser
 60 65 70

GCG CCC CTC TTC ATG TTG GAC CTA TAC CAC GCC ATG ACC GAT GAC GAC 353
 Ala Pro Leu Phe Met Leu Asp Leu Tyr His Ala Met Thr Asp Asp Asp
 75 80 85

GAC GGC GGG CCA CCA CAG GCT CAC TTA GGC CGT GCC GAC CTG GTC ATG 401
 Asp Gly Gly Pro Pro Gln Ala His Leu Gly Arg Ala Asp Leu Val Met
 90 95 100

AGC TTC GTC AAC ATG GTG GAA CGC GAC CGT ACC CTG GGC TAC CAG GAG Ser Phe Val Asn Met Val Glu Arg Asp Arg Thr Leu Gly Tyr Gln Glu 105 110 115	449
CCA CAC TGG AAG GAA TTC CAC TTT GAC CTA ACC CAG ATC CCT GCT GGG Pro His Trp Lys Glu Phe His Phe Asp Leu Thr Gln Ile Pro Ala Gly 120 125 130 135	497
GAG GCT GTC ACA GCT GCT GAG TTC CGG ATC TAC AAA GAA CCC AGC ACC Glu Ala Val Thr Ala Ala Glu Phe Arg Ile Tyr Lys Glu Pro Ser Thr 140 145 150	545
CAC CCG CTC AAC ACA ACC CTC CAC ATC AGC ATG TTC GAA GTG GTC CAA His Pro Leu Asn Thr Thr Leu His Ile Ser Met Phe Glu Val Val Gln 155 160 165	593
GAG CAC TCC AAC AGG GAG TCT GAC TTG TTC TTT TTG GAT CTT CAG ACG Glu His Ser Asn Arg Glu Ser Asp Leu Phe Phe Leu Asp Leu Gln Thr 170 175 180	641
CTC CGA TCT GGG GAC GAG GGC TGG CTG GTG CTG GAC ATC ACA GCA GCC Leu Arg Ser Gly Asp Glu Gly Trp Leu Val Leu Asp Ile Thr Ala Ala 185 190 195	689
AGT GAC CGA TGG CTG CTG AAC CAT CAC AAG GAC CTG GGA CTC CGC CTC Ser Asp Arg Trp Leu Leu Asn His His Lys Asp Leu Gly Leu Arg Leu 200 205 210 215	737
TAT GTG GAA ACC GCG GAT GGG CAC AGC ATG GAT CCT GGC CTG GCT GGT Tyr Val Glu Thr Ala Asp Gly His Ser Met Asp Pro Gly Leu Ala Gly 220 225 230	785
CTG CTT GGA CGA CAA GCA CCA CGC TCC AGA CAG CCT TTC ATG GTA ACC Leu Leu Gly Arg Gln Ala Pro Arg Ser Arg Gln Pro Phe Met Val Thr 235 240 245	833
TTC TTC AGG GCC AGC CAG AGT CCT GTG CGG GCC CCT CGG GCA GCG AGA Phe Phe Arg Ala Ser Gln Ser Pro Val Arg Ala Pro Arg Ala Ala Arg 250 255 260	881
CCA CTG AAG AGG AGG CAG CCA AAG AAA ACG AAC GAG CTT CCG CAC CCC Pro Leu Lys Arg Arg Gln Pro Lys Lys Thr Asn Glu Leu Pro His Pro 265 270 275	929
AAC AAA CTC CCA GGG ATC TTT GAT GAT GGC CAC GGT TCC CGC GGC AGA Asn Lys Leu Pro Gly Ile Phe Asp Asp Gly His Gly Ser Arg Gly Arg 280 285 290 295	977
GAG GTT TGC CGC AGG CAT GAG CTC TAC GTC AGC TTC CGT GAC CTT GGC Glu Val Cys Arg Arg His Glu Leu Tyr Val Ser Phe Arg Asp Leu Gly 300 305 310	1025
TGG CTG GAC TGG GTC ATC GCC CCC CAG GGC TAC TCT GCC TAT TAC TGT Trp Leu Asp Trp Val Ile Ala Pro Gln Gly Tyr Ser Ala Tyr Tyr Cys 315 320 325	1073
GAG GGG GAG TGT GCT TTC CCA CTG GAC TCC TGT ATG AAC GCC ACC AAC Glu Gly Glu Cys Ala Phe Pro Leu Asp Ser Cys Met Asn Ala Thr Asn	1121

330	335	340	
CAT GCC ATC TTG CAG TCT CTG GTG CAC CTG ATG AAG CCA GAT GTT GTC			1169
His Ala Ile Leu Gln Ser Leu Val His Leu Met Lys Pro Asp Val Val			
345 350		355	
CCC AAG GCA TGC TGT GCA CCC ACC AAA CTG. AGT GCC ACC TCT GTG CTG			
1217			
Pro Lys Ala Cys Cys Ala Pro Thr Lys Leu Ser Ala Thr Ser Val Leu			
360 365	370	375	
TAC TAT GAC AGC AGC AAC AAT GTC ATC CTG CGT AAA CAC CGT AAC ATG			1265
Tyr Tyr Asp Ser Ser Asn Asn Val Ile Leu Arg Lys His Arg Asn Met			
380 385		390	
GTG GTC AAG GCC TGT GGC TGC CAC TGAGGCCCG CCCAGCATCC TGCTTCTACT			1319
Val Val Lys Ala Cys Gly Cys His			
395			
ACCTTACCAT CTGGCCGGGC CCCTCTCCAG AGGCAGAAAC CCTTCTATGT TATCATAGCT			1379
CAGACAGGGG CAATGGGAGG CCCTTCACTT CCCCTGGCCA CTCCTGCTA AAATTCTGGT			1439
CTTTCCCAGT TCCTCTGTCC TTCATGGGGT TTCGGGGCTA TCACCCCGCC CTCTCCATCC			1499
TCCTACCCCA AGCATAGACT GAATGCACAC AGCATCCCAG AGCTATGCTA ACTGAGAGGT			1559
CTGGGGTCAG CACTGAAGGC CCACATGAGG AAGACTGATC CTTGGCCATC CTCAGCCAC			1619
AATGGCAAAT TCTGGATGGT CTAAGAAGGC CCTGGAATTC TAAACTAGAT GATCTGGGCT			1679
CTCTGCACCA TTCATTGTGG CAGTTGGGAC ATTTTtaggt ATAACAGACA CATACTTA			1739
GATCAATGCA TCGCTGTACT CCTTGAAATC AGAGCTAGCT TGTTAGAAAA AGAATCAGAG			1799
CCAGGTATAG CGGTGCATGT CATTAATCCC AGCGCTAAAG AGACAGAGAC AGGAGAATCT			1859
CTGTGAGTTC AAGGCCACAT AGAAAGAGCC TGTCTCGGGA GCAGGAAAAA AAAAAAAC			1919
GGAATTC			1926

SEQ ID NO :23

Met Ala Met Arg Pro Gly Pro Leu Trp Leu Leu Gly Leu Ala Leu Cys	
1 5 10 15	
Ala Leu Gly Gly Gly His Gly Pro Arg Pro Pro His Thr Cys Pro Gln	
20 25 30	
Arg Arg Leu Gly Ala Arg Glu Arg Arg Asp Met Gln Arg Glu Ile Leu	
35 40 45	
Ala Val Leu Gly Leu Pro Gly Arg Pro Arg Pro Arg Ala Gln Pro Ala	
50 55 60	
Ala Ala Arg Gln Pro Ala Ser Ala Pro Leu Phe Met Leu Asp Leu Tyr	
65 70 75 80	

SEQ ID NO:24

generic sequence 1

```
<220>
<221> MISC_FEATURE
<222> (2)..(2)
<223> Xaa at res. 2 = (Tyr or Lys)

<220>
<221> MISC_FEATURE
<222> (3)..(3)
<223> Xaa at res. 3 = Val or Ile)

<220>
<221> MISC_FEATURE
<222> (4)..(4)
<223> Xaa at res. 4 = (Ser, Asp or Glu)

<220>
<221> MISC_FEATURE
<222> (6)..(6)
<223> Xaa at res. 6 = (Arg, Gln, Ser, Lys or Ala)

<220>
<221> MISC_FEATURE
<222> (7)..(7)
<223> Xaa at res. 7 = (Asp or Glu)

<220>
<221> MISC_FEATURE
<222> (8)..(8)
<223> Xaa at res. 8 = (Leu, Val or Ile)

<220>
<221> MISC_FEATURE
<222> (11)..(11)
<223> Xaa at res. 11 = (Gln, Leu, Asp, His, Asn or Ser)

<220>
<221> MISC_FEATURE
<222> (12)..(12)
<223> Xaa at res. 12 = (Asp, Arg, Asn or Glu)

<220>
<221> MISC_FEATURE
<222> (13)..(13)
<223> Xaa at res. 13 = (Trp or Ser)

<220>
<221> MISC_FEATURE
<222> (14)..(14)
<223> Xaa at res. 14 = (Ile or Val)

<220>
<221> MISC_FEATURE
<222> (15)..(15)
```

<223> Xaa at res. 15 = (Ile or Val)

<220>
<221> MISC_FEATURE
<222> (16)..(16)
<223> Xaa at res. 16 (Ala or Ser)

<220>
<221> MISC_FEATURE
<222> (18)..(18)
<223> Xaa at res. 18 = (Glu, Gln, Leu, Lys, Pro or Arg)

<220>
<221> MISC_FEATURE
<222> (19)..(19)
<223> Xaa at res. 19 = (Gly or Ser)

<220>
<221> MISC_FEATURE
<222> (20)..(20)
<223> Xaa at res. 20 = (Tyr or Phe)

<220>
<221> MISC_FEATURE
<222> (21)..(21)
<223> Xaa at res. 21 = (Ala, Ser, Asp, Met, His, Gln, Leu or Gly)

<220>
<221> MISC_FEATURE
<222> (23)..(23)
<223> Xaa at res. 23 = (Tyr, Asn or Phe)

<220>
<221> MISC_FEATURE
<222> (26)..(26)
<223> Xaa at res. 26 = (Glu, His, Tyr, Asp, Gln, Ala or Ser)

<220>
<221> MISC_FEATURE
<222> (28)..(28)
<223> Xaa at res. 28 = (Glu, Lys, Asp, Gln or Ala)

<220>
<221> MISC_FEATURE
<222> (30)..(30)
<223> Xaa at res. 30 = (Ala, Ser, Pro, Gln, Ile or Asn)

<220>
<221> MISC_FEATURE
<222> (31)..(31)
<223> Xaa at res. 31 = (Phe, Leu or Tyr)

<220>
<221> MISC_FEATURE
<222> (33)..(33)
<223> Xaa at res. 33 = (Leu, Val or Met)

<220>

<221> MISC_FEATURE
<222> (34)..(34)
<223> Xaa at res. 34 = (Asn, Asp, Ala, Thr or Pro)

<220>
<221> MISC_FEATURE
<222> (35)..(35)
<223> Xaa at res. 35 = (Ser, Asp, Glu, Leu, Ala or Lys)

<220>
<221> MISC_FEATURE
<222> (36)..(36)
<223> Xaa at res. 36 = (Tyr, Cys, His, Ser or Ile)

<220>
<221> MISC_FEATURE
<222> (37)..(37)
<223> Xaa at res. 37 = (Met, Phe, Gly or Leu)

<220>
<221> MISC_FEATURE
<222> (38)..(38)
<223> Xaa at res. 38 = (Asn, Ser or Lys)

<220>
<221> MISC_FEATURE
<222> (39)..(39)
<223> Xaa at res. 39 = (Ala, Ser, Gly or Pro)

<220>
<221> MISC_FEATURE
<222> (40)..(40)
<223> Xaa at res. 40 = (Thr, Leu or Ser)

<220>
<221> MISC_FEATURE
<222> (44)..(44)
<223> Xaa at res. 44 = (Ile, Val or Thr)

<220>
<221> MISC_FEATURE
<222> (45)..(45)
<223> Xaa at res. 45 = (Val, Leu, Met or Ile)

<220>
<221> MISC_FEATURE
<222> (46)..(46)
<223> Xaa at res. 46 = (Gln or Arg)

<220>
<221> MISC_FEATURE
<222> (47)..(47)
<223> Xaa at res. 47 = (Thr, Ala or Ser)

<220>
<221> MISC_FEATURE
<222> (48)..(48)
<223> Xaa at res. 48 = (Leu or Ile)

```

<220>
<221> MISC_FEATURE
<222> (49)..(49)
<223> Xaa at res. 49 = (Val or Met)

<220>
<221> MISC_FEATURE
<222> (50)..(50)
<223> Xaa at res. 50 = (His, Asn or Arg)

<220>
<221> MISC_FEATURE
<222> (51)..(51)
<223> Xaa at res. 51 = (Phe, Leu, Asn, Ser, Ala or Val)

<220>
<221> MISC_FEATURE
<222> (52)..(52)
<223> Xaa at res. 52 = (Ile, Met, Asn, Ala, Val, Gly or Leu)

<220>
<221> MISC_FEATURE
<222> (53)..(53)
<223> Xaa at res. 53 = (Asn, Lys, Ala, Glu, Gly or Phe)

<220>
<221> MISC_FEATURE
<222> (54)..(54)
<223> Xaa at res. 54 = (Pro, Ser or Val)

<220>
<221> MISC_FEATURE
<222> (55)..(55)
<223> Xaa at res. 55 = (Glu, Asp, Asn, Gly, Val, Pro or Lys)

<220>
<221> MISC_FEATURE
<222> (56)..(56)
<223> Xaa at res. 56 = (Thr, Ala, Val, Lys, Asp, Tyr, Ser, Gly, Ile
or
His)

<220>
<221> MISC_FEATURE
<222> (57)..(57)
<223> Xaa at res. 57 = (Val, Ala or Ile)

<220>
<221> MISC_FEATURE
<222> (58)..(58)
<223> Xaa at res. 58 = (Pro or Asp)

<220>
<221> MISC_FEATURE
<222> (59)..(59)
<223> Xaa at res. 59 = (Lys, Leu or Glu)

```

<220>
<221> MISC_FEATURE
<222> (60)..(60)
<223> Xaa at res. 60 = (Pro, Val or Ala)

<220>
<221> MISC_FEATURE
<222> (63)..(63)
<223> Xaa at res. 63 = (Ala or Val)

<220>
<221> MISC_FEATURE
<222> (65)..(65)
<223> Xaa at res. 65 = (Thr, Ala or Glu)

<220>
<221> MISC_FEATURE
<222> (66)..(66)
<223> Xaa at res. 66 = (Gln, Lys, Arg or Glu)

<220>
<221> MISC_FEATURE
<222> (67)..(67)
<223> Xaa at res. 67 = (Leu, Met or Val)

<220>
<221> MISC_FEATURE
<222> (68)..(68)
<223> Xaa at res. 68 = (Asn, Ser, Asp or Gly)

<220>
<221> MISC_FEATURE
<222> (69)..(69)
<223> Xaa at res. 69 = (Ala, Pro or Ser)

<220>
<221> MISC_FEATURE
<222> (70)..(70)
<223> Xaa at res. 70 = (Ile, Thr, Val or Leu)

<220>
<221> MISC_FEATURE
<222> (71)..(71)
<223> Xaa at res. 71 = (Ser, Ala or Pro)

<220>
<221> MISC_FEATURE
<222> (72)..(72)
<223> Xaa at res. 72 = (Val, Leu, Met or Ile)

<220>
<221> MISC_FEATURE
<222> (74)..(74)
<223> Xaa at res. 74 = (Tyr or Phe)

<220>
<221> MISC_FEATURE
<222> (75)..(75)


```

<223>  Xaa at res. 75 = (Phe, Tyr, Leu or His)

<220>
<221>  MISC_FEATURE
<222>  (76)..(76)
<223>  Xaa at res. 76 = (Asp, Asn or Leu)

<220>
<221>  MISC_FEATURE
<222>  (77)..(77)
<223>  Xaa at res. 77 = (Asp, Glu, Asn, Arg or Ser)

<220>
<221>  MISC_FEATURE
<222>  (78)..(78)
<223>  Xaa at res. 78 = (Ser, Gln, Asn, Tyr or Asp)

<220>
<221>  MISC_FEATURE
<222>  (79)..(79)
<223>  Xaa at res. 79 = (Ser, Asn, Asp, Glu or Lys)

<220>
<221>  MISC_FEATURE
<222>  (80)..(80)
<223>  Xaa at res. 80 = (Asn, Thr or Lys)

<220>
<221>  MISC_FEATURE
<222>  (82)..(82)
<223>  Xaa at res. 82 = (Ile, Val or Asn)

<220>
<221>  MISC_FEATURE
<222>  (84)..(84)
<223>  Xaa at res. 84 = (Lys or Arg)

<220>
<221>  MISC_FEATURE
<222>  (85)..(85)
<223>  Xaa at res. 85 = (Lys, Asn, Gln, His, Arg or Val)

<220>
<221>  MISC_FEATURE
<222>  (86)..(86)
<223>  Xaa at res. 86 = (Tyr, Glu or His)

<220>
<221>  MISC_FEATURE
<222>  (87)..(87)
<223>  Xaa at res. 87 = (Arg, Gln, Glu or Pro)

<220>
<221>  MISC_FEATURE
<222>  (88)..(88)
<223>  Xaa at res. 88 = (Asn, Glu, Trp or Asp)

<220>

```

```

<221> MISC_FEATURE
<222> (90)..(90)
<223> Xaa at res. 90 = (Val, Thr, Ala or Ile

<220>
<221> MISC_FEATURE
<222> (92)..(92)
<223> Xaa at res. 92 = (Arg, Lys, Val, Asp, Gln or Glu)

<220>
<221> MISC_FEATURE
<222> (93)..(93)
<223> Xaa at res. 93 = (Ala, Gly, Glu or Ser)

<220>
<221> MISC_FEATURE
<222> (95)..(95)
<223> Xaa at res. 95 = (Gly or Ala)

<220>
<221> MISC_FEATURE
<222> (97)..(97)
<223> Xaa at res. 97 = (His or Arg)

```

```

Leu Xaa Xaa Xaa Phe Xaa Xaa Xaa Gly Trp Xaa Xaa Xaa Xaa Xaa Xaa
1           5           10           15

```

```

Pro Xaa Xaa Xaa Xaa Ala Xaa Tyr Cys Xaa Gly Xaa Cys Xaa Xaa Pro
          20           25           30

```

```

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Asn His Ala Xaa Xaa Xaa Xaa Xaa
          35           40           45

```

```

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Cys Xaa Pro
          50           55           60

```

```

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Leu Xaa Xaa Xaa Xaa Xaa Xaa Xaa
65           70           75           80

```

```

Val Xaa Leu Xaa Xaa Xaa Xaa Xaa Met Xaa Val Xaa Xaa Cys Xaa Cys
          85           90           95

```

Xaa

SEQ ID NO:25

```

<220>
<223> generic sequence 2

```

<220>
<221> MISC_FEATURE
<222> (7)..(7)
<223> Xaa = (Tyr or Lys)

<220>
<221> MISC_FEATURE
<222> (8)..(8)
<223> Xaa = (Val or Ile)

<220>
<221> MISC_FEATURE
<222> (9)..(9)
<223> Xaa = (Ser, Asp or Glu)

<220>
<221> MISC_FEATURE
<222> (11)..(11)
<223> Xaa = (Arg, Gln, Ser, Lys or Ala)

<220>
<221> MISC_FEATURE
<222> (12)..(12)
<223> Xaa = (Asp or Glu)

<220>
<221> MISC_FEATURE
<222> (13)..(13)
<223> Xaa = (Leu, Val or Ile)

<220>
<221> MISC_FEATURE
<222> (16)..(16)
<223> Xaa = (Gln, Leu, Asp, His, Asn or Ser)

<220>
<221> MISC_FEATURE
<222> (17)..(17)
<223> Xaa = (Asp, Arg, Asn or Glu)

<220>
<221> MISC_FEATURE
<222> (18)..(18)
<223> Xaa = (Trp or Ser)

<220>
<221> MISC_FEATURE
<222> (19)..(19)
<223> Xaa = (Ile or Val)

<220>
<221> MISC_FEATURE
<222> (20)..(20)
<223> Xaa = (Ile or Val)

<220>
<221> MISC_FEATURE
<222> (21)..(21)

<223> Xaa = (Ala or Ser)

<220>
<221> MISC_FEATURE
<222> (23)..(23)
<223> Xaa = (Glu, Gln, Leu, Lys, Pro or Arg)

<220>
<221> MISC_FEATURE
<222> (24)..(24)
<223> Xaa = (Gly or Ser)

<220>
<221> MISC_FEATURE
<222> (25)..(25)
<223> Xaa = (Tyr or Phe)

<220>
<221> MISC_FEATURE
<222> (26)..(26)
<223> Xaa = (Ala, Ser, Asp, Met, His, Gln, Leu or Gly)

<220>
<221> MISC_FEATURE
<222> (28)..(28)
<223> Xaa = (Tyr, Asn or Phe)

<220>
<221> MISC_FEATURE
<222> (31)..(31)
<223> Xaa = (Glu, His, Tyr, Asp, Gln, Ala or Ser)

<220>
<221> MISC_FEATURE
<222> (33)..(33)
<223> Xaa = (Glu, Lys, Asp, Gln or Ala)

<220>
<221> MISC_FEATURE
<222> (35)..(35)
<223> Xaa = (Ala, Ser, Pro, Gln, Ile or Asn)

<220>
<221> MISC_FEATURE
<222> (36)..(36)
<223> Xaa = (Phe, Leu or Tyr)

<220>
<221> MISC_FEATURE
<222> (38)..(38)
<223> Xaa at res. 33 = (Leu, Val or Met)

<220>
<221> MISC_FEATURE
<222> (39)..(39)
<223> Xaa = (Asn, Asp, Ala, Thr or Pro)

<220>

<221> MISC_FEATURE
<222> (40)..(40)
<223> Xaa = (Ser, Asp, Glu, Leu, Ala or Lys)

<220>
<221> MISC_FEATURE
<222> (41)..(41)
<223> Xaa = (Tyr, Cys, His, Ser or Ile)

<220>
<221> MISC_FEATURE
<222> (42)..(42)
<223> Xaa = (Met, Phe, Gly or Leu)

<220>
<221> MISC_FEATURE
<222> (43)..(43)
<223> Xaa = (Asn, Ser or Lys)

<220>
<221> MISC_FEATURE
<222> (44)..(44)
<223> Xaa = (Ala, Ser, Gly or Pro)

<220>
<221> MISC_FEATURE
<222> (45)..(45)
<223> Xaa (Thr, Leu or Ser)

<220>
<221> MISC_FEATURE
<222> (49)..(49)
<223> Xaa = (Ile, Val or Thr)

<220>
<221> MISC_FEATURE
<222> (50)..(50)
<223> Xaa = (Val, Leu, Met or Ile)

<220>
<221> MISC_FEATURE
<222> (51)..(51)
<223> Xaa = (Gln or Arg)

<220>
<221> MISC_FEATURE
<222> (52)..(52)
<223> Xaa = (Thr, Ala or Ser)

<220>
<221> MISC_FEATURE
<222> (53)..(53)
<223> Xaa = (Leu or Ile)

<220>
<221> MISC_FEATURE
<222> (54)..(54)
<223> Xaa = (Val or Met)

<220>
<221> MISC_FEATURE
<222> (55)..(55)
<223> Xaa = (His, Asn or Arg)

<220>
<221> MISC_FEATURE
<222> (56)..(56)
<223> Xaa = (Phe, Leu, Asn, Ser, Ala or Val)

<220>
<221> MISC_FEATURE
<222> (57)..(57)
<223> Xaa = (Ile, Met, Asn, Ala, Val, Gly or Leu)

<220>
<221> MISC_FEATURE
<222> (58)..(58)
<223> Xaa = (Asn, Lys, Ala, Glu, Gly or Phe)

<220>
<221> MISC_FEATURE
<222> (59)..(59)
<223> Xaa = (Pro, Ser or Val)

<220>
<221> MISC_FEATURE
<222> (60)..(60)
<223> Xaa = (Glu, Asp, Asn, Gly, Val, Pro or Lys)

<220>
<221> MISC_FEATURE
<222> (61)..(61)
<223> Xaa = (Thr, Ala, Val, Lys, Asp, Tyr, Ser, Gly, Ile or His)

<220>
<221> MISC_FEATURE
<222> (62)..(62)
<223> Xaa = (Val, Ala or Ile)

<220>
<221> MISC_FEATURE
<222> (63)..(63)
<223> Xaa = (Pro or Asp)

<220>
<221> MISC_FEATURE
<222> (64)..(64)
<223> Xaa = (Lys, Leu or Glu)

<220>
<221> MISC_FEATURE
<222> (65)..(65)
<223> Xaa = (Pro, Val or Ala)

<220>

<221> MISC_FEATURE
<222> (68)..(68)
<223> Xaa = (Ala or Val)

<220>
<221> MISC_FEATURE
<222> (70)..(70)
<223> Xaa = (Thr, Ala or Glu)

<220>
<221> MISC_FEATURE
<222> (71)..(71)
<223> Xaa = (Gln, Lys, Arg or Glu)

<220>
<221> MISC_FEATURE
<222> (72)..(72)
<223> Xaa = (Leu, Met or Val)

<220>
<221> MISC_FEATURE
<222> (73)..(73)
<223> Xaa = (Asn, Ser, Asp or Gly)

<220>
<221> MISC_FEATURE
<222> (74)..(74)
<223> Xaa = (Ala, Pro or Ser)

<220>
<221> MISC_FEATURE
<222> (75)..(75)
<223> Xaa = (Ile, Thr, Val or Leu)

<220>
<221> MISC_FEATURE
<222> (76)..(76)
<223> Xaa = (Ser, Ala or Pro)

<220>
<221> MISC_FEATURE
<222> (77)..(77)
<223> Xaa = (Val, Leu, Met or Ile)

<220>
<221> MISC_FEATURE
<222> (79)..(79)
<223> Xaa = (Tyr or Phe)

<220>
<221> MISC_FEATURE
<222> (80)..(80)
<223> Xaa = (Phe, Tyr, Leu or His)

<220>
<221> MISC_FEATURE
<222> (81)..(81)
<223> Xaa = (Asp, Asn or Leu)

<220>
<221> MISC_FEATURE
<222> (82)..(82)
<223> Xaa = (Asp, Glu, Asn, Arg or Ser)

<220>
<221> MISC_FEATURE
<222> (83)..(83)
<223> Xaa = (Ser, Gln, Asn, Tyr or Asp)

<220>
<221> MISC_FEATURE
<222> (84)..(84)
<223> Xaa = (Ser, Asn, Asp, Glu or Lys)

<220>
<221> MISC_FEATURE
<222> (85)..(85)
<223> Xaa = (Asn, Thr or Lys)

<220>
<221> MISC_FEATURE
<222> (87)..(87)
<223> Xaa = (Ile, Val or Asn)

<220>
<221> MISC_FEATURE
<222> (89)..(89)
<223> Xaa = (Lys or Arg)

<220>
<221> MISC_FEATURE
<222> (90)..(90)
<223> Xaa = (Lys, Asn, Gln, His, Arg or Val)

<220>
<221> MISC_FEATURE
<222> (91)..(91)
<223> Xaa = (Tyr, Glu or His)

<220>
<221> MISC_FEATURE
<222> (92)..(92)
<223> Xaa = (Arg, Gln, Glu or Pro)

<220>
<221> MISC_FEATURE
<222> (93)..(93)
<223> Xaa = (Asn, Glu, Trp or Asp)

<220>
<221> MISC_FEATURE
<222> (95)..(95)
<223> Xaa = (Val, Thr, Ala or Ile)

<220>
<221> MISC_FEATURE

<222> (97)..(97)
 <223> Xaa = (Arg, Lys, Val, Asp, Gln or Glu)

<220>
 <221> MISC_FEATURE
 <222> (98)..(98)
 <223> Xaa = (Ala, Gly, Glu or Ser)

<220>
 <221> MISC_FEATURE
 <222> (100)..(100)
 <223> Xaa = (Gly or Ala)

<220>
 <221> MISC_FEATURE
 <222> (102)..(102)
 <223> Xaa = (His or Arg)

<400> 25

Cys	Xaa	Xaa	Xaa	Xaa	Leu	Xaa	Xaa	Xaa	Phe	Xaa	Xaa	Xaa	Gly	Trp	Xaa
1				5					10					15	

Xaa	Xaa	Xaa	Xaa	Xaa	Pro	Xaa	Xaa	Xaa	Xaa	Ala	Xaa	Tyr	Cys	Xaa	Gly
			20					25					30		

Xaa	Cys	Xaa	Xaa	Pro	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Asn	His	Ala
	35						40					45			

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
	50					55					60				

Xaa	Cys	Cys	Xaa	Pro	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Leu	Xaa	Xaa
65					70					75					80

Xaa	Xaa	Xaa	Xaa	Xaa	Val	Xaa	Leu	Xaa	Xaa	Xaa	Xaa	Xaa	Met	Xaa	Val
				85				90						95	

Xaa	Xaa	Cys	Xaa	Cys	Xaa
			100		

SEQ ID NO:26

<220>
 <223> generic sequence

<220>
 <221> MISC_FEATURE
 <222> (2)..(2)

<223> Xaa at res. 2 = (Lys, Arg, Ala or Gln)

<220>
<221> MISC_FEATURE
<222> (3)..(3)
<223> Xaa at res. 3 = (Lys, Arg or Met)

<220>
<221> MISC_FEATURE
<222> (4)..(4)
<223> Xaa at res. 4 = (His, Arg or Gln)

<220>
<221> MISC_FEATURE
<222> (5)..(5)
<223> Xaa at res. 5 = (Glu, Ser, His, Gly, Arg, Pro, Thr, or Tyr)

<400> 26

Cys Xaa Xaa Xaa Xaa
1 5

SEQ ID NO:27

<220>
<223> generic sequence 3

<220>
<221> MISC_FEATURE
<222> (1)..(1)
<223> Xaa at res. 1 = (Phe, Leu or Glu)

<220>
<221> MISC_FEATURE
<222> (2)..(2)
<223> Xaa at res. 2 = (Tyr, Phe, His, Arg, Thr, Lys, Gln, Val or Glu)

<220>
<221> MISC_FEATURE
<222> (3)..(3)
<223> Xaa at res. 3 = (Val, Ile, Leu or Asp)

<220>
<221> MISC_FEATURE
<222> (4)..(4)
<223> Xaa at res. 4 = (Ser, Asp, Glu, Asn or Phe)

<220>
<221> MISC_FEATURE
<222> (5)..(5)
<223> Xaa at res. 5 = (Phe or Glu)

<220>
<221> MISC_FEATURE
<222> (6)..(6)

<223> Xaa at res. 6 = (Arg, Gln, Lys, Ser, Glu, Ala or Asn)

<220>
<221> MISC_FEATURE
<222> (7)..(7)
<223> Xaa at res. 7 = (Asp, Glu, Leu, Ala or Gln)

<220>
<221> MISC_FEATURE
<222> (8)..(8)
<223> Xaa at res. 8 = (Leu, Val, Met, Ile or Phe)

<220>
<221> MISC_FEATURE
<222> (9)..(9)
<223> Xaa at res. 9 = (Gly, His or Lys)

<220>
<221> misc_feature
<222> (10)..(10)
<223> Xaa at res. 10 = (Trp or Met)

<220>
<221> MISC_FEATURE
<222> (11)..(11)
<223> Xaa at res. 11 = (Gln, Leu, His, Glu, Asn, Asp, Ser or Gly)

<220>
<221> MISC_FEATURE
<222> (12)..(12)
<223> Xaa at res. 12 = (Asp, Asn, Ser, Lys, Arg, Glu or His)

<220>
<221> MISC_FEATURE
<222> (13)..(13)
<223> Xaa at res. 13 = (Trp or Ser)

<220>
<221> MISC_FEATURE
<222> (14)..(14)
<223> Xaa at res. 14 = (Ile or Val)

<220>
<221> MISC_FEATURE
<222> (15)..(15)
<223> Xaa at res. 15 = (Ile or Val)

<220>
<221> MISC_FEATURE
<222> (16)..(16)
<223> Xaa at res. 16 = (Ala, Ser, Tyr or Trp)

<220>
<221> MISC_FEATURE
<222> (18)..(18)
<223> Xaa at res. 18 = (Glu, Lys, Gln, Met, Pro, Leu, Arg, His or Lys)

<220>
<221> MISC_FEATURE
<222> (19)..(19)
<223> Xaa at res. 19 = (Gly, Glu, Asp, Lys, Ser, Gln, Arg or Phe)

<220>
<221> MISC_FEATURE
<222> (20)..(20)
<223> Xaa at res. 20 = (Tyr or Phe)

<220>
<221> MISC_FEATURE
<222> (21)..(21)
<223> Xaa at res. 21 = (Ala, Ser, Gly, Met, Gln, His, Glu, Asp,
Leu,
Asn, Lys or Thr)

<220>
<221> MISC_FEATURE
<222> (22)..(22)
<223> Xaa at res. 22 = (Ala or Pro)

<220>
<221> MISC_FEATURE
<222> (23)..(23)
<223> Xaa at res. 23 = (Tyr, Phe, Asn, Ala or Arg)

<220>
<221> MISC_FEATURE
<222> (24)..(24)
<223> Xaa at res. 24 = (Tyr, His, Glu, Phe or Arg)

<220>
<221> MISC_FEATURE
<222> (26)..(26)
<223> Xaa at res. 26 = (Glu, Asp, Ala, Ser, Tyr, His, Lys, Arg, Gln
or
Gly)

<220>
<221> MISC_FEATURE
<222> (28)..(28)
<223> Xaa at res. 28 = (Glu, Asp, Leu, Val, Lys, Gly, Thr, Ala or
Gln)

<220>
<221> MISC_FEATURE
<222> (30)..(30)
<223> Xaa at res. 30 = (Ala, Ser, Ile, Asn, Pro, Glu, Asp, Phe, Gln
or
Leu)

<220>
<221> MISC_FEATURE
<222> (31)..(31)
<223> Xaa at res. 31 = (Phe, Tyr, Leu, Asn, Gly or Arg)

<220>

<221> MISC_FEATURE
<222> (32)..(32)
<223> Xaa at res. 32 = (Pro, Ser, Ala or Val)

<220>
<221> MISC_FEATURE
<222> (33)..(33)
<223> Xaa at res. 33 = (Leu, Met, Glu, Phe or Val)

<220>
<221> MISC_FEATURE
<222> (34)..(34)
<223> Xaa at res. 34 = (Asn, Asp, Thr, Gly, Ala, Arg, Leu or Pro)

<220>
<221> MISC_FEATURE
<222> (35)..(35)
<223> Xaa at res. 35 = (Ser, Ala, Glu, Asp, Thr, Leu, Lys, Gln or His)

<220>
<221> MISC_FEATURE
<222> (36)..(36)
<223> Xaa at res. 36 = (Tyr, His, Cys, Ile, Arg, Asp, Asn, Lys, Ser, Glu or Gly)

<220>
<221> MISC_FEATURE
<222> (37)..(37)
<223> Xaa at res. 37 = (Met, Leu, Phe, Val, Gly or Tyr)

<220>
<221> MISC_FEATURE
<222> (38)..(38)
<223> Xaa at res. 38 = (Asn, Glu, Thr, Pro, Lys, His, Gly, Met, Val or Arg)

<220>
<221> MISC_FEATURE
<222> (39)..(39)
<223> Xaa at res. 39 = (Ala, Ser, Gly, Pro or Phe)

<220>
<221> MISC_FEATURE
<222> (40)..(40)
<223> Xaa at res. 40 = (Thr, Ser, Leu, Pro, His or Met)

<220>
<221> MISC_FEATURE
<222> (41)..(41)
<223> Xaa at res. 41 = (Asn, Lys, Val, Thr or Gln)

<220>
<221> MISC_FEATURE
<222> (42)..(42)
<223> Xaa at res. 42 = (His, Tyr or Lys)

<220>
<221> MISC_FEATURE
<222> (43)..(43)
<223> Xaa at res. 43 = (Ala, Thr, Leu or Tyr)

<220>
<221> MISC_FEATURE
<222> (44)..(44)
<223> Xaa at res. 44 = (Ile, Thr, Val, Phe, Tyr, Met or Pro)

<220>
<221> MISC_FEATURE
<222> (45)..(45)
<223> Xaa at res. 45 = (Val, Leu, Met, Ile or His)

<220>
<221> MISC_FEATURE
<222> (46)..(46)
<223> Xaa at res. 46 = (Gln, Arg or Thr)

<220>
<221> MISC_FEATURE
<222> (47)..(47)
<223> Xaa at res. 47 = (Thr, Ser, Ala, Asn or His)

<220>
<221> MISC_FEATURE
<222> (48)..(48)
<223> Xaa at res. 48 = (Leu, Asn or Ile)

<220>
<221> MISC_FEATURE
<222> (49)..(49)
<223> Xaa at res. 49 = (Val, Met, Leu, Pro or Ile)

<220>
<221> MISC_FEATURE
<222> (50)..(50)
<223> Xaa at res. 50 = (His, Asn, Arg, Lys, Tyr or Gln)

<220>
<221> MISC_FEATURE
<222> (51)..(51)
<223> Xaa at res. 51 = (Phe, Leu, Ser, Asn, Met, Ala, Arg, Glu, Gly
or
Gln)

<220>
<221> MISC_FEATURE
<222> (52)..(52)
<223> Xaa at res. 52 = (Ile, Met, Leu, Val, Lys, Gln, Ala or Tyr)

<220>
<221> MISC_FEATURE
<222> (53)..(53)
<223> Xaa at res. 53 = (Asn, Phe, Lys, Glu, Asp, Ala, Gln, Gly, Leu
or

Val)

<220>
<221> MISC_FEATURE
<222> (54)..(54)
<223> Xaa at res. 54 = (Pro, Asn, Ser, Val or Asp)

<220>
<221> MISC_FEATURE
<222> (55)..(55)
<223> Xaa at res. 55 = (Glu, Asp, Asn, Lys, Arg, Ser, Gly, Thr,
Gln,
Pro or His)

<220>
<221> MISC_FEATURE
<222> (56)..(56)
<223> Xaa at res. 56 = (Thr, His, Tyr, Ala, Ile, Lys, Asp, Ser, Gly
or
Arg)

<220>
<221> MISC_FEATURE
<222> (57)..(57)
<223> Xaa at res. 57 = (Val, Ile, Thr, Ala, Leu or Ser)

<220>
<221> MISC_FEATURE
<222> (58)..(58)
<223> Xaa at res. 58 = (Pro, Gly, Ser, Asp or Ala)

<220>
<221> MISC_FEATURE
<222> (59)..(59)
<223> Xaa at res. 59 = (Lys, Leu, Pro, Ala, Ser, Glu, Arg or Gly)

<220>
<221> MISC_FEATURE
<222> (60)..(60)
<223> Xaa at res.: 60 = (Pro, Ala, Val, Thr or Ser)

<220>
<221> MISC_FEATURE
<222> (61)..(61)
<223> Xaa at res. 61 = (Cys, Val or Ser)

<220>
<221> MISC_FEATURE
<222> (63)..(63)
<223> Xaa at res. 63 = (Ala, Val or Thr)

<220>
<221> MISC_FEATURE
<222> (65)..(65)
<223> Xaa at res. 65 = (Thr, Ala, Glu, Val, Gly, Asp or Tyr)

<220>
<221> MISC_FEATURE

<222> (66)..(66)
<223> Xaa at res. 66 = (Gln, Lys, Glu, Arg or Val)

<220>
<221> MISC_FEATURE
<222> (67)..(67)
<223> Xaa at res. 67 = (Leu, Met, Thr or Tyr)

<220>
<221> MISC_FEATURE
<222> (68)..(68)
<223> Xaa at res. 68 = (Asn, Ser, Gly, Thr, Asp, Glu, Lys or Val)

<220>
<221> MISC_FEATURE
<222> (69)..(69)
<223> Xaa at res. 69 = (Ala, Pro, Gly or Ser)

<220>
<221> MISC_FEATURE
<222> (70)..(70)
<223> Xaa at res. 70 = (Ile, Thr, Leu or Val)

<220>
<221> MISC_FEATURE
<222> (71)..(71)
<223> Xaa at res. 71 = (Ser, Pro, Ala, Thr, Asn or Gly)

<220>
<221> MISC_FEATURE
<222> (72)..(72)
<223> Xaa at res. 2 = (Val, Ile, Leu or Met)

<220>
<221> MISC_FEATURE
<222> (74)..(74)
<223> Xaa at res. 74 = (Tyr, Phe, Arg, Thr, Tyr or Met)

<220>
<221> MISC_FEATURE
<222> (75)..(75)
<223> Xaa at res. 75 = (Phe, Tyr, His, Leu, Ile, Lys, Gln or Val)

<220>
<221> MISC_FEATURE
<222> (76)..(76)
<223> Xaa at res. 76 = (Asp, Leu, Asn or Glu)

<220>
<221> MISC_FEATURE
<222> (77)..(77)
<223> Xaa at res. 77 = (Asp, Ser, Arg, Asn, Glu, Ala, Lys, Gly or Pro)

<220>
<221> MISC_FEATURE
<222> (78)..(78)


```

<223> Xaa at res. 78 = (Ser, Asn, Asp, Tyr, Ala, Gly, Gln, Met,
Glu,
      Asn or Lys)

<220>
<221> MISC_FEATURE
<222> (79)..(79)
<223> Xaa at res. 79 = (Ser, Asn, Glu, Asp, Val, Lys, Gly, Gln or
Arg)

<220>
<221> MISC_FEATURE
<222> (80)..(80)
<223> Xaa at res. 80 = (Asn, Lys, Thr, Pro, Val, Ile, Arg; Ser or
Gln)

<220>
<221> MISC_FEATURE
<222> (81)..(81)
<223> Xaa at res. 81 = (Val, Ile, Thr or Ala)

<220>
<221> MISC_FEATURE
<222> (82)..(82)
<223> Xaa at res. 82 = (Ile, Asn, Val, Leu, Tyr, Asp or Ala)

<220>
<221> misc_feature
<222> (83)..(83)
<223> Xaa at res. 83 = (Leu, Tyr, Lys, or Ile)

<220>
<221> MISC_FEATURE
<222> (84)..(84)
<223> Xaa at res. 84 = (Lys, Arg, Asn, Tyr, Phe, Thr, Glu or Gly)

<220>
<221> MISC_FEATURE
<222> (85)..(85)
<223> Xaa at res. 85 = (Lys, Arg, His, Gln, Asn, Glu or Val)

<220>
<221> MISC_FEATURE
<222> (86)..(86)
<223> Xaa at res. 86 = (Tyr, His, Glu or Ile)

<220>
<221> MISC_FEATURE
<222> (87)..(87)
<223> Xaa at res. 87 = (Arg, Glu, Gln, Pro or Lys)

<220>
<221> MISC_FEATURE
<222> (88)..(88)
<223> Xaa at res. 88 = (Asn, Asp, Ala, Glu, Gly or Lys)

<220>
<221> MISC_FEATURE

```

```

<222> (89)..(89)
<223> Xaa at res. 89 = (Met or Ala)

<220>
<221> MISC_FEATURE
<222> (90)..(90)
<223> Xaa at res. 90 = (Val, Ile, Ala, Thr, Ser or Lys)

<220>
<221> MISC_FEATURE
<222> (91)..(91)
<223> Xaa at res. 91 = (Val or Ala)

<220>
<221> MISC_FEATURE
<222> (92)..(92)
<223> Xaa at res. 92 = (Arg, Lys, Gln, Asp, Glu, Val, Ala, Ser or
Thr)

<220>
<221> MISC_FEATURE
<222> (93)..(93)
<223> Xaa at res. 93 = (Ala, Ser, Glu, Gly, Arg or Thr)

<220>
<221> MISC_FEATURE
<222> (95)..(95)
<223> Xaa at res. 95 = (Gly, Ala or Thr)

<220>
<221> MISC_FEATURE
<222> (97)..(97)
<223> Xaa at res. 97 = (His, Arg, Gly, Leu or Ser)

<400> 27

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
1 5 10 15

Pro Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Gly Xaa Cys Xaa Xaa Xaa
20 25 30

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
35 40 45

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Pro
50 55 60

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
65 70 75 80

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Cys
85 90 95

```

Xaa

SEQ ID NO:28

<220>
<223> generic sequence 4

<220>
<221> MISC_FEATURE
<222> (6)..(6)
<223> Xaa = (Phe, Leu or Glu)

<220>
<221> MISC_FEATURE
<222> (7)..(7)
<223> Xaa = (Tyr, Phe, His, Arg, Thr, Lys, Gln, Val or Glu)

<220>
<221> MISC_FEATURE
<222> (8)..(8)
<223> Xaa = (Val, Ile, Leu or Asp)

<220>
<221> MISC_FEATURE
<222> (9)..(9)
<223> Xaa = (Ser, Asp, Glu, Asn or Phe)

<220>
<221> MISC_FEATURE
<222> (10)..(10)
<223> Xaa = (Phe or Glu)

<220>
<221> MISC_FEATURE
<222> (11)..(11)
<223> Xaa = (Arg, Gln, Lys, Ser, Glu, Ala or Asn)

<220>
<221> MISC_FEATURE
<222> (12)..(12)
<223> Xaa = (Asp, Glu, Leu, Ala or Gln)

<220>
<221> MISC_FEATURE
<222> (13)..(13)
<223> Xaa = (Leu, Val, Met, Ile or Phe)

<220>
<221> MISC_FEATURE
<222> (14)..(14)
<223> Xaa = (Gly, His or Lys)

<220>
<221> misc_feature
<222> (15)..(15)

<223> Xaa = (Trp or Met)

<220>
<221> MISC_FEATURE
<222> (16)..(16)
<223> Xaa = (Gln, Leu, His, Glu, Asn, Asp, Ser or Gly)

<220>
<221> MISC_FEATURE
<222> (17)..(17)
<223> Xaa = (Asp, Asn, Ser, Lys, Arg, Glu or His)

<220>
<221> MISC_FEATURE
<222> (18)..(18)
<223> Xaa = (Trp or Ser)

<220>
<221> MISC_FEATURE
<222> (19)..(19)
<223> Xaa = (Ile or Val)

<220>
<221> MISC_FEATURE
<222> (20)..(20)
<223> Xaa = (Ile or Val)

<220>
<221> MISC_FEATURE
<222> (21)..(21)
<223> Xaa = (Ala, Ser, Tyr or Trp)

<220>
<221> MISC_FEATURE
<222> (23)..(23)
<223> Xaa = (Glu, Lys, Gln, Met, Pro, Leu, Arg, His or Lys)

<220>
<221> MISC_FEATURE
<222> (24)..(24)
<223> Xaa = (Gly, Glu, Asp, Lys, Ser, Gln, Arg or Phe)

<220>
<221> MISC_FEATURE
<222> (25)..(25)
<223> Xaa = (Tyr or Phe)

<220>
<221> MISC_FEATURE
<222> (26)..(26)
<223> Xaa Ala, Ser, Gly, Met, Gln, His, Glu, Asp, Leu,
Asn, Lys or Thr)

<220>
<221> MISC_FEATURE
<222> (27)..(27)
<223> Xaa = (Ala or Pro)

<220>
<221> MISC_FEATURE
<222> (28)..(28)
<223> Xaa = (Tyr, Phe, Asn, Ala or Arg)

<220>
<221> MISC_FEATURE
<222> (29)..(29)
<223> Xaa = (Tyr, His, Glu, Phe or Arg)

<220>
<221> MISC_FEATURE
<222> (31)..(31)
<223> Xaa = (Glu, Asp, Ala, Ser, Tyr, His, Lys, Arg, Gln or Gly)

<220>
<221> MISC_FEATURE
<222> (33)..(33)
<223> Xaa = (Glu, Asp, Leu, Val, Lys, Gly, Thr, Ala or Gln)

<220>
<221> MISC_FEATURE
<222> (35)..(35)
<223> Xaa = (Ala, Ser, Ile, Asn, Pro, Glu, Asp, Phe, Gln or Leu)

<220>
<221> MISC_FEATURE
<222> (36)..(36)
<223> Xaa = (Phe, Tyr, Leu, Asn, Gly or Arg)

<220>
<221> MISC_FEATURE
<222> (37)..(37)
<223> Xaa = (Pro, Ser, Ala or Val)

<220>
<221> MISC_FEATURE
<222> (38)..(38)
<223> Xaa = (Leu, Met, Glu, Phe or Val)

<220>
<221> MISC_FEATURE
<222> (39)..(39)
<223> Xaa = (Asn, Asp, Thr, Gly, Ala, Arg, Leu or Pro)

<220>
<221> MISC_FEATURE
<222> (40)..(40)
<223> Xaa = (Ser, Ala, Glu, Asp, Thr, Leu, Lys, Gln or His)

<220>
<221> MISC_FEATURE
<222> (41)..(41)
<223> Xaa = (Tyr, His, Cys, Ile, Arg, Asp, Asn, Lys, Ser, Glu or Gly)

```

<220>
<221> MISC_FEATURE
<222> (42)..(42)
<223> Xaa = (Met, Leu, Phe, Val, Gly or Tyr)

<220>
<221> MISC_FEATURE
<222> (43)..(43)
<223> Xaa = (Asn, Glu, Thr, Pro, Lys, His, Gly, Met, Val or
Arg)

<220>
<221> MISC_FEATURE
<222> (44)..(44)
<223> Xaa = (Ala, Ser, Gly, Pro or Phe)

<220>
<221> MISC_FEATURE
<222> (45)..(45)
<223> Xaa = (Thr, Ser, Leu, Pro, His or Met)

<220>
<221> MISC_FEATURE
<222> (46)..(46)
<223> Xaa = (Asn, Lys, Val, Thr or Gln)

<220>
<221> MISC_FEATURE
<222> (47)..(47)
<223> Xaa = (His, Tyr or Lys)

<220>
<221> MISC_FEATURE
<222> (48)..(48)
<223> Xaa = (Ala, Thr, Leu or Tyr)

<220>
<221> MISC_FEATURE
<222> (49)..(49)
<223> Xaa = (Ile, Thr, Val, Phe, Tyr, Met or Pro)

<220>
<221> MISC_FEATURE
<222> (50)..(50)
<223> Xaa = (Val, Leu, Met, Ile or His)

<220>
<221> MISC_FEATURE
<222> (51)..(51)
<223> Xaa = (Gln, Arg or Thr)

<220>
<221> MISC_FEATURE
<222> (52)..(52)
<223> Xaa = (Thr, Ser, Ala, Asn or His)

<220>
<221> MISC_FEATURE

```

<222> (53)..(53)
<223> Xaa = (Leu, Asn or Ile)

<220>
<221> MISC_FEATURE
<222> (54)..(54)
<223> Xaa = (Val, Met, Leu, Pro or Ile)

<220>
<221> MISC_FEATURE
<222> (55)..(55)
<223> Xaa = (His, Asn, Arg, Lys, Tyr or Gln)

<220>
<221> MISC_FEATURE
<222> (56)..(56)
<223> Xaa = (Phe, Leu, Ser, Asn, Met, Ala, Arg, Glu, Gly or Gln)

<220>
<221> MISC_FEATURE
<222> (57)..(57)
<223> Xaa = (Ile, Met, Leu, Val, Lys, Gln, Ala or Tyr)

<220>
<221> MISC_FEATURE
<222> (58)..(58)
<223> Xaa = (Asn, Phe, Lys, Glu, Asp, Ala, Gln, Gly, Leu or Val)

<220>
<221> MISC_FEATURE
<222> (59)..(59)
<223> Xaa = (Pro, Asn, Ser, Val or Asp)

<220>
<221> MISC_FEATURE
<222> (60)..(60)
<223> Xaa = (Glu, Asp, Asn, Lys, Arg, Ser, Gly, Thr, Gln, Pro or His)

<220>
<221> MISC_FEATURE
<222> (61)..(61)
<223> Xaa = (Thr, His, Tyr, Ala, Ile, Lys, Asp, Ser, Gly or Arg)

<220>
<221> MISC_FEATURE
<222> (62)..(62)
<223> Xaa = (Val, Ile, Thr, Ala, Leu or Ser)

<220>
<221> MISC_FEATURE
<222> (63)..(63)
<223> Xaa = (Pro, Gly, Ser, Asp or Ala)

<220>

<221> MISC_FEATURE
<222> (64)..(64)
<223> Xaa = (Lys, Leu, Pro, Ala, Ser, Glu, Arg or Gly)

<220>
<221> MISC_FEATURE
<222> (65)..(65)
<223> Xaa = (Pro, Ala, Val, Thr or Ser)

<220>
<221> MISC_FEATURE
<222> (66)..(66)
<223> Xaa = (Cys, Val or Ser)

<220>
<221> MISC_FEATURE
<222> (68)..(68)
<223> Xaa = (Ala, Val or Thr)

<220>
<221> MISC_FEATURE
<222> (70)..(70)
<223> Xaa = (Thr, Ala, Glu, Val, Gly, Asp or Tyr)

<220>
<221> MISC_FEATURE
<222> (71)..(71)
<223> Xaa = (Gln, Lys, Glu, Arg or Val)

<220>
<221> MISC_FEATURE
<222> (72)..(72)
<223> Xaa = (Leu, Met, Thr or Tyr)

<220>
<221> MISC_FEATURE
<222> (73)..(73)
<223> Xaa = (Asn, Ser, Gly, Thr, Asp, Glu, Lys or Val)

<220>
<221> MISC_FEATURE
<222> (74)..(74)
<223> Xaa = (Ala, Pro, Gly or Ser)

<220>
<221> MISC_FEATURE
<222> (75)..(75)
<223> Xaa = (Ile, Thr, Leu or Val)

<220>
<221> MISC_FEATURE
<222> (76)..(76)
<223> Xaa = (Ser, Pro, Ala, Thr, Asn or Gly)

<220>
<221> MISC_FEATURE
<222> (77)..(77)
<223> Xaa = (Val, Ile, Leu or Met)

<220>
<221> MISC_FEATURE
<222> (79)..(79)
<223> Xaa = (Tyr, Phe, Arg, Thr, Tyr or Met)
<220>
<221> MISC_FEATURE
<222> (80)..(80)
<223> Xaa = (Phe, Tyr, His, Leu, Ile, Lys, Gln or Val)

<220>
<221> MISC_FEATURE
<222> (81)..(81)
<223> Xaa = (Asp, Leu, Asn or Glu)

<220>
<221> MISC_FEATURE
<222> (82)..(82)
<223> Xaa = (Asp, Ser, Arg, Asn, Glu, Ala, Lys, Gly or Pro)

<220>
<221> MISC_FEATURE
<222> (83)..(83)
<223> Xaa = (Ser, Asn, Asp, Tyr, Ala, Gly, Gln, Met, Glu,
Asn or Lys)

<220>
<221> MISC_FEATURE
<222> (84)..(84)
<223> Xaa = (Ser, Asn, Glu, Asp, Val, Lys, Gly, Gln or Arg)

<220>
<221> MISC_FEATURE
<222> (85)..(85)
<223> Xaa = (Asn, Lys, Thr, Pro, Val, Ile, Arg; Ser or Gln)

<220>
<221> MISC_FEATURE
<222> (86)..(86)
<223> Xaa = (Val, Ile, Thr or Ala)

<220>
<221> MISC_FEATURE
<222> (87)..(87)
<223> Xaa = (Ile, Asn, Val, Leu, Tyr, Asp or Ala)

<220>
<221> misc_feature
<222> (88)..(88)
<223> Xaa = (Leu, Tyr, Lys, or Ile)

<220>
<221> MISC_FEATURE
<222> (89)..(89)
<223> Xaa = (Lys, Arg, Asn, Tyr, Phe, Thr, Glu or Gly)

<220>
<221> MISC_FEATURE

```

<222> (90)..(90)
<223> Xaa = (Lys, Arg, His, Gln, Asn, Glu or Val)

<220>
<221> MISC_FEATURE
<222> (91)..(91)
<223> Xaa = (Tyr, His, Glu or Ile)

<220>
<221> MISC_FEATURE
<222> (92)..(92)
<223> Xaa = (Arg, Glu, Gln, Pro or Lys)

<220>
<221> MISC_FEATURE
<222> (93)..(93)
<223> Xaa = (Asn, Asp, Ala, Glu, Gly or Lys)

<220>
<221> MISC_FEATURE
<222> (94)..(94)
<223> Xaa = (Met or Ala)

<220>
<221> MISC_FEATURE
<222> (95)..(95)
<223> Xaa = (Val, Ile, Ala, Thr, Ser or Lys)

<220>
<221> MISC_FEATURE
<222> (96)..(96)
<223> Xaa = (Val or Ala)

<220>
<221> MISC_FEATURE
<222> (97)..(97)
<223> Xaa = (Arg, Lys, Gln, Asp, Glu, Val, Ala, Ser or Thr)

<220>
<221> MISC_FEATURE
<222> (98)..(98)
<223> Xaa = (Ala, Ser, Glu, Gly, Arg or Thr)

<220>
<221> MISC_FEATURE
<222> (100)..(100)
<223> Xaa = (Gly, Ala or Thr)

<220>
<221> MISC_FEATURE
<222> (102)..(102)
<223> Xaa = (His, Arg, Gly, Leu or Ser)

<400> 28

Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
1      5      10      15

```

```

Xaa Xaa Xaa Xaa Xaa Pro Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Gly
      20                      25                      30
Xaa Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
      35                      40                      45
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
      50                      55                      60
Xaa Xaa Cys Xaa Pro Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Leu Xaa Xaa
      65                      70                      75                      80
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
      85                      90                      95
Xaa Xaa Cys Xaa Cys Xaa
      100

```

SEQ ID NO:29

```

<220>
<223>  generic sequence 5

<220>
<221>  MISC_FEATURE
<222>  (2)..(2)
<223>  Xaa at res. 2 = (Lys or Arg)

<220>
<221>  MISC_FEATURE
<222>  (3)..(3)
<223>  Xaa at res. 3 = (Lys or Arg)

<220>
<221>  MISC_FEATURE
<222>  (11)..(11)
<223>  Xaa at res. 11 = (Arg or Gln)

<220>
<221>  MISC_FEATURE
<222>  (16)..(16)
<223>  Xaa at res. 16 = (Gln or Leu)

<220>
<221>  MISC_FEATURE
<222>  (19)..(19)
<223>  Xaa. at res. 19 = (Ile or Val)

<220>

```

```
<221> MISC_FEATURE
<222> (23)..(23)
<223> Xaa at res. 23 = (Glu or Gln)

<220>
<221> MISC_FEATURE
<222> (26)..(26)
<223> Xaa at res. 26 = (Ala or Ser)

<220>
<221> MISC_FEATURE
<222> (35)..(35)
<223> Xaa at res. 35 = (Ala or Ser)

<220>
<221> misc_feature
<222> (39)..(39)
<223> Xaa can be any naturally occurring amino acid

<220>
<221> MISC_FEATURE
<222> (41)..(41)
<223> Xaa at res. 41 = (Tyr or Cys)

<220>
<221> MISC_FEATURE
<222> (50)..(50)
<223> Xaa at res. 50 = (Val or Leu)

<220>
<221> MISC_FEATURE
<222> (52)..(52)
<223> Xaa at res. 52 = (Ser or Thr)

<220>
<221> MISC_FEATURE
<222> (56)..(56)
<223> Xaa at res. 56 = (Phe or Leu)

<220>
<221> MISC_FEATURE
<222> (57)..(57)
<223> Xaa at res. 57 = (Ile or Met)

<220>
<221> MISC_FEATURE
<222> (58)..(58)
<223> Xaa at res. 58 = (Asn or Lys)

<220>
<221> MISC_FEATURE
<222> (60)..(60)
<223> Xaa at res. 60 = (Glu, Asp or Asn)

<220>
<221> MISC_FEATURE
<222> (61)..(61)
<223> Xaa at res. 61 = (Thr, Ala or Val)
```

<220>
<221> MISC_FEATURE
<222> (65)..(65)
<223> Xaa at res. 65 = (Pro or Ala)

<220>
<221> MISC_FEATURE
<222> (71)..(71)
<223> Xaa at res. 71 = (Gln or Lys)

<220>
<221> MISC_FEATURE
<222> (73)..(73)
<223> Xaa at res. 73 = (Asn or Ser)

<220>
<221> MISC_FEATURE
<222> (75)..(75)
<223> Xaa at res. 75 = (Ile or Thr)

<220>
<221> MISC_FEATURE
<222> (80)..(80)
<223> Xaa at res. 80 = (Phe or Tyr)

<220>
<221> MISC_FEATURE
<222> (82)..(82)
<223> Xaa at res. 82 = (Asp or Ser)

<220>
<221> MISC_FEATURE
<222> (84)..(84)
<223> Xaa at res. 84 = (Ser or Asn)

<220>
<221> MISC_FEATURE
<222> (89)..(89)
<223> Xaa at res. 89 = (Lys or Arg)

<220>
<221> MISC_FEATURE
<222> (91)..(91)
<223> Xaa at res. 91 = (Tyr or His)

<220>
<221> misc_feature
<222> (96)..(96)
<223> Xaa can be any naturally occurring amino acid

<220>
<221> MISC_FEATURE
<222> (97)..(97)
<223> Xaa at res. 97 = (Arg or Lys)

<400> 29

Cys Xaa Xaa His Glu Leu Tyr Val Ser Phe Xaa Asp Leu Gly Trp Xaa
1 5 10 15

Asp Trp Xaa Ile Ala Pro Xaa Gly Tyr Xaa Ala Tyr Tyr Cys Glu Gly
20 25 30

Glu Cys Xaa Phe Pro Leu Xaa Ser Xaa Met Asn Ala Thr Asn His Ala
35 40 45

Ile Xaa Gln Xaa Leu Val His Xaa Xaa Xaa Pro Xaa Xaa Val Pro Lys
50 55 60

Xaa Cys Cys Ala Pro Thr Xaa Leu Xaa Ala Xaa Ser Val Leu Tyr Xaa
65 70 75 80

Asp Xaa Ser Xaa Asn Val Ile Leu Xaa Lys Lys Arg Asn Met Val Xaa
85 90 95

Ala Cys Gly Cys His
100

SEQ ID NO:30

<220>

<223> proteolytic site

<220>

<221> misc_feature

<222> (2)..(3)

<223> Xaa can be any naturally occurring amino acid

<400> 30

Arg Xaa Xaa Arg
1

SEQ ID NO:31

Gly Gly Pro Pro
1

All cited references (including scientific publications, abstracts, etc.),
patents, patent application publications are hereby incorporated herein by reference.